

Sally 11, 1977
UNDER THE
FOR L
EQU UNDER NO. 6735-170

TIDE

TOXAPHENE-SULFUR 20-40 DUST

AN AGRICULTURAL INSECTICIDE - FUNGICIDE

ACTIVE INGREDIENTS:

Toxaphene 20.0%
Sulfur 40.0%

INERT INGREDIENTS

Technical Chlorinated Camphene (Chlorine Content 67% to 69%) 100.0%

DIRECTIONS FOR USE

Apply the specified amounts of this product uniformly to the foliage when air is calm, using ground equipment or aircraft. Apply at the first sign of infestation and infection and repeat at 7 to 10 day intervals, depending upon the severity of the disease and insect conditions, but observe no limit on the number of applications.

For the control of disease, persistent after the prescribed time limits have expired, an effective fungicide should be used alone in a manner which will not leave residue in excess of established tolerances.

WARNING

Keep Out of Reach of Children

HARMFUL IF SWALLOWED.

AVOID BREATHING DUST, CONTACT WITH EYES, SKIN AND CLOTHING.

WASH THOROUGHLY AFTER HANDLING.

DO NOT CONTAMINATE FOOD AND FEED

OBSERVE THE FOLLOWING USE PRECAUTIONS

Observe use limitations given on the label for specific crops.

In order that pesticide residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals, or onto adjacent crops, feed or pasture crops. The grower is responsible for residues on his crop, as well as for damage caused by drift from his property to that of others.

Highly toxic to fish and wildlife. Fish feeding on treated areas may be killed. Do not contaminate any body of water. Do not apply to any area not specified on the label.

This product is toxic to bees and should not be applied when bees are actively visiting the bloom.

Do not use this product on food or forage crops to be fed to dairy animals or to livestock from the time of application.

Do not use Toxaphene on citrus, or on other fruit crops, or on grape, currant, or other berry crops, or on any other fruit crop, unless specifically labeled for such use.

Sulfur may cause damage to plants under conditions of high temperatures and high humidity, or other similar conditions. Do not apply sulfur to sensitive crops when maximum temperature exceeds 90 degrees Fahrenheit.

Consult State Agricultural Experiment Stations or State Agricultural Extension Service for additional information. The time of day, number and rate of applications, weather will vary with local conditions.

Not for use or storage in or around the home environment.

Crush empty containers and bury at least 18 inches deep.

RECOMMENDATIONS

ALFALFA: Armyworms, Flea Beetle, Leafhoppers, Lygus Bug, Thrips. Apply at rate of 10 to 15 lb. per acre. Do not feed treated forage to dairy animals or animals being finished for slaughter.

BEANS: Armyworms, Bean Leaf Roller and Leafhopper, Climbing Cutworms, apply only in ~~green~~ stage. Corn Earworm, Cowpea Curculio, Lygus, Powdery Mildew, Rust, Salt-marsh Caterpillar, Thrips. Use 10 lb. per acre. Do not apply to green or snap beans within 7 days of harvest. Do not feed treated forage to dairy animals or animals being finished for slaughter.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER: Cabbage Looper, Corn Earworm, Flea Beetle, Imported Cabbageworm, Cutworms (clubfoot and sulfur-feeding), Fall Armyworm, Spiny-legged Leaf Miner, Thrips. Use 10 lb. per acre. Flea Beetles. Use 15 lb. per acre. Do not apply after edible parts start to form or within 30 days of harvest. On cabbage do not apply after heads begin to form.

COTTON: Overwintered Boll Weevil, Flea Beetles, Leafhoppers, Lygus bugs, or other insects. Apply 10 lb. per acre. Make 2 to 4 applications at 7 to 10 day intervals. Fall Armyworm, Leafminer, other Moths. Use 10 to 15 lb. per acre. Boll Weevil. Use 15 to 20 lb. per acre. Apply at 4 to 5 day intervals until control is obtained. Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season.

PEPPERS: Armyworm, Blister Beetle, Cabbage Root Fly, Colorado Potato Beetle, Colorado Potato Beetle, Flea Beetle, Hornworm, Serpentine, Leaf Miner. Use 10 to 15 lb. per acre. Fruitworm, Pepper Weevil. Use 15 lb. per acre. Do not apply within 5 days of harvest.

TOMATOES: Armyworms, Blister Beetle, Cabbage Root Fly, Colorado Potato Beetle, Flea Beetle, Lygus, Cutworm (sulfur-feeding), Flea Beetle, Leafworm, Root-Minor, Spiny-legged Leaf Miner, Fruitworm, Tomato Rootworm. Use 10 to 20 lb. per acre. Do not apply within 3 days of harvest.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with the directions under normal conditions of use but neither this warranty nor any other warranty or merchantability or fitness for a particular purpose express or implied, extends to the use of this product contrary to the instructions, or under abnormal conditions, or under conditions not reasonably foreseeable, and buyer assumes the risk of any such use.

USDA Reg. No. 6735-170

GPS 3411 IM 3692 D 11

A PRODUCT OF
TIDE PRODUCTS, INC.

EDINBURG, TEXAS 78539

50 LB. NET