

June 25, 1971
FOR L...
ED UNDER 0735-189

TIDE

SEVIN* - ZINEB 10-6.5

AGRICULTURAL INSECTICIDE - FUNGICIDE

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-Methylcarbamate) 10%
Zineb (Zinc Ethylene Bisdithiocarbamate) 6.5%
(Zinc expressed as Metallic — 1.42%)

INERT INGREDIENTS

83.5%
100.0%

*Trade-Mark of UCC for 1-naphthyl N-methylcarbamate.

RECOMMENDATIONS

BEANS (FIELD, LIMA, SNAP) ^{bean} Insect control -- Mexican Bean Beetle, use 10 pounds per acre; Bean Leaf Beetle, Cucumber Beetle, Flea Beetle, Japanese Beetle, Leafhopper, Velvet Bean Caterpillar, Armyworms, Corn Earworms, Stink Bugs and Tarnished Plant Bug use 20 pounds per acre. Disease control -- Anthracnose, Downy Mildew and Rust - Use 20 pounds per acre. Begin use when disease threatens or after first bloom appears. Do not apply within 7 days of harvest.

CUCUMBER, MELONS, PUMPKIN, AND SQUASH: Insect control -- Pickleworm, Melonworm, Cucumber Beetle, Flea Beetle, Leafhopper and Squash Bugs. Disease control -- Alternaria Blight, Angular Leaf Spot, Anthracnose, Blossom Blight, Downy Mildew and Gummy Stem Blight - Use 10 pounds per acre, begin treatment for disease control when disease appears. To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days. Sevin is injurious to watermelons in Florida. No time limitations.

EGG PLANT, TOMATO AND PEPPER: Insect control -- Colorado Potato Beetle, Flea Beetle and Leafhopper -- Use 20 pounds per acre; Fall Armyworms, Lace Bugs, Tomato Fruitworms, Tomato Hornworms, Tarnished Plant Bugs and Stink Bugs, use 30 pounds per acre. Disease control -- Egg Plant -- Alternaria Leaf Spots (Early Blight), Anthracnose, Downy Mildew, Peppers -- Anthracnose, Bacterial Leaf Spot, Downy Mildew, Cercospora Leaf Spot and Phytophthora. Tomatoes -- Anthracnose, Early and Late Blights, Septoria Leaf Spot and Southern Blight. Use 20 to 30 pounds per acre when disease threatens. No time limitations.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE AND CAULIFLOWER: Insect control -- Flea Beetles, Harlequin Bug use 20 pounds per acre. Disease control -- Downy Mildew and Leaf Spots; use 20 pounds per acre. Do not apply within 7 days of harvest.

PEANUTS: Insect control -- Blister Beetle, Mexican Bean Beetle, Cucumber Beetle, Leafhoppers and Thrips use 15 pounds per acre. Disease control -- Cercospora Leaf Spot. Use 15 pounds per acre. Begin treatment when first spots appear on lower leaves. To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days. No time limitations. Do not feed treated top to livestock.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty or merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

CAUTION

- KEEP OUT OF REACH OF CHILDREN AND ANIMALS
- HARMFUL IF SWALLOWED OR INHALED.
- DO NOT TAKE INTERNALLY.
- DO NOT BREATHE DUST.
- AVOID PROLONGED OR REPEATED BREATHING OF DUST.
- SKIN CONTACT MAY BE HARMFUL. AVOID CONTACT.
- WEAR REGULAR LONG SLEEVED WORK CLOTHING. CHANGE TO CLEAN CLOTHING DAILY.
- WASH HANDS BEFORE EATING. TAKE SHOWER OR BATH AFTER WORK.
- NOTE FOR PHYSICIANS: CARBARYL IS A MODERATE REVERSIBLE, CHOLINESTERASE INHIBITOR. ATROPINE IS ANTIDOTAL.

OBSERVE THE FOLLOWING USE PRECAUTIONS

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified.

This product is highly toxic to bees exposed to direct treatment or residue on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

Apply when air is calm, using ground equipment, aircraft, or hand sprayer. Make thorough coverage of plants with a uniform deposit of dust. Apply at the first sign of infestation and repeat at regular intervals when needed to maintain control.

DO NOT REUSE EMPTY CONTAINER. DESTROY IT BY BURYING WITH WASTE OR BURNING. STAY AWAY FROM SMOKE OR FUMES.

A PRODUCT OF
TIDE PRODUCTS, INC.
EDINBURG, TEXAS 78539