

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

SEVIN* -PARATHION 10-1

AGRICULTURAL INSECTICIDE

ACTIVE INGREDIENTS:
Parathion (O,O-diethyl O-p-nitrophenyl thiophosphate)
Carbaryl (1-Naphthyl N-methylcarbamate)

INERT INGREDIENTS

*Trade-mark of UCC for 1-Naphthyl N-Methylcarbamate

DIRECTIONS FOR USE

This product is an insecticide dust formulation for use by the commercial grower or commercial applicator. Apply when air is calm. To obtain most effective results, apply with a ground blower or by airplane. Make thorough coverage of plants with a uniform deposit of dust.

RECOMMENDATIONS

COTTON

- For control of early season Thrips and Cotton Fleahopper. Apply 8 to 15 lbs per acre when insects first appear.
- For control of Mid and Late Season Boll Weevil, Cotton Leafworm, Lygus Bugs, and Cotton Leafhopper. Apply 15 to 25 lbs per acre at 5-7 day intervals when insects first appear.
- For control of Pink Bollworm. Apply 22-25 lbs per acre at 5-7 day intervals.
- On cotton do not apply within 5 days of hand picking.

BEANS: For control of Mexican Bean Beetle apply 10-15 lbs per acre. Leafhoppers apply 10-15 lbs per acre and Tarnished Plant Bug, apply 12-18 lbs per acre at 7-10 day intervals. Do not apply within 7 days of harvest.

POTATOES: For control of Colorado Potato Beetle, Potato Leafhopper, Six Spotted Leafhopper and Potato Flea Beetle, apply 7-10 lbs per acre, and Lygus Bugs apply 12-18 lbs per acre at 7-10 day intervals. Foliage application only. Do not apply within 5 days of harvest.

STRAWBERRIES: For control of Meadow Spittlebug and Strawberry Leaf Roller, apply 12-18 lbs per acre at 7-10 day intervals up to 14 days before harvest.

CUCUMBERS and SUMMER SQUASH: For control of Striped and Spotted Cucumber Beetles, Spittlebugs and Flea Beetles, apply 8-10 lbs per acre at 7-10 day intervals. Do not apply within 15 days of harvest and not before vining on cucumbers.

TOMATOES, EGGPLANT and PEPPERS: For control of Flea Beetles apply 7-10 lbs per acre at 7-10 day intervals. For control of European Corn Borer, Armyworm, Tomato Fruitworm and Tomato Hornworm apply 15-20 lbs per acre at 7-10 day intervals. For control of Stink Bugs apply 25-30 lbs per acre. Do not apply within 10 days of harvest on Tomatoes and 15 days before harvest on Eggplant and Peppers.

CORN: For control of Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetle, Japanese Beetle, apply 10-20 lbs per acre at 7-10 day intervals up to day of harvest. Except when used for control do not apply within 14 days of harvest or cutting for forage. Application of this product to the sil region of corn during the pollen shed period will seriously reduce bee populations.

LETTUCE: For control of Flea Beetles, Leafhoppers and Harlequin Bugs, apply 7-10 lbs per acre. To control the Six Spotted Leafhopper, apply 10-15 lbs per acre. To control Corn Earworm, Armyworm, Cabbage Caterpillars, Spittlebugs, Lygus and Stink Bugs, apply 10-20 lbs per acre. Make application at 7-10 day interval. On Head Lettuce do not apply later than 7 days before harvest. On Leaf Lettuce do not apply within 14 days of harvest.

ORRA: For control of Corn Earworm and Stink Bugs, apply 12-20 lbs per acre at a 5-7 day interval up to 21 days of harvest.

ALFALFA, CLOVERS, COMPEAS (MIDWESTERN OR FIELD PEAS), PEANUTS, SOYBEANS: For control of Blister Beetle and Mexican Bean Beetle, apply 7 to 14 lbs per acre. To control Thrips, Leafhoppers, Spittlebugs, Armyworm, Cucumber Beetle, Bean Leaf Hopper, Japanese Beetle, Flea Beetles, Lygus Bugs, and Stink Bugs, apply 12-15 lbs per acre. To control Armyworm, Leafhopper, Corn Earworm and Stink Bugs, apply 14-15 lbs per acre. Do not apply within 15 days of harvest.

SORGHUMS: (including MILO and GRAIN SORGHUM) For control of Armyworm, Weevils, Corn Earworms and Stink Bugs, apply 14-20 lbs per acre. Do not apply within 21 days of harvest on grain sorghum or 12 days of harvest for forage.

EPA Est. No. 6735-TX-10
EPA Registration No. 6735-100

A PRODUCT OF
TIDE PRODUCTS, INC.
EDINBURG, TEXAS 78539

BEST DOCUMENT AVAILABLE

OBSERVE FOLLOWING PRECAUTIONS

- This product is highly toxic to bees exposed to direct treatment or to crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.
- Do not use on feed or forage to be fed to dairy animals or to animals finished for slaughter.
- Avoid dust drift onto adjoining food and forage crops about to be harvested.
- Apply in early morning or evening when there is little or no wind to protect bees and to prevent drift.
- This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment. Apply this product only as specified on this label.
- The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME

DANGER-POISON

- POISONING SYMPTOMS:** Headache, blurred vision, weakness, nausea, diarrhea and discomfort in chest.
- ANTIDOTE:** Atropine is the specific antidote. Obtain emergency supply of atropine 1-100 grain tablets from your physician. Keep on hand.
- CALL A PHYSICIAN IMMEDIATELY** if persons handling or working with Parathion develop poisoning symptoms.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

FIRST AID TREATMENT

IF SWALLOWED — Give tablespoonful of salt in a glass of warm water. Repeat until vomit fluid is clear. Have victim lie down and keep quiet.

INHALATION — Remove patient from contaminated atmosphere.

SKIN — In case of contact, remove contaminated clothing and immediately wash skin thoroughly with soap and water.

EYES — In case of contact, flush immediately with water for at least 15 minutes.

- INFORMATION FOR PHYSICIAN (Treatment)** — Give atropine preferably by injection, grains 1.30 to 1.15 (2 to 4 mg.), at once and every hour or oftener as needed to achieve and maintain full atropinization. 2-PAM is also antidote and may be administered in conjunction with atropine. Never give morphine. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously for 48 hours.

DANGER

POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. RAPIDLY ABSORBED THROUGH SKIN. DO NOT GET IN EYES OR ON SKIN.

Wear natural rubber gloves, protective clothing and goggles. Wear a mask or respirator of a type passed by the U.S. Department of Agriculture for Parathion protection.

Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated.

Do not contaminate feed and foodstuffs. Wash all contaminated clothing with soap and hot water before re-use. Wash hands, arms, and face thoroughly with soap and water before eating or smoking.

NOTICE — Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty or merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

READ LABEL BEFORE USING—DESTROY EMPTY CONTAINER—BURY OR BURN—STAY OUT OF SMOKE OR FUMES

BEST DOCUMENT AVAILABLE