

ACCEPTED
Feb 17 1969

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND HERBICIDE ACT
FOR ECONOMIC PESTS REGISTERED
UNDER NO. 6735-189

DO NOT USE

with skin. Avoid breathing spray
or food products. Keep children and
been washed into the soil.
physician at once.

weed control, particularly for grassy
IT IS UNNECESSARY TO ADD
N. Best results are obtained on
70° F.

the spray equipment reservoir about
with agitation. Finish filling the
thoroughly by flushing with water

AS TO ITS CHEMICAL DESCRIP-
TION STATED ON THE LABEL WHEN
NORMAL CONDITIONS OF USE.
WARRANTY OR MERCHANTABILITY
EXPRESS OR IMPLIED, EXTENDS TO
INSTRUCTIONS, OR UNDER AB-
SEASONABLY FORSEEABLE TO
SUCH USE.

GAL NET



TIDE DISO



HERBICIDE

For Selective Post-Emergence Weed Control

ACTIVE INGREDIENT:

Disodium Methanearsonate*

21.8%

INERT INGREDIENTS:

Total Arsenic (as elemental) all in water soluble form 8.9%

78.2%

*Product contains 3.6 pounds Disodium Methanearsonate Hexahydrate per gallon

CAUTION

KEEP OUT OF REACH OF CHILDREN

See left side panel for additional cautions

CONTAINER DISPOSAL: Do not reuse container. Wash thoroughly with water and detergent, crush if possible, and discard in a safe place.

PACKAGED BY

TIDE PRODUCTS, INC.

EDINBURG, TEXAS 78539

ONLY the BEST . . . is fit to be TIDE

READ ENTIRE LABEL BEFORE USING

COTTON: TIDE DISO Herbicide is useful as a directed application in cotton for post-emergent control of weeds and grasses such as Johnsongrass, nutsedge, Dallisgrass, cocklebur, barnyardgrass, ragweed, sandbur and puncturevine.

Add 1 gallon of TIDE DISO Herbicide to 40 gallons of water and mix thoroughly. Do not add any surfactant to the spray solution. Apply spray solution at 40 gallons per acre. Weed control is most effective at temperatures above 70° F. For band application in 40 inch rows, apply 1 gallon of spray solution per inch of band width per acre. Direct the spray solution to the base of the cotton plant. Nozzles should be placed so as to avoid spraying the cotton foliage. A second application may be required during humid growing conditions to control perennials such as Johnsongrass and nutcracker. Apply the second application 1 to 3 weeks later. Slight burning and a reddish discoloration of the cotton leaf may occasionally occur following the recommended treatment, but the cotton plant will develop normally.

APPLY ONLY WHEN COTTON IS 3 INCHES HIGH TO FIRST BLOOM. DO NOT APPLY AFTER FIRST BLOOM. DO NOT FEED TREATED FOLIAGE TO LIVESTOCK OR GRAZE TREATED AREAS.

NON-CROP: TIDE DISO Herbicide is useful for controlling Johnsongrass, nutsedge, watergrass and certain other weeds in drainage ditchbanks, rights-of-way, storage yards, and similar noncrop areas.

Mix TIDE DISO Herbicide at a rate of 1 to 2 gallons in 100 gallons of water. No additional surfactant is necessary. Spray the unwanted vegetation thoroughly to just short of runoff. Any spray equipment that gives good coverage may be used. If regrowth occurs, reapply as required.

DO NOT CONTAMINATE WATERS USED FOR DOMESTIC CONSUMPTION, OR BY ANIMALS, WILDLIFE AND AQUATIC LIFE, OR FOR IRRIGATION PURPOSES.

LAWN AND ORNAMENTAL TURF: TIDE DISO Herbicide is useful for control of Dallisgrass, sandbur, bahiagrass, nutsedge, barnyardgrass, chickenweed and wood sorrel with little or no injury to good lawn grasses. On new lawns, do not treat until after three mowings. Good grasses may be temporarily discolored. Zoysia, bluegrasses, and Bermuda are quite tolerant. Do not use on St. Augustine or centipede.

Mow lawns 1 to 1½ inches high before treatment. Mix 4 fluid ounces of TIDE DISO Herbicide in 5 gallons of water and apply to an area of 1,000 square feet. Spray thoroughly to wet all undesirable plants. Repeat applications, 10 to 14 days apart, may be needed for good control.

GPS 2 1 00 1 1