

READ LABEL BEFORE USING

EYES — In case of contact, flush immediately with water for at least 15 minutes.

(Treatment) — Give atropine preferably by injection, grains 1/30 to 1/15 (2 to 4 mg.), at once and every hour or oftener as needed to achieve and maintain full atropinization. 2-PAM is also antidotal and may be administered in conjunction with atropine. Never give morphine. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Treat symptomatically. Observe patient continuously for 48 hours. PARATHION is a Cholinesterase inhibitor and treatment of poisoning should include the use of ATROPINE. Toxaphene produces central nervous system excitation which ordinarily responds to barbituric acid derivatives, but this drug should be considered only if complete atropinization fails to relieve symptoms.

Do not apply to cucurbits (cantaloupe, cucumber, squash, melons) as injury to the plant may result.

Avoid drift to adjoining food and forage crops about to be harvested, or to crops in bloom, or areas occupied by unprotected humans or beneficial animals.

Do not apply to feed or forage crops to be fed to dairy animals or to animals being finished for slaughter.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Co-operative Agricultural Extension Service.

This product is toxic to fish, shrimp, crab, birds and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

TIDE
Symbol of Quality
EXPANDER

EMULSIFIABLE LIQUID-AGRICULTURAL INSECTICIDE

Contains 1.5 lbs. Methyl Parathion and 6 lbs. Toxaphene per Gallon

ACTIVE INGREDIENTS:

- *O, O-dimethyl O-p-nitrophenyl thiophosphate 13.95%
- **Toxaphene 55.80%
- Xylene 26.40%

INERT INGREDIENTS:

- *Methyl Parathion 3.85%
- **Technical chlorinated camphene (chlorine content 67% to 69%) 100.00%

DANGER POISON



SEE ANTIDOTE STATEMENT AND OTHER REQUIRED WARNING STATEMENT ON LEFT PANEL

PROTECT FROM TEMPERATURES BELOW 32 °F. DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME

A PRODUCT OF TIDE PRODUCTS, INC.

1 GALLON NET EDINBURG, TEXAS 78539

ONLY the BEST . . . is fit to be TIDE

E. P. A. Registration No. 6735-218

EPA Est. No. 6735-TX-1

FOR USE ONLY BY OPERATORS EXPLICITLY TRAINED IN PARATHION INSURE

DIRECTIONS:

Application should be made by trained personnel using appropriate equipment. Do not apply with more than the specified amount of the concentrate per spray tank. Add buffer material to spray tank. Keep agitator running during operations. Do not apply to plants if mixture does not separate as an oily emulsion. Plants may result in powdery residues unless proven physically. Thoroughly emulsify mixture. Half of total weight should be in powder form.

SUGGESTED WATER AND GROUND APPLICATION RATES: Required to provide adequate coverage. Except as specified, application rates are therefore based on a range of conditions.

Crop: Vegetable and Field Crops

RECOMMENDATIONS:

(The dosage rates are incorporated with suitable coverage).

COLE CROPS (Brassicae, Cauliflowers, etc.): Imported Cabbage Rootworm, Cabbage Loopers, Cabbage Cutworms (cutworms), Cabbage Moth, Flea Beetles, etc. the rate of 1 1/2 to 2 quarts per acre. Do not apply more than once within 30 days.

COTTON: For bollworm, species apply 1 1/2 to 2 quarts per acre. For Fleahopper, Rapid Pyralis, Tarnished Plant Bug, etc. apply 1 1/2 to 2 quarts per acre.

Wear natural rubber gloves, protective clothing and goggles.
Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for Methyl Parathion protection.
Airplane pilots should wear full face canister-type mask.
Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be reentered until drifting insecticide and volatile residues have dissipated.
Do not contaminate feed and foodstuffs.
Wash hands, arms, and face thoroughly with soap and water before eating or smoking.
Wash all contaminated clothing with soap and hot water before re-use.
NOT FOR USE OR STORAGE IN OR AROUND THE HOME

DANGER POISON

Headache, blurred vision, weakness, nausea, cramps, diarrhea and discomfort in chest.


Atropine is the specific antidote. Obtain emergency supply of Atropine 1/100 grain tablets from your physician. Keep on Hand.

If persons handling or working with Methyl Parathion develop poisoning symptoms.

IF SWALLOWED — Give tablespoonful of salt in a glass of warm water. Repeat until vomit fluid is clear. Have victim lie down and keep quiet.

INHALATION — Remove patient from contaminated atmosphere.

SKIN — In case of contact, remove contaminated clothing and immediately wash skin thoroughly with soap and water.



TIDE

Symbol of Quality

EXPANDER

EMULSIFIABLE LIQUID-AGRICULTURAL INSECTICIDE

Contains 1.5 lbs. Methyl Parathion and 6 lbs. Toxaphene per Gallon

ACTIVE INGREDIENTS:

*O, O-dimethyl O-p-nitrophenyl thiophosphate
 **Toxaphene
 Xylene

13.95%
 55.87%
 26.40%
 3.85%

INERT INGREDIENTS:

*Methyl Parathion
 **Technical chlorinated camphene (chlorine content 67% to 69%).

100.00%

DANGER POISON

Keep Out of Reach of Children



SEE ANTIDOTE STATEMENT AND OTHER REQUIRED WARNING STATEMENT ON LEFT PANEL

PROTECT FROM TEMPERATURES BELOW 32 °F.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME

A PRODUCT OF

TIDE PRODUCTS, INC.

1 GAL. NET EDINBURG, TEXAS 78539

ONLY the BEST . . . is fit to be TIDE

READ LABEL BEFORE USING

FOR USE ONLY IN POWER SPRAYERS BY OPERATORS EXPERIENCED OR ADEQUATELY TRAINED IN THE APPLICATION OF PARATHION INSECTICIDES.

DIRECTIONS FOR USE

Application should be made only by trained personnel using approved protective equipment. Do not apply with hand equipment. Pour specified amount of this product into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. If mixture does not mix readily, but tends to separate as an oily layer, do not use as injury to plants may result. Do not combine with wettable powders unless previous use of the mixture has proven physically compatible and safe. Always thoroughly emulsify this product with at least half of total water before adding wettable powder.

SUGGESTED WATER RATES FOR AIRCRAFT AND GROUND APPLICATION. (The actual rate required to provide thorough, uniform coverage varies with plant growth at time of application. Except as specified for certain uses, the following rates are therefore intended to cover a broad range of conditions.)

Crop	Gal. Water/Acre	
	Aircraft	Ground
Vegetable and Field Crops	1-20	5-100

RECOMMENDATIONS

(The dosage rates given below are to be incorporated with sufficient water to provide adequate coverage).

COLE CROPS (Broccoli, Brussel Sprouts, Cabbage, Cauliflower): Aphids, Cabbage Looper, Imported Cabbageworms, Army Worms, Climbing-Cutworms (certain species), Diamond Back Moth, Flea Beetles, Leafhoppers, Mites. Apply at the rate of 1½ to 2 qts. per acre at 7-10 day intervals. Do not apply after edible parts begin to form or within 30 days of harvest.

COTTON: For Climbing-Cutworms (certain species) apply 1½ qt. per acre. For Army Worms, Eleahopper, Rapid Plant Bug, Southern Stink Bug, Tarnished Plant Bug, Lygus, Aphid, Boll

Weevil, Bollworm, Red Spider Mites, Thrips. Apply at the rate of 1½-2 qts. per acre at 5 day intervals. Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season. Do not hand pick or harvest within 7 days of application. Workers entering fields within 24 hours after application should wear protective clothing.

PEPPERS: Aphid, Army Worms, Flea Beetles, Leafhopper, Spider Mites. Apply at the rate of 1½ qts. per acre. Do not apply within 15 days of harvest.

TOMATOES: Aphids, Army Worms, Climbing Cutworms (certain species), Tomato Fruitworm, Cabbage Loopers, Tomato Hornworm, Blister Beetles, Colorado Potato Beetle, Flea Beetle, Psyllids, Thrips, Leafhoppers, Spider Mites, Russet Mites, Serpentine Leafminer. Apply at the rate of 1½-2 qts. per acre. Do not apply within 15 days of harvest.

RICE: Climbing cutworms, Armyworms, Grasshoppers, Rice Stinkbug — Apply 1½-2 qts. per acre. Do not feed treated forage to dairy animals or animals being finished for slaughter. Crayfish, shrimp and crabs will be killed at application rates recommended on this label. Do not use where these are important resources. Do not apply within 15 days of harvest.

CONTAINER DISPOSAL

1. Empty container into spray tank. Then drain in verticle position for 30 seconds.
2. Add a measured amount of rinse water so container is ¼ to 1/5 full.
3. Rinse container thoroughly, pour into tank, and drain for 30 seconds. Repeat 3 times. Add enough fluid (water) to bring tank up to level.
4. Crush pesticide container. Sell as scrap for recycling or bury. Do not reuse.

NOTICE: SELLER WARRANTS THAT THE PRODUCT CONFORMS TO ITS CHEMICAL DESCRIPTION AND IS REASONABLY FIT FOR THE PURPOSE STATED ON THE LABEL WHEN USED IN ACCORDANCE WITH DIRECTIONS UNDER NORMAL CONDITIONS OF USE, BUT NEITHER THIS WARRANTY NOR ANY OTHER WARRANTY OR MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, EXPRESS OR IMPLIED, EXTENDS TO THE USE OF THIS PRODUCT CONTRARY TO LABEL INSTRUCTIONS, OR UNDER ABNORMAL CONDITIONS, OR UNDER CONDITIONS NOT REASONABLY FORSEEABLE TO SELLER AND BUYER ASSUMES THE RISK OF ANY SUCH USE.

GPS 10-74 10-(11137) H-74 H-80

E. P. A. Registration No. 6735-218
 EPA Est. No. 6735-TX-1