


DANGER-POISON-PRECAUTIONS



POISONOUS IF SWALLOWED
This product can kill you if swallowed even in small amounts; spray mist or dust may be fatal if swallowed.



POISONOUS BY SKIN CONTACT
Poisonous if touched by hands or spilled or splashed on skin. In eyes or on clothing (liquid goes through clothes.)



POISONOUS IF BREATHED
Breathing vapors, spray mist or dust may be fatal.

**RQ ORGANOPHOSPHOROS PESTICIDE,
LIQUID, N.O.S. UN-2783**

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses authorized by the Certified Applicator's certification.

BUDMOR 33

Emulsifiable Concentrate Insecticide

KEEP OUT OF REACH OF CHILDREN

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

POISON DANGER POISON

DANGER

Poisonous if swallowed, inhaled or absorbed through the skin. Rapidly absorbed through the skin or eyes. Do not get in eyes, on skin or on clothing. Do not breathe the spray mist or vapor. Rapid inhalation or skin contact may, without symptoms, progressively increase susceptibility to poisoning. Clothing wet with this material must be removed immediately. Exposed persons must receive prompt medical attention or they may die.

USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE EQUIPMENT AND CLOTHING

Protective Clothing Required:

INFORMATION FOR PHYSICIAN

Give 2 to 4 mg. (1/30 to 1/15 grain) of atropine immediately, preferably by intravenous injection, and repeat in 5 to 10 minute intervals until signs of atropinization appear (dry, flushed skin, dilated pupils, rapid pulse). Maintain full atropinization and give 1 mg. (adult dosage) Protopam (2-PAM) intravenously, slowly, to restore normal erythrocyte cholinesterase levels. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Methyl Parathion and EPN are strong cholinesterase inhibitors affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At the first sign of pulmonary edema, the patient should be given supplementary oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement. VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until cholinesterase regeneration has been attained as determined by a blood test.

ENVIRONMENTAL HAZARDS

Toxic to fish and wildlife. Birds and other mammals out of range.

POISON

See side panels for precautionary and first aid statements



DANGER

KEEP OUT OF REACH OF CHILDREN with COMMENTS in EPA Letter D-67

NOT FOR HOME USE

NOV 28 1983

to direct seeds. Do to blooming tment area.

ACTIVE INGREDIENTS:

1. 0-p-nitrophenyl phosphorothioate*
s favor ds
phenyl phenyl phosphonothioate**
omatic Solvent

INGREDIENTS

TOTAL

Wear clean protective clothing, goggles, and respirator approved by NIOSH or the American National Standards Institute when applying or handling, or when reentering fields within 24 hours of treatment.

The following protective clothing must be worn: lightweight, unlined, natural rubber gloves at least mid-forearm in length; a wide brimmed waterproof hat or waterproof hood; a protective suit or coveralls of a non-permeable, non-cloth material covering the body from ankles to wrists; lightweight, unlined, natural rubber boots at least mid-calf in length; full-face respirators are recommended; half-face respirators and goggles are required.

Aerial applicators in positive pressure cockpits and other applicators in comparable ground equipment with appropriate filters at all air intakes need not comply with these protective clothing requirements.

POISON SIGNS (Symptoms)

WARNING SYMPTOMS Include weakness, headache, tightness in the chest, labored respiration, blurred vision, nausea, abdominal cramps, diarrhea, sweating, vomiting, salivation and redreactive p-nitrophenyl pupils.

FIRST AID TREATMENT

IF SWALLOWED—Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or blunt object. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

INHALATION—Remove victim from contaminated atmosphere.

SKIN—In case of contact, immediately remove patient from the vicinity of insecticides. Remove contaminated clothing and immediately wash skin thoroughly with soap and water.

EYES—In case of contact, immediately remove victim from the vicinity of the pesticide. Flush eyes immediately with water for at least 15 minutes.

IF WARNING SYMPTOMS (See Above) OCCUR—Keep victim prone and quiet. Start artificial respiration if victim is not breathing. Transport victim to nearest hospital immediately.

ANTIDOTE. ATROPINE is the specific antidote for Methyl parathion poisoning. Obtain emergency supply of atropine 1/100 grain tablets from your physician. Keep on hand.

ter, agitate, and the proper amount of water to the spray tank with water while still agitating. Continue to agitate while using.

For aerial applications, flagging must be by fully automated mechanical means or by humans working in totally enclosed vehicles.

1. Prohibitions
Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty containers.

2. Pesticide Disposal
Pesticide, spray mixture or rinse water that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

3. Container Disposal
Triple rinse, and offer for recycling, reconditioning, or disposal in approved landfill, or bury in a safe place.

4. General
Contact federal, state or local disposal authorities for approved alternatives.

NOTICE Seller warrants that the product conforms to its contract description and is free from any defect in the product stated on the label when used in accordance with directions under normal conditions of use. But neither this warranty nor any other warranty of merchantability or fitness for a particular purpose expressed or implied extends to the use of this product contrary to label instructions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

NET CONTENTS: _____ GAL.

E.P.A. Reg. No. 6735-238
E.P.A. Est. No. 6735-TX-10

A PRODUCT OF
TIDE PRODUCTS, INC.
EDINBURG, TEXAS 78539

ONLY the BEST . . . is fit to be TIDE

BEST DOCUMENT A

**RQ ORGANOPHOSPHOROS PESTICIDE,
LIQUID, N.O.S. UN-2783**

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses approved by the Certified Applicator's certification.

BUDMOR 33

Emulsifiable Concentrate Insecticide

KEEP OUT OF REACH OF CHILDREN

POISON
See side panels
for precautionary
and first aid statements



DANGER
KEEP OUT OF REACH OF CHILDREN
with COMMENTS
in EPA Label Panel

NOT FOR HOME USE

NOV 28 1983

to direct
seeds. Do
to blooming
tment area.

ACTIVE INGREDIENTS:

Disopropyl 1-naphthyl phosphorothioate*	29.83%
Disopropyl 2-naphthyl phosphorothioate*	1.57%
Diethyl 1-naphthyl phenyl phosphonothioate**	31.40%
Diethyl 2-naphthyl phenyl phosphonothioate**	26.14%
Inert Solvent	11.06%
TOTAL	100.00%

E.P.A. Reg. No. 6735-238

E.P.A. Est. No. 6735-TX-10

NET CONTENTS: _____ GAL.

BEST DOCUMENT AVAILABLE

A PRODUCT OF
TIDE PRODUCTS, INC.
EDINBURG, TEXAS 78539
ONLY the BEST . . . is fit to be TIDE

CROP	INSECT	RATE OF BUDMOR 33 PER ACRE	APPLICATION DIRECTIONS AND LIMITATIONS			
Cotton	Aphids	2/3 to 1-1/3 pints	Do not apply to blooming cotton treatment area. Do not feed treated forage or silage to livestock.			
	Hill Weevil					
	Cotton Leafhopper					
	Cotton Leafworm					
	Fall Armyworm					
	Fleahoppers					
	Garden Webworm					
	Grasshoppers					
	Lycus Bugs (and other Miridae)					
	Red Spider Mites					
	Thrips					
	Darkling Ground Beetle			2 1/2 pints		
	Mexican Bean Beetle					
	Pink Bollworm					
	Plant Bug (Rapid, Tarnished)					
Russet Mites						
Serpentine Leafminer						
Southern Armyworm						
Sugarbeet Webworm						
Two-Spotted Mite						
Corn	Beet Armyworms	2-2/3 pints	Repeat application as necessary. Do not apply within 14 days of harvest.			
	Bollworm					
	Cabbage Looper					
	Cotton Leaf Perforator					
	Saltmarsh Caterpillar					
	Tobacco Budworm					
	Yellow Striped Armyworm					
	Chimney Cutworms			1/2 pint		
	Corn Leaf Aphids					
	Corn Rootworm (adult)					
	Fall Armyworms					
	Stinkbugs					
	Armyworms (up to third instar)			2/3 pint		
	European Corn Borer (sweet corn)			2/3 to 3/4 pints		
	European Corn Borer (field corn)			2/3 pint		
Southwestern Corn Borer (second to third generation)	2/3 pint					
Beans (Dry, green)	Lycus Bugs	1 1/2 pints	Make first application when 25% of plants show evidence of leaf feeding. Continue at 5 day intervals as long as feeding is evident. From 1 to 3 treatments may be required. For control of second brood borers apply when eggs begin to hatch. Do not apply within 14 days of harvest.			
	Darkling Ground Beetle					
	Becetia					
	Mexican Bean Beetle					
	Russet Mite					
	Serpentine Leafminer					
	Sugarbeet Webworm					
	Thrips					
	Tomato Fruitworm					
	Two-Spotted Mite					
	Soybeans			Chimney Cutworm	1 pint	Use restricted to the southeast. Do not make more than 2 applications during the growing season. Do not harvest or graze or feed forage from treated fields to livestock within 21 days after the last application.
				Stinkbugs		
				Velvetbean Caterpillar		
				Aphids		
				Buster Beetle		
Cotton Bollworm						
Garden Webworm						
Mexican Bean Beetle						
Three Corned Alfalfa Hopper						
Two-spotted Spider Mite						
Cabbage Looper		2 pints				
Corn Earworm						
Green Cloverworm						
Fall Armyworm (up to third instar)		2-2/3 pints				
Tomatoes		Potato Aphid	1 1/2 to 1 3/4 pints	Do not apply within 21 days of harvest.		
	Serpentine Leafminer					
	Southern Armyworm					
	Tomato Fruitworm					
	Mites	2-2/3 pints				
	Psyllids					
	Russet Mites					
	Sugar Beet Armyworm					
	Thrips					
	Two-spotted Mite					
	Yellow Striped Armyworm					
	Fire Beetle	2-2/3 pints				
	Cabbage Looper					

TANK MIXTURES FOR COTTON

This product may be tank mixed with Galleon 4E, Fundal 4EC, Fundal 4SP, Lannate, Lannate L, Nudrin 1B or Nudrin 60. When using these products in a tank mix observe all limitations and precautions which appear on the labels. Use the rate recommendations and application timings for the insect pest that is to be controlled as stated on the labels. Galleon 4E is a trademark of the Ciba-Geigy Corp. Fundal 4E and Fundal 4SP are trademarks of Schering A. E. Berlin/Berksham and distributed by NOH-AM Agricultural Products, Inc. Lannate and Lannate L are trademarks of E. I. DuPont Nemours & Co., Inc. Nudrin 1B and Nudrin 60 are trademarks of Shell Chemical Company.

PESTICIDE
783

PESTICIDE
Applicators or per-
sonnel for those users cav-
tion.

33

Insecticide

CHILDREN

DANGER

KEEP OUT OF REACH OF CHILDREN
with COMMENTS
In EPA Letter Dated

NOV 28 1983

6735-235

29.83%
1.57%
31.40%
28.14%
11.06%

TOTAL 100.00%

6735-238

6735-TX-10

DOCUMENT AVAILABLE


be TIDE

CROP	INSECT	RATE OF BUDMOR 33 PER ACRE	APPLICATION DIRECTIONS AND LIMITATIONS		
Cotton	Aphids	2/3 to 1-1/3 pints	Do not apply to blooming cotton if bees are visiting the treatment area. Do not feed treated forage or sila trash to livestock.		
	Holl Weevil				
	Cotton Leafhopper				
	Cotton Leafworm				
	Fall Armyworm				
	Fleahoppers				
	Garden Webworm				
	Grasshoppers				
	Lycus Bugs (and other				
	lice)				
	Red Spider Mites				
	Thrips				
	Darkling Ground Beetle			2 1/2 pints	
	Mexican Bean Beetle				
	Pink Bollworm				
Plant Bug (Rapid Tarnished)					
Russet Mites					
Serpentine Leafminer					
Southern Armyworm					
Sugarbeet Webworm					
Two-Spotted Mites					
Corn	Beet Armyworms	2-2/3 pints			
	Hollworm				
	Cabbage Looper				
	Cotton Leaf Perforator				
	Saltmarsh Caterpillar				
	Tobacco Budworm				
	Yellow Striped Armyworm				
	Climbing Cutworms	1/2 pint	Repeat application as necessary. Do not apply within 14 days of harvest.		
	Corn Leaf Aphids				
	Corn Rootworm (adult)				
	Fall Armyworms				
	Stinkbugs				
	Armyworms (up to third instar)	2/3 pint	Make first application when 25% of the plants show evidence of leaf feeding. Continue at 5 day intervals as long as feeding is evident. From 1 to 3 treatments may be required. For control of second brood borers apply when eggs begin to hatch. Do not apply within 14 days of harvest.		
	European Corn Borer (sweet corn)	2/3 to 3/4 pints			
	European Corn Borer (field corn)	2/3 pint	Make application 10 days after eggs first hatch or when 75% of the plants show evidence of feeding in leaf whorls. Repeat 7 to 10 days later if new evidence of leaf feeding is observed. For control of second brood borers apply when eggs begin to hatch. Do not apply within 14 days of harvest.		
Southwestern Corn Borer (second to third generations)	2/3 pint	Apply when 10% of the plant population have egg masses. Repeat at 5 to 10 day intervals as needed. Do not apply within 14 days of harvest.			
Beans (Dry, green)	Lycus Bugs				
	Mites				
	Darkling Ground Beetle	1 1/2 pints			
	Beetles				
	Mexican Bean Beetle				
	Russet Mite				
	Serpentine Leafminer				
	Sugarbeet Webworm				
	Thrips				
	Tomato Fruitworm				
	Two-Spotted Mite				
	Soybeans	Climbing Cutworm	1/2 pint	Use restricted to the southeast.	
		Stinkbugs	1 pint	Do not make more than 2 applications during the growing season.	
		Velvetbean Caterpillar		Do not harvest or graze or feed forage from treated fields to livestock within 31 days after the last application.	
		Aphids	1-1/3 pints		
Bitter Beetle					
Cotton Bollworm					
Garden Webworm					
Mexican Bean Beetle					
Three Cornered Alfalfa Hopper					
Two-spotted Spider Mite					
Cabbage Looper		2 pints			
Corn Earworm					
Green Cloverworm					
Fall Armyworm (up to third instar)		2-2/3 pints			
Tomatoes		Potato Aphid	1 1/2 to 1 3/4 pints	Do not apply within 21 days of harvest.	
	Serpentine Leafminer				
	Southern Armyworm				
	Tomato Fruitworm				
	Mites	2-2/3 pints			
	Psyllids				
	Russet Mites				
	Sugar Beet Armyworm				
	Thrips				
	Two-spotted Mites				
	Yellow Striped Armyworm				
	Flea Beetle	2-2/3 pints			
	Cabbage Looper				

TANK MIXTURES FOR COTTON

This product may be tank mixed with Galecron 4E, Fendal 4FC, Fendal 4SP, Lannate, Lannate L, Nudrin 18 or Nudrin 90. When using these products in a tank mix observe all limitations and precautions which appear on the labels. Use the rate recommendations and application timings for the insect pest that is to be controlled as stated on the label.
*Galecron 4E is a trademark of the Ciba-Geigy Corp.
*Fendal 4E and Fendal 4SP are trademarks of Schering. A. E. Berlin, Bensheim and distributed by NOR-AM Agricultural Products, Inc.
*Lannate and Lannate L are trademarks of E. I. DuPont Nemours & Co., Inc.
*Nudrin 18 and Nudrin 90 are trademarks of Shell Chemical Company.

DANGER-POISON-PRECAUTIONS



**RQ ORGANOPHOSPHOROS PESTICIDE
LIQUID, N.O.S. UN-2783**

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

BUDMOR 33

Emulsifiable Concentrate Insecticide

KEEP OUT OF REACH OF CHILDREN

POISON

See side panels
for precautionary
and first aid statements



DANGER

KEEP OUT OF
REACH OF
CHILDREN
with COMMENTS
in EPA Letter D-67

NOT FOR
HOME
USE

NOV 28 1983

Wear clean, heavy duty, natural rubber gloves, waterproof pants, hat, rubber boots or rubber overboots and safety goggles when loading, mixing or applying this product. A pesticide label is hereby approved by the Environmental Enforcement and Safety Administration, formerly the U.S. Bureau of Plant Industry and by the National Institute for Occupational Safety and Health under the provisions of 30CFR PART 11.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

Do not apply this product when weather conditions favor drift from treated areas.

ACTIVE INGREDIENTS:

2,2-bis(4-chlorophenyl) ethyl dimethyl phosphorothioate*	2
2,2-bis(4-chlorophenyl) ethyl dimethyl phosphonothioate**	3
Nonionic Solvent	1
TOTAL	10

*Methyl Parathion
**EPN.

Contains 2.85 lbs. */gal. Methyl Parathion by isomer specific AOAC Methods 8.600/8.603 and 3 lbs. **/gal. EPN.

E.P.A. Reg. No. 6735-238
E.P.A. Est. No. 6735-TX-10

NET CONTENTS: _____ GAL.

BEST DOCUMENT A

A PRODUCT OF
TIDE PRODUCTS, INC.
EDINBURG, TEXAS 78539

ONLY the BEST . . . is fit to be TIDE

WORK SAFETY RULES

Keep all unprotected persons and children away from tree area where there is danger of drift. Do not rub eyes or mouth with hands. If you feel sick any way STOP work and get help right away. Call a doctor (physician), clinic or hospital — immediately. Explain to the doctor that you have been exposed to a Methyl Parathion-EPN combination and describe his condition. After first aid is given (see First Aid Treatment Section) and if a doctor cannot come, take victim to clinic or hospital. **IF SWALLOWED** Before removing gloves, wash them with water. Always wash hands, face and arms with water before smoking, eating or drinking. **AFTER WORK** Take off all work clothes and shoes using soap and water. Wear only clean clothes during your job. Do not wear contaminated clothing. Wash protective clothing and protective equipment with soap and water after each use. Respirator should be cleaned and filter replaced according to instructions included with respirator.

CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING. FIRST AID PROCEDURES, ANTIDOTES AND PHYSICIAN'S INSTRUCTIONS.

POISON SIGNS (Symptoms)

WARNING SYMPTOMS include weakness, headache, tightness in the chest, labored respiration, blurred vision, runny nose, muscular cramps, diarrhea, sweating, vomiting, salivation and nonreactive pupils.

FIRST AID TREATMENT

IF SWALLOWED—Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or blunt object. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.
INHALATION—Remove victim from contaminated atmosphere.
SKIN—In case of contact, immediately remove patient from the vicinity of insecticide. Remove contaminated clothing and immediately wash skin thoroughly with soap and water.
EYES—In case of contact, immediately remove victim from the vicinity of the pesticide. Flush eyes immediately with water for at least 15 minutes.
IF WARNING SYMPTOMS (See Above) OCCUR—Keep victim prone and quiet. Start artificial respiration if victim is not breathing. Transport victim to nearest hospital immediately.
ANTIDOTE—ATROPINE is the specific antidote for Methyl parathion poisoning. Obtain emergency supply of atropine 1/100 grain tablets from your physician. Keep on hand.


When a dosage rate is given, use the lower rate on small, young plants or where insect infestation is light. Use the higher rate on large or mature plants or where insect infestation is heavy. BUDMOR 33 may be applied by ground or aerial application. When making ground applications, apply the recommended dosage in at least 3 gallons of water per acre, using sufficient water to give good coverage of the foliage. For aerial application, use 3 to 10 gallons of water per acre. To make spray mixture, fill half of the spray tank with water, agitate, add the proper amount of BUDMOR 33, then fill the spray tank with water while still agitating. Continue to agitate while using.

For aerial applications, flying must be by fully automated mechanical means or by humans working in totally enclosed vehicles.

- 1. Prohibitions**
Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty container.
- 2. Pesticide Disposal**
Pesticide spray mixture or rinse water that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.
- 3. Container Disposal**
Triple rinse, and offer for recycling, reconditioning, or disposal in approved landfill, or bury in a safe place.
- 4. General**
Consult federal, state or local disposal authorities for approved alternatives.

NOTICE Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose expressed or implied extends to the use of this product contrary to label instructions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

DANGER-POISON-PRECAUTIONS



POISONOUS IF SWALLOWED
This product can kill you if swallowed even in a small amount; spray mist or dust may be fatal if swallowed.



POISONOUS BY SKIN CONTACT
Poisonous if touched by hands or spilled or splashed on skin, in eyes or on clothing (liquid goes through clothes.)



POISONOUS IF BREATHED
Breathing vapors, spray mist or dust may be fatal.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

POISON **POISON**
DANGER

Poisonous if swallowed, inhaled or absorbed through the skin. Rapidly absorbed through the skin or eyes. Do not get in eyes, on skin or on clothing. Do not breathe the spray mist or vapor. Rapid inhalation or skin contact may, without symptoms, progressively increase susceptibility to poisoning. Clothing wet with this material must be removed immediately. Exposed persons must receive prompt medical attention or they may die.

USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE EQUIPMENT AND CLOTHING

Wear clean, heavy duty, natural rubber gloves, waterproof pants, a hat, rubber boots or rubber overboots and safety goggles when mixing, spraying or applying this product. Wear a pesticide resistant hoodily approved by the Environmental Protection Agency, and Safety Administration, U.S. Bureau of Mines and by the National Institute for Occupational Safety and Health under the provisions of 30CFR PART 11.

WORK SAFETY RULES

Keep all unprotected persons and children away from treated area where there is danger of drift. Do not rub eyes or mouth with hands. If you feel sick in any way STOP work and get help right away. Call a doctor (Physician), clinic or hospital — immediately. Explain that the victim has been exposed to a Methyl Parathion-EPN combination and describe his condition. After first aid is given (see First Aid Treatment section) and if a doctor cannot come, take victim to clinic or hospital. If ALERT! Before removing gloves, wash them with soap and water. Always wash hands, face and arms with soap and water before smoking, eating or drinking. AFTER WORK, take off and wash clothes and shoes. Shower, using soap and water. Wear only clean clothes when leaving job. Do not wear contaminated clothing. Wash protective clothing and protective equipment with soap and water after each use. Respirator should be cleaned and later replaced according to instructions included with respirator.

CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING. FIRST AID PROCEDURES, ANTIDOTES AND PHYSICIAN'S INSTRUCTIONS.

POISON SIGNS (Symptoms)

WARNING SYMPTOMS include weakness, headache, tightness in the chest, labored respiration, dilated pupils, muscular abdominal cramps, diarrhea, sweating, vomiting, salivation and nonreactive pinpoint pupils.

FIRST AID TREATMENT

IF SWALLOWED—Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or blunt object. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.
INHALATION—Remove victim from contaminated atmosphere.
SKIN—In case of contact, immediately remove patient from the vicinity of insecticides. Remove contaminated clothing and immediately wash skin thoroughly with soap and water.
EYES—In case of contact, immediately remove victim from the vicinity of the pesticide. Flush eyes immediately with water for at least 15 minutes.
IF WARNING SYMPTOMS (See Above) OCCUR—Keep victim prone and quiet. Start artificial respiration if victim is not breathing. Transport victim to nearest hospital immediately.
ANTIDOTE, ATROPINE is the specific antidote for Methyl parathion poisoning. Obtain emergency supply of atropine 1/100 grain tablets from your physician. Keep on hand.

INFORMATION FOR PHYSICIAN

Give 2 to 4 mg. (0.20 to 1.15 grains) of atropine, immediately, preferably by intravenous injection, and repeat at 5 to 10 minute intervals until signs of atropinization appear (dry, flushed skin, dilated pupils, rapid pulse). Maintain full atropinization and give 1 mg. adult dosage Protopam (2-PAM) intravenously, slowly, to restore normal erythrocyte cholinesterase levels. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Methyl Parathion and EPN are strong cholinesterase inhibitors affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At the first sign of pulmonary edema, the patient should be given supplementary oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement. VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until cholinesterase regeneration has been attained as determined by a blood test.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL. This product is highly toxic to bees exposed to direct treatment and should not be applied to blooming crops or honey bees in bloom. Obtain more information from your Cooperative Agricultural Extension Service.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Re-entry Statement
Workers entering treated fields within 48 hours of application should wear protective clothing.
Where a dose rate is given, use the lower rate on small young plants or where insect infestation is light; use the higher rates on large or mature plants or where insect infestation is heavy. BUDMOR 33 may be applied by ground or aerial application. When making ground applications, apply the recommended dosage in at least 3 gallons of water per acre, using sufficient water to give good coverage of the leaves. For aerial application, use 3 to 10 gallons of water per acre.
To make spray mixture, fill half of the spray tank with water, agitate, add the proper amount of BUDMOR 33, then fill the spray tank with water while still agitating. Continue to agitate while using.
POST TREATED AREA
Consult your State Agricultural Extension Service or Experiment Station regarding posting of treated areas.

STORAGE AND DISPOSAL

1. Prohibitions
Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open burning is prohibited. Do not reuse empty containers.
2. Pesticide Disposal
Pesticide spray mixture or rinse water that cannot be used or which is left over should be disposed of in a landfill approved for pesticides or buried in a safe place away from water sources.
3. Container Disposal
If possible, offer for recycling, reconditioning or disposal in an approved landfill, or bury in a safe place.
4. General
Consult federal, state or local disposal authorities for approved alternatives.

NOTICE—Tide warrants that the product conforms to its chemical description and is suitable for the purpose stated on the label when used in a proper and directed manner. It does not warrant or imply that it is a warranty for any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

**RQ ORGANOPHOSPHOROS PESTICIDE,
LIQUID, N.O.S. UN-2783**

RESTRICTED USE PESTICIDE
For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

BUDMOR 33

Emulsifiable Concentrate Insecticide

KEEP OUT OF REACH OF CHILDREN

POISON
See side panels for precautionary and first aid statements



DANGER
KEEP OUT OF REACH OF CHILDREN with COMMENT in EPA Letter to

NOT FOR HOME USE

NOV 28 198

ACTIVE INGREDIENTS:
O,O-Dimethyl O-p-nitrophenyl phosphorothioate*
Related compounds
O-Ethyl O-(p-nitrophenyl) phenyl phosphonothioate**
Xylene Range Aromatic Solvent
INERT INGREDIENTS

TOTAL

*Methyl Parathion
**EPN.
Contains 2.85 lbs **/gal Methyl Parathion by isomer specific AOAC Methods 6400/6405 and 3 lbs **/gal EPN.
E.P.A. Reg. No. 6735-238
E.P.A. Est. No. 6735-TX-10

NET CONTENTS: _____ GAL.
BEST DOCUMENT
A PRODUCT OF
TIDE PRODUCTS, INC.
EDINBURG, TEXAS 78539
ONLY the BEST . . . is fit to be TIDE

Protective Clothing Required: ← **BOLD TYPE**

Wear clean protective clothing, goggles, and respirator approved by NIOSH or the American National Standards Institute when applying or handling, or when reentering fields within 24 hours of treatment.

The following protective clothing must be worn: lightweight, unlined, natural rubber gloves at least mid-forearm in length; a wide brimmed waterproof hat or waterproof hood; a protective suit or coveralls of a non-permeable, non-cloth material covering the body from ankles to wrists; lightweight, unlined, natural rubber boots at least mid-calf in length; full-face respirators are recommended; half-face respirators and goggles are required.

Aerial applicators in positive pressure cockpits and other applicators in comparable ground equipment with appropriate filters at all air intakes need not comply with these protective clothing requirements.

RQ ORGANOPHOSPHOROS PESTICIDE, LIQUID, N.O.S. UN-2783

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

BUDMOR 33

Emulsifiable Concentrate Insecticide

KEEP OUT OF REACH OF CHILDREN

POISON

See side panels
for precautionary
and first aid statements



DANGER

KEEP OUT OF
REACH OF
CHILDREN

with COMMENTS
in EPA 1600-1-110-1

NOV 28 1983

NOT FOR
HOME
USE

ACTIVE INGREDIENTS:

O,O-Dimethyl O-p-nitrophenyl phosphorothioate*	29.83%
Related compounds	1.57%
O-Ethyl O-(p-nitrophenyl) phenyl phosphonothioate**	31.40%
Xylene Range Aromatic Solvent	26.14%
INERT INGREDIENTS	11.06%

TOTAL 100.00%

*Methyl Parathion
**EPN.

Contains 2.5 lbs. * gal. Methyl Parathion by isomer specific AOAC
Methods 6400/6405 and 3 lbs. ** gal. EPN

E.P.A. Reg. No. 6735-238

E.P.A. Est. No. 6735-TX-10

NET CONTENTS.....GAL.

BEST DOCUMENT AVAILABLE

A PRODUCT OF
TIDE PRODUCTS, INC.
EDINBURG, TEXAS 78539

ONLY the BEST . . . is fit to be TIDE

CROP	INSECT	RATE OF BUDMOR 33 PER ACRE	APPLICATION DIREC AND LIMITATION	
Cotton	Aphids			
	16:1 Weevil			
	Cotton Leafhopper	2/3 to 1-1/3 pints	Make application at 5 to 7 until control is obtained. Do not apply within 7 days ing or harvest. Do not feed treated forage of livestock.	
	Cotton Leafworm			
	Fall Armyworm			
	Flea-hoppers			
	Garden Webworm			
	Grasshoppers			
	Lygus Bugs (and other Miridae)			
	Red Spider Mites			
	Thrips			
	Darkling Ground Beetle	2 1/2 pints		
	Mexican Bean Beetle			
	Pink Bollworm			
Plant Bug (Stapid, Tarnished)				
Russet Mites				
Serpentine Leafminer				
Southern Armyworm				
Sugarbeet Webworm				
Two-Spotted Mites				
Beet Armyworms	2-2/3 pints			
Bollworm				
Cabbage Looper				
Cotton Leaf Perforator				
Saltmarsh Caterpillar				
Tobacco Budworm				
Yellow Striped Armyworm				
Corn	Girdling Caterpillars	1/2 pint	Repeat application as necessa Do not apply within 14 days	
	Corn Leaf Aphids			
	Corn Rootworm (adult)			
	Fall Armyworms			
	Stinkbugs			
	Armyworms (up to third instar)	2/3 pint	Make first application when plants show evidence of leaf continue at 5 day intervals if feeding is evident. From 1 to may be required. For control of second brood when eggs begin to hatch. Do not apply within 14 days	
	Ku Ban Corn Borer (sweet corn)	2/3 to 3/4 pints		
	European Corn Borer (field corn)	2/3 pint	Make application 10 days at hatch or when 75% of the evidence of feeding in leaf w 7 to 10 days later if new at feeding is observed. For con brood borers apply when a hatch. Do not apply within harvest.	
	Southeastern Corn Borer (second to third generational field corn)	2/3 pint	Apply when 10% of the plant have egg masses. Repeat at intervals as needed. Do not apply within 14 days	
	Beans (Dry, green, lima)	Aphids Armyworms (up to third instar) Flea Beetles Leafhoppers Lygus Bugs Mites	1 1/2 pints	Do not apply within 21 days
	Beans (Dry, green)	Darkling Ground Beetles Mexican Bean Beetle Red Spider Mite Serpentine Leafminer Sugarbeet Webworm Thrips Tomato Fruitworm Two Spotted Mite	1 1/2 pints	
Soybeans	Girdling Caterpillars	1/2 pint	Use restricted to the Do not make more than 11 during the growing season Do not harvest or graze of from treated fields to livestock days after the last application	
	Stinkbugs	1 pint		
	Yellowish Green Caterpillar			
	Aphids			
	Borer Beetle			
	Cotton Bollworm			
	Garden Webworm			
	Mexican Bean Beetle			
	Three Cornered Alfalfa Hopper			
	Two-spined Spider Mite			
	Cabbage Looper			
Corn Earworm				
Green Cloverworm				
Pink Armyworm (up to third instar)	2-3/4 pints			
Tomatoes	Tomato Aphid			
	Scout Aphid			
	Serpentine Leafminer			
	Southern Armyworm			
	Tomato Fruitworm			
	Mites			
	Psyllids	2-2.3 pints		
	Harvest Mites			
	Sugar Beet Armyworm			
	Thrips			
	Two-spined Mites			
Yellow Striped Armyworm				
Flea Beetle				
Cabbage Looper	2-2.3 pints			

TANK MIXTURES FOR COTTON

This product may be tank mixed with Othello 45P, Funtal 45P, Farnatec 160, Nudrin 90, Wadon 45P to the extent permitted by a tank mix observe all label directions which appear on the labels for the rate of application and application interval. Do not apply to crops that are over 2500 lbs. per acre. Do not apply to crops that are over 2500 lbs. per acre. Do not apply to crops that are over 2500 lbs. per acre.

*EPA Reg. No. 6735-238 and E.P.A. Reg. No. 6735-238 are trademarks of Shell Chemical Company.

**EPA Reg. No. 6735-238 and E.P.A. Reg. No. 6735-238 are trademarks of Shell Chemical Company.

©1983 TIDE PRODUCTS, INC. EDINBURG, TEXAS

©1983 TIDE PRODUCTS, INC. EDINBURG, TEXAS

©1983 TIDE PRODUCTS, INC. EDINBURG, TEXAS

11/28

Mr. Edward H. Tefertiller
Tide Products Inc.
P.O. Box 1020
Edinburg, TX 78539

BEST DOCUMENT AVAILABLE

Dear Mr. Tefertiller:

Subject: Revised labeling per EPI Notice
of September 23, 1983
Tide EPN-5E,
EPA File Symbol 876-328-AA-6735
Budnor 42, EPA Registration No. 6735-154
Budnor 33, EPA Registration No. 6735-238
Tide EPN-4E, EPA Registration No. 6735-142

With the exception of the label for "Tide EPN-5E" (distributor) the labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy of each of the above labels is enclosed for your records. The distributor label is also accepted, but will remain with our records for this product.

Sincerely yours,

Jay S. Ellenberger
Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure

TS-767:ELLENBERGER:DCR-28434:WANG-1248C:vjr:Raven:479-2013:11/16/83

CONCURRENCES							
SYMBOL							
SURNAME							
DATE							

DIPHOSPHOROS PESTICIDE, D.O.S. UN-2783

RECOMMENDED USE PESTICIDE

Use only by Certified Applicators or under supervision, and only for those uses on the applicator's certification.

FOR 33

Concentrate Insecticide

KEEP OUT OF REACH OF CHILDREN



DANGER
KEEP OUT OF REACH OF CHILDREN
with COMMENTS in EPA Label Text

NOT FOR HOME USE

NOV 28 1983

Diphosphorothioate*	29.83%
Diphosphorothioate**	1.57%
	31.40%
	26.14%
	11.06%
TOTAL	100.00%

BEST AVAILABLE COPY

Former specific AOAC

E.P.A. Reg. No. 6735-238

E.P.A. Est. No. 6735-TX-10

CROP	INSECT	RATE OF DIPHOSPHOROS 33 PER ACRE	APPLICATION DIRECTIONS AND LIMITATIONS			
Cotton	Aphids	2 3/4 to 1-1/3 pints	Do not apply to bloom treatment area. Do not apply to livestock.			
	Earl Weevil					
	Cotton Leafhopper					
	Cotton Leafworm					
	Fall Armyworm					
	Fleahoppers					
	Garden Webworm					
	Grasshoppers					
	Lycus Bugs (and other Miridae)					
	Red Spider Mites					
	Thrips					
	Darling Ground Beetle			2 1/2 pints		
	Mexican Bean Beetle					
	Pink Bollworm					
	Plant Bug (Rapid, Tarnished)					
Russet Mite						
Serpentine Leafminer						
Southern Armyworm						
Sugarbeet Webworm						
Two-Spotted Mite	2-2 3/4 pints					
Belt Armyworms						
Heliothis						
Cabbage Looper						
Cotton Leaf Perforator						
Saltmarsh Caterpillar						
Tobacco Budworm						
Yellow Striped Armyworm						
Corn	Climbing Cutworm	1/2 pint	Repeat application as necessary. Do not apply within 14 days of harvest.			
	Corn Leaf Aphid					
	Corn Rootworm (adult)					
	Fall Armyworms					
	Stinkbugs					
Armyworms (up to third instar)	2 3/4 pints	Make first application when 25% of plants show evidence of leaf feeding. Continue at 5 day intervals as long as feeding is evident. From 1 to 3 treatments may be required. For control of second brood borers when eggs begin to hatch. Do not apply within 14 days of harvest.				
European Corn Borer (sweet corn)	2 3/4 to 3 1/4 pints					
European Corn Borer (field corn)	2 3/4 pints	Make application 10 days after eggs hatch or when 75% of the plants show evidence of feeding in leaf whorls. Repeat 7 to 10 days later if new evidence of feeding is observed. For control of second brood borers apply when eggs begin to hatch. Do not apply within 14 days of harvest.				
Southern Corn Borer (second to third generation)	2 3/4 pints	Apply when 10% of the plant population have egg masses. Repeat at 5 to 10 day intervals as needed. Do not apply within 14 days of harvest.				
Beans (Dry, green)	Lycus Bugs	1 1/2 pints	Do not apply to corn during the pollen-shed period if bees are visiting the treatment area.			
	Mites					
	Darling Ground Beetle					
	Beetles					
	Mexican Bean Beetle					
	Russet Mite					
	Serpentine Leafminer					
	Sugarbeet Webworm					
	Thrips					
	Tomato Fruitworm					
	Two-Spotted Mite					
	Soybeans			Climbing Cutworm	1/2 pint	Use restricted to the southern states. Do not make more than 2 applications during the growing season. Do not harvest or graze or feed from treated fields to livestock with 14 days after the last application.
				Stinkbugs		
				Vegetarian Caterpillar		
				Aphids		
Blister Beetle						
Cotton Bollworm						
Garden Webworm						
Mexican Bean Beetle						
Three Corned Alfalfa Hopper						
Two-spotted Spider Mite						
Cabbage Looper		2 pints				
Corn Earworm						
Green Cloverworm						
Fall Armyworm (up to third instar)		2-2 3/4 pints				
Tomatoes		Potato Aph	1 1/2 to 1 3/4 pints	Do not apply within 21 days of harvest.		
	Serpentine Leafminer					
	Southern Armyworm					