

PROFESSIONAL DDVP No. 4 INSECTICIDE EMULSIBLE CONCENTRATE

(CONTAINING 4.0 LBS. DDVP INSECTICIDE PER GALLON)

ACTIVE INGREDIENTS
 2,2-Dichlorovinyl Dimethyl Phosphate
 Related Compounds
 Petroleum Hydrocarbons
INERT INGREDIENTS

BY WEIGHT
 49.9%
 3.1%
 45.0%
 10.0%
 100.0%

GENERAL DIRECTIONS

DDVP Insecticide Emulsible Concentrate (4 pounds per gallon) is a specially prepared concentrate for use in conventional application equipment.

USE AND DIRECTIONS FOR APPLICATION

USE: Household, Restaurants, Theaters, Food Processing Plants, Industrial Plants, Warehouses.

INSECT: Flies, Mosquitoes, Wasps, Gnats, Flying Moths, Cockroaches, Ants, Spiders, Silverfish.

DOSAGE ACTUAL INSECTICIDE: 2 Pints to 25 Gals. of Water (0.5% Actual DDVP).

APPLICATION: Apply as a coarse wet Spray according to Standard PCO Practices.

DIRECTIONS FOR APPLICATION: TO BE APPLIED BY PEST CONTROL OPERATORS ONLY. Apply with low pressure sprayer to localized areas insects may infest. Treat infested areas around baseboards, cracks, walls, door and window frames, and localized areas of floors. Do not apply to areas which will be contacted frequently by children and do not allow them in treated areas until surfaces are dry. Do not contaminate food, water, food containers, or cooking utensils. Do not treat animals.

PRECAUTIONS IN USING

This insecticide concentrate and the diluted emulsion are poisonous. Therefore, precautions must be observed when handling them. (See antidote and warnings below). Store the concentrate in a safe place. Children or animals must not be able to reach or handle the concentrate or diluted emulsion. It is recommended that only the amount of diluted emulsion be prepared from the concentrate that will be used immediately. Mix the concentrate into water outdoors or in a well ventilated room. Wash off the outside of the bottle of concentrate in running water before storing away. Do not contaminate lakes or streams. Do not contaminate feed, water or food stuffs. *Handwritten:* This product is toxic to fish.

This insecticide concentrate is poisonous if swallowed, inhaled or absorbed through the skin or eyes. Do not get on skin or eyes. Do not breathe fumes. Avoid prolonged exposure to fumes. Wear natural rubber gloves, protective clothing and goggles when handling the concentrate to prepare emulsions. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for DDVP protection. Keep all unprotected persons out of operational areas. In case of contact, wash immediately with soap and water. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

KEEP OUT OF REACH OF CHILDREN

ANTIDOTE & POISON &

ANTIDOTE AND FIRST AID TREATMENT

ATROPINE is the emergency antidote for DDVP Insecticide poisoning. Consult your physician about obtaining an adequate supply of 1-100 grain atropine tablets for emergency use. Call a **PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.**

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). **NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.**

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.

IF WARNING SYMPTOMS APPEAR: Administer two 1-100 grain atropine tablets immediately. **NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR.** (See **WARNING SYMPTOMS** below).

Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.

NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine should be injected at 10 minute intervals, in doses of 1 to 2 milligrams until complete atropinization has occurred.

Atropine is contraindicated. Clean chest by postural drainage. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.

USE: Household, Restaurants, Theaters, Food Processing Plants, Industrial Plants, Warehouses.

INSECT: Flies, Mosquitoes, Wasps, Gnats, Flying Moths, Cockroaches, Ants, Spiders, Silverfish.

DOSAGE ACTUAL INSECTICIDE: 2 Pints to 25 Gals. of Water (0.5% Actual DDVP).

APPLICATION: Apply as a coarse wet Spray according to Standard PCO Practices.

INSTRUCTIONS FOR APPLICATION: TO BE APPLIED BY PEST CONTROL OPERATORS ONLY. Apply with low pressure sprayer to localized areas insects may infest. Treat infested areas around baseboards, cracks, walls, door and window frames, and localized areas of floors. Do not apply to areas which will be contacted frequently by children, and do not allow them in treated areas until surfaces are dry. Do not contaminate food, water, food containers, or cooking utensils. Do not treat animals.

PRECAUTIONS IN USING

The concentrate and the diluted emulsion are poisonous. Therefore, precautions must be observed when handling them. (See antidote and warnings below). Store the concentrate in a safe place. Children or animals must not be able to reach or handle the concentrate or diluted emulsion. It is recommended that only the amount of diluted emulsion be prepared from the concentrate that will be used immediately.

Mix the concentrate into water outdoors or in a well ventilated room. Wash off the outside of the bottle of concentrate in running water before storing away. Do not contaminate lakes or streams. Do not contaminate feed, water or food stuffs. *This product is toxic to fish.*

This insecticide concentrate is poisonous if swallowed, inhaled or absorbed through the skin or eyes. Do not get on skin or eyes. Do not breathe fumes. Avoid prolonged exposure to fumes. Wear natural rubber gloves, protective clothing and goggles when handling the concentrate to prepare emulsions. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for DDVP protection. Keep all unprotected persons out of operational areas.

In case of contact, wash immediately with soap and water. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

KEEP OUT OF REACH OF CHILDREN

NOT A POISON

ANTIDOTE AND FIRST AID TREATMENT

ATROPINE is the emergency antidote for DDVP Insecticide poisoning. Consult your physician about obtaining an adequate supply of 1/100 grain atropine tablets for emergency use. **CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.**

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). **NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.**

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.

IF WARNING SYMPTOMS APPEAR: Administer two 1/100 grain atropine tablets immediately. **NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR.** (See WARNING SYMPTOMS below).

Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.

NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine should be injected at 10 minute intervals, in doses of 1 to 2 milligrams until complete atropinization has occurred. Atropine is contraindicated. Clean chest by postural drainage. Observe patient continuously for 48 hours.

Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.

NET CONTENTS 1 GALLON

W. H. BACH PESTICIDE CORP.

3111 PEACHTREE ROAD, N. E.
ATLANTA, GEORGIA 30319

USDA REG. NO. 6754-14

INDUSTRIAL AND BACTERIOLOGICAL LAB CONTROL

EP 3441

6754-14

Jan 16 1968