

Jan 26, 1965 4 (1421-183)  
Comments  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 6754-16

# MALATHION FLY BAIT

WITH D. D. V. P.  
**Attracts and Kills  
Resistant House Flies**

## ACTIVE INGREDIENTS:

Malathion (0, 0-Dimethyl Dithiophosphate  
of Diethylmercaptosuccinate ..... 1.0%  
0, 0 Dimethyl 2, 2 Dichloro Vinyl Phosphate ..... .5%

INERT INGREDIENTS ..... 98.5%

**CAUTION: READ CAUTION STATEMENT  
KEEP OUT OF REACH OF CHILDREN**

## DIRECTIONS

Apply Fly Bait daily while flies are abundant. Sprinkle thinly in strips on floor and other places where flies gather, but not where it will contaminate feed or food. Usually, about 1 oz. (2 tbsp.) is needed for each 500 sq. ft. depending on the number of flies. If they are extremely numerous, several times this quantity may be necessary. Fly Bait is especially recommended for use in barns, stables, pens, feed rooms, poultry houses, kennels, manure piles, garbage cans, and other outside areas. In dairy barns, treat the feed room floor and the barn floor, but do not feed the floor sweepings to livestock. Repeat as needed. In poultry houses where poultry is in cages, use Fly Bait on walkways in and around the building, including feed rooms and egg houses. Where poultry is on ground or litter, apply only outside building or in yards where poultry cannot feed on Fly Bait.

To kill fly maggots, apply Fly Bait evenly and thoroughly on manure at the rate of 1 to 1½ lbs. per 100 sq. ft. (2 to 2½ oz. per sq. yd.). If in small mounds, apply on sides as well as top. Repeat in 5 to 7 days, and afterwards as needed.

TO MAKE A LIQUID BAIT: Dissolve in water at rate of 1 pound per gallon. Spray solution in coarse spray and apply to point of run-off. Apply daily until fly population is under control. Repeat applications as necessary. Do not use as a spray in dairy barns.

CAUTION: Harmful if swallowed or absorbed through the skin. Avoid skin contact. Wash thoroughly with soap and water after using. Keep container out of reach of children and animals. Avoid contamination of food, feed, and utensils used for food or feed. Do not use inside homes or where milk and other foods are stored or processed. Repeated or careless use may cause Cholinesterase inhibition. Atropine is antidotal.

U.S.D.A. REG. NO. 6754-16

NET WEIGHT      POUND

ACCEPTED WITH COMMENTS