

ED

USE DIRECTIONS

PESTS Indoors: General Information: Use ORKINBAN 2E insecticide as a spot treatment to control cockroaches, ants, crickets, firebrats, silverfish, spiders and brown dog ticks in and around residential, institutional and commercial buildings including homes, apartment buildings, hospitals, stores and warehouses. Dilute ORKINBAN 2E with water or with a suitable base oil to make a 0.25% or 0.5% chlorpyrifos spray as shown in the following SPRAY DILUTION CHART. Apply according to directions given for pests listed. Repeat the treatment as needed.

SPRAY DILUTION CHART

Gallons of Spray Mixture Desired	Fluid Ounces of ORKINBAN 2E to Use	
	0.25% spray	0.50% spray
Water-base spray:		
1	1 1/3	? 2/3
3	4	3
5	6 2/3	13 1/3
10	13 1/3	26 2/3
Oil-base spray:		
1	1	2
3	3	6
5	5	10
10	10	21

*Note: 1 fluid ounce = 2 tablespoonsful or 6 teaspoonsful
 **Calculated on basis of deodorized kerosene

Note: Where quick "knockdown" and/or "flushing action" is desired, dichlorvos (DDVP) or pyrethrins may be added to the spray mixture containing ORKINBAN 2E insecticide. For best results, use dichlorvos at a concentration of 0.25% in both water and oil-base sprays; use pyrethrins at 0.25% in water-base and 0.05% in oil-base sprays. Consult labeling for the dichlorvos or pyrethrins product used to determine the amount needed to give the desired spray concentration. Also read and observe all use and handling precautions for these products.

Cockroaches (including strains resistant to certain other insecticides). **Ants, Crickets, Firebrats, Silverfish, Spiders:** Use a 0.25% spray to control light infestations and a 0.5% spray to quickly reduce heavy infestations. Apply as a coarse spray or with a paint brush to localized areas where these pests are found or normally occur, including dark corners of rooms and closets; cracks and crevices in walls; along and behind baseboards; beneath and behind sinks, stoves, refrigerators and cabinets, around plumbing and other utility installations. For ants apply to ant trails, also around doors and windows and wherever else these pests may find entrance. Note: A period of 7 to 10 days is normally required for maximum effect on cockroaches.

Brown Dog Ticks: Thoroughly apply a 0.5% spray as a spot treatment to infested areas such as pet beds and resting quarters; nearby cracks and crevices; along and behind baseboards, window and door frames and localized areas of floor and floor covering where these pests may be present. Old bedding of pets should be removed and replaced with clean, fresh bedding after treatment. Do not treat pets with this product.

Warning (Pests Indoors): For Spot Application Only. Do not use in poultry houses, other animal buildings or in edible products areas of food processing plants, restaurants, or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed. Do not allow spray to contact food, feedstuffs and water supplies. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated by application of this product. Do not allow children or pets to contact treated surfaces until spray has dried. Remove pets and cover fish bowls (tanks) before spraying.

APPLICATIONS WITHIN FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS Limited to Crack and Crevice Treatment Only—Apply a small amount of material directly into cracks and crevices such as expansion joints between different elements of construction or between equipment bases and the floor, wall voids or hollow equipment legs where cockroaches, ants, brown dog ticks, silverfish, spiders may hide. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces. Orkinban 2E is not recommended for use in conduits, motor housings or switch and junction boxes because of shock hazard.

Application of this product in the food areas of food handling establishments, other than as a crack and crevice treatment, are not permitted.

PROFESSIONAL ORKINBAN 2E

Contains DURSBAN* Insecticide

CONTROLS PESTS OF HOUSEHOLDS (COCKROACHES, ANTS, BROWN DOG TICKS, SILVERFISH, SPIDERS), TURF (CHINCH BUGS, SOD WEBWORMS, ANTS, BROWN DOG TICKS, CUTWORMS, EARWIGS, CRICKETS, GRASSHOPPERS, HYPERODES WEEVIL) AND ORNAMENTAL PLANTS (BAGWORMS, GRASSHOPPERS, LEAFHOPPERS, SPITTLEBUGS, MEALYBUGS, WHITEFLIES, EXPOSED THRIPS, MITES, CUTWORMS). ALSO RECOMMENDED FOR MOSQUITO CONTROL AND AREA CONTROL OF TICKS.

To Be Applied Only By or Under the Supervision of Pest Control Operators, Public Health Organizations, Mosquito Abatement Districts and Other Trained Personnel Responsible for Insect Control Programs

ACTIVE INGREDIENTS:

Chlorpyrifos [O,O-diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate] 22.5%
 Aromatic petroleum derivative solvent 70.9%

INERT INGREDIENTS:

Contains 2 pounds of chlorpyrifos per gallon

*Dursban is Trademark The Dow Chemical Company—U. S. Patent #3,244,586.

WARNING

**KEEP OUT OF REACH OF CHILDREN
 MAY BE FATAL IF SWALLOWED
 MAY BE ABSORBED THROUGH SKIN
 MAY BE IRRITATING TO THE EYES AND SKIN
 DO NOT GET IN EYES OR ON SKIN**

Do Not Wear Contaminated Clothing. Wash All Contaminated Clothing With Soap and Hot Water Before Reuse. Avoid Breathing Spray Particles. Handle in Ventilated Area.

Do not use in edible products areas of food processing plants, restaurants, or other areas where food is commercially prepared or processed, except when using the Crack and Crevice method of treatment.

Do not use in serving areas while food is exposed. Flush contaminated Eyes and Skin with plenty of Water and for Eyes Get Medical Attention.

Note to Physician: Orkinban 2E Concentrate is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

Keep Container Closed and Away from Heat or Open Flame

COMBUSTIBLE LIQUID

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment, or disposal of wastes. Keep out of lakes, streams or ponds. Apply this product only as specified on this label. Do not apply where runoff is likely to occur.

Do not reuse empty container. Destroy by perforating or crushing and burying in a location isolated from water. Do not use, pour, spill or store near heat or open flame.

GENERAL USE PRECAUTIONS

This product is highly toxic to bees exposed to direct treatment or residues on plants. Protective information may be obtained from your Cooperative Agricultural Extension Service. ORKINBAN 2E also is toxic to fish, wildlife and birds.

Do not contaminate water by cleaning equipment or disposal of wastes. Promptly crush empty containers and burn them and wastes in locations away from domestic water supplies.

Use this product only as specified on this label.

EXCLUSIVE FOR



EXTERMINATORS

EPA Reg. No. 6754-37

EPA Est. No. 6754-GA-1

DETTELBACH

Pesticide

CORPORATION

4111 PEACHTREE ROAD, N.E. - ATLANTA, GEORGIA 30319

PESTS OUT-OF-DOORS: General Information: Use ORKINBAN 2E for control of certain pests of lawns and ornamental plants, for area control of ticks and for mosquito control. Consult the following directions for dosage and application details to control the pests listed.

Pests of Lawns and other Ornamental and Recreational Turf Grass Areas: Use ORKINBAN 2E insecticide to control ants, brown dog ticks, chinch bugs, crickets, cutworms, earwigs, grasshoppers and sod webworms (lawn moths). Mix ORKINBAN 2E with water according to directions given in the following table and apply as a coarse low pressure spray using suitable application equipment. For best results, the lawn should be moist at time of treatment, and care should be taken to obtain complete and uniform coverage of pest infested areas. Spray when pests first appear, retreat when needed.

DOSAGE AND DILUTIONS FOR USE ON TURF

Area to be Treated	Amount of ORKINBAN 2E	Dilute with Water to Give
500 sq. ft.	3/4 fl. oz.	15 gallons
1,000 sq. ft.	1 1/2 fl. oz.	30 gallons
5,000 sq. ft.	7 1/2 fl. oz.	150 gallons
10,700 sq. ft.	1 pint	325 gallons
One Acre	2 quarts	1,300 gallons

To control turfgrass (Hyperodes' weevils) use twice the amount of ORKINBAN 2E shown in the above TURF table, i.e. 1 gallon diluted in 1,300 gallons of water per acre. Spray suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialist for proper time to treat in your area.

Note: To help prevent infestation of buildings by ants, crickets, earwigs and sowbugs (pillbugs), treat a band of soil 6 to 10 feet wide around and adjacent to the building, also the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use the same dosage and dilution recommended for control of these pests on turf as shown in the above table. Apply as a coarse spray to thoroughly and uniformly wet the band area.

Pests of Ornamentals: Use ORKINBAN 2E to control bagworms, cutworms, grasshoppers, leafhoppers, mealybugs, mites, spittlebugs, thrips (exposed) and whiteflies on flowers, shrubs, vines, shade and flowering trees and evergreens found to be infested with these pests. Dilute ORKINBAN 2E with water according to directions given in the following table and apply using suitable hand or power spray equipment in a manner to provide complete and uniform coverage of infested plants. Avoid overspraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals, if needed. **Note**—Do not use on azaleas, camellias, poinsettias, rose bushes or variegated ivy because of possible injury to these plants.

DOSAGE AND DILUTIONS FOR USE ON ORNAMENTAL PLANTS

Pests Controlled	Amount of ORKINBAN 2E in Water to Make		
	3 gallons	10 gallons	100 gallons
Bagworms			
Grasshoppers			
Mites	0.5 fl. oz.	1.6 fl. oz.	1 pint
Spittlebugs			
Whiteflies			
Cutworms			
Mealybugs			
Leafhoppers	1 fl. oz.	3.2 fl. oz.	1 quart
Thrips (exposed)			

*Equivalent to 4 ounces of chlorpyrifos per 100 gallons

**Equivalent to 8 ounces of chlorpyrifos per 100 gallons

Note: Mosquitoes coming to rest on areas treated for control of turf and ornamental plant pests will be killed for varying periods of time after treatment, depending on exposure of treated areas to weathering conditions.

Warning (Lawn and Ornamental Plant Pests): Keep children and pets off treated areas until spray has dried. Keep out of lawn and garden fishpools and other bodies of water. Fish may be killed if their waters are contaminated with this product. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated.

Area Control of Ticks: Use ORKINBAN 2E to control ticks infesting non-crop land areas such as roadsides, foot paths and trails, picnic and camping sites, parks and other recreational areas where these pests are present and create a nuisance or possible public health problem. Apply ORKINBAN 2E in water at the rate of 1 pint per acre (equivalent to 1 fluid ounce per 2,700 sq. ft.) using a hydraulic sprayer, mist applicator, knapsack sprayer or other suitable hand or power spray equipment. Treat low underbrush, grassy areas, weeds and ground surface and debris using enough spray volume to obtain thorough coverage, usually 40 to 100 gallons per acre.

Warning (Area Control of Ticks): Do not allow public use of treated areas during application or until spray has dried. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff into lakes, streams, ponds or other bodies of water is likely to occur. Fish and crustaceans may be killed if their waters are contaminated with this product.

Mosquitoes: Apply ORKINBAN 2E to mosquito infested, non-crop areas according to the conditions and rates shown in the following table. Mix ORKINBAN 2E with either water or oil, such as kerosene or No. 2 fuel oil, using sufficient spray volume to give uniform coverage of area treated. Apply the spray using suitable application equipment such as hand and power ground sprayers, mist applicators, and aerial spray equipment.

CONDITIONS AND RATES TO USE FOR MOSQUITO CONTROL

Condition of Area to be Treated	Amount to Use per Acre		Number Acres 1 Gallon of ORKINBAN 2E Will Treat	
	ORKINBAN 2E (fluid ounces)	Chlorpyrifos (pounds)		
Larval Control	None to medium vegetative cover	0.8 to 1.6	0.0125 to 0.025	160 to 80
	Medium to heavy vegetative cover	1.6 to 3.2	0.025 to 0.05	80 to 40
Adult Control	Light to medium vegetative cover	1.6	0.025	80
	Medium to heavy vegetative cover	3.2	0.05	40

*The higher dosage will also give faster knockdown and longer residual control where there is little or no vegetative cover.

Warning (Mosquito Control): ORKINBAN 2E is toxic to shrimp, crabs and fish. Do not apply to water containing crustaceans and fish. Consult your State Fish and Game Department before using this product for mosquito control.