

PROFESSIONAL DDVP No. 2 INSECTICIDE EMULSIFIABLE CONCENTRATE

(CONTAINING 2.0 LBS. DDVP INSECTICIDE PER GALLON)

ACTIVE INGREDIENTS	By Weight
2, 2-Dichlorovinyl Dimethyl Phosphate	22.0%
Related Compounds	1.7%
Heavy Aromatic Naphtha	71.3%
INERT INGREDIENTS	5.0%
	100.0%

DANGER KEEP OUT OF REACH OF CHILDREN

PRECAUTIONS IN USING

Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed, except when using the Crack and Crevice method of treatment. Do not use in serving areas while food is exposed.

DANGER: This insecticide concentrate and the diluted emulsion are poisonous. Therefore precautions must be observed when handling them. (See antidote and warnings on side panel.) Store in a safe place. Children or animals must not be able to reach or handle the concentrate or diluted emulsion. It is recommended that only the amount of diluted emulsion be prepared from the concentrate that will be used immediately.

Mix the concentrate into water outdoors or in a well ventilated room. Wipe off the outside of the bottle of concentrate in running water before storing away.

Do not contaminate lakes or streams. Do not contaminate feed, water or food stuffs. This insecticide is poisonous if swallowed, inhaled or absorbed through the skin or eyes. Do not get on skin or eyes. Do not breathe fumes. Avoid exposure to fumes. Wear natural rubber gloves, protective clothing and goggles when handling the concentrate to prepare emulsions. Wear a mask or respirator of a type passed by the U. S. Bureau of Mines for DDVP protection. Keep all unprotected persons out of operational areas.

In case of contact, wash immediately with soap and water. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse. Dispose of empty containers, spillage, and spray wastes by burying in non-crop lands.

NET CONTENTS

GALLONS

EPA Reg. No. 6754-50

EPA Est. No. 6754 GA-1

EXCLUSIVE FOR

ORKIN

EXTERMINATORS

DETTELBACH

Pesticide

CORPORATION

4111 PEACHTREE ROAD, N.E. - ATLANTA, GEORGIA 30319

FOR INDOOR USE ONLY

GENERAL DIRECTIONS

DDVP #2 Insecticide Emulsifiable Concentrate (2 pounds per gallon) is a specially prepared concentrate for use in conventional application equipment.

USE AND DIRECTIONS FOR APPLICATION

USE: Houses, Restaurants, Theaters, Food Processing Plants, Industrial Plants, Warehouses.

INSECT: Flies, Mosquitoes, Wasps, Gnats, Small Fly (e. g. Midge), Cockroaches, Ants, Spiders, Silverfish, Fleas, Brown Dog Ticks.

DOSAGE ACTUAL INSECTICIDE: 4 Pints to 25 Gallons of Water (1:5 to 1:10 Actual DDVP).

APPLICATION: Apply as a coarse wet spray according to Standard POC practices.

DIRECTIONS FOR APPLICATION: TO BE APPLIED BY PEST CONTROL OPERATORS ONLY. Apply with low pressure sprayer to exposed areas where insects may infest. Treat infested areas around doorways, cracks, walls, floor and window frames, and other areas of interest.

FOOD HANDLING ESTABLISHMENTS (Other than private residences in which food is held, prepared, packaged and/or served). For control of roaches, ants, spiders, silverfish, crickets and cockroaches.

NON-FOOD AREAS

Apply to areas of insect infestation, for example, lawns, lawns, lawns, and lawns. (See side panel for more information.) Do not use in areas where children or animals may be present.

Apply as a coarse wet spray by means of a low pressure sprayer to base of walls, around window and door surfaces behind and beneath sinks, tubs, toilets, showers, and other areas where pests hide or through which they may enter.

FOOD AREAS

Includes the areas where food is held, frozen, raw, packaging, canning, bottling, washing, drying, preparing, packaging, slicing, cooking, broiling, roasting, etc. Includes food processing systems, mills, canneries, etc. **Limited to Crack and Crevice Treatment Only!**

Apply as a coarse wet spray by means of a low pressure sprayer to small amount of crack and crevices where cockroaches, ants and spiders hide or through which they may enter the building. Includes expansion joints, cracks between equipment and floors, and openings leading to outdoors and basements in walk-in equipment, etc. and bases. Care should be taken to avoid getting the product into exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces. Do not use DDVP No. 2 INSECTICIDE EMULSIFIABLE CONCENTRATE in a kitchen, dining room, or switch and junction boxes because of shock hazard.

APPLICATIONS OF DDVP NO. 2 INSECTICIDE EMULSIFIABLE CONCENTRATE IN FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS, OTHER THAN AS A CRACK AND CREVICE TREATMENT ARE NOT PERMITTED.

6

DANGER KEEP OUT OF REACH OF CHILDREN

POISON

ATROPINE is the emergency antidote for DDVP insecticide poisoning. 2-PAM is also antidotal and may be administered in conjunction with atropine. Consult your physician about obtaining an adequate supply of 1/100 grain atropine tablets for emergency use. Call a PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the material, remove all contaminated clothing and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.

IF WARNING SYMPTOMS APPEAR: Administer two (2) 1/100 grain atropine tablets immediately. NEVER ADMINISTER ATROPINE UNLESS SYMPTOMS APPEAR. (See WARNING SYMPTOMS below).

Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.

NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10 minute intervals, in doses of 1 to 2 milligrams until complete atropinization has occurred.

Microhine is contraindicated. Clean chest by percutaneous drainage. Observe patient continuously for 48 hours.

Repeated exposure to cholinesterase inhibitors may, without warning, cause pronounced susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until tolerance for cholinesterase re-education has been allowed as determined by blood test.