

FRONT PANEL

MICROBAN

HOSPITAL SPRAY PLUS

MILDEWCIDE • BACTERICIDE • FUNGICIDE
DEODORANT AND DISINFECTANT (Anti-Microbial)

FOR INSTITUTIONAL, HOUSEHOLD, HEALTH CARE, FARM PREMISE USES AND ANIMAL QUARTERS

Microban is a Broad Spectrum Disinfectant Against most Bacteria, Fungus,
Mildew, Odors, Caused by Bacteria and Germs

ACTIVE INGREDIENTS

o-Phenylphenol	0.22%
Diisobutylphenoxyethoxy ethyl dimethyl benzyl ammonium chloride monohydrate	0.70%
INERT INGREDIENTS	99.08%
TOTAL	100.00%

E P.A. Reg. 6768-

E P.A. EST 6768-PA 1

KEEP OUT OF REACH OF CHILDREN

WARNING

Statement of Practical Treatment

If Inhaled: Remove patient to fresh air. If not breathing, give artificial respiration. If breathing is difficult, give oxygen. Get immediate medical attention.

If Swallowed: Drink promptly a large quantity of water. Do not induce vomiting. Avoid alcohol. Get medical attention.

If In Eyes Or On Skin: Immediately flush with plenty of water. Get medical attention if irritation persists.

See other precautions on side panel

Net Contents 1 PT. 1 QT. 1 GAL. 5 GAL. 55 GAL.

Mfg by: MICROBAN GERMICIDE COMPANY P.O. Box 777, Braddock, PA 15104

SHAKE WELL BEFORE AND DURING USE

U.S. Patent No. 3,458,629

ACCEPTED

12 APR 1993

Order on File & License, Imports, and
Residue Act. of 1990 for the
pesticide registered at
EPA Reg. No. 6768-14

BEST AVAILABLE COPY

GENERAL METHOD OF APPLICATION

Surfaces and objects must be cleaned of gross filth with suitable detergent before treatment. Spray surfaces and articles until thoroughly wet, use enough so that treated surfaces and objects remain wet for at least 10 minutes. May be applied with a sponge to impervious covers and surfaces, allowing to dry for 10 minutes. For fogging and other large volume applications use appropriate spraying equipment, protective clothing, gloves and respirator. Heavily treated spaces should be adequately ventilated and not re-entered for at least one hour after treatment.

AS A MILDEWICIDE, FUNGICIDE AND BROAD SPECTRUM DISINFECTANT: (Kills most gram negative and gram positive bacteria and influenza virus). In hospitals, operating rooms, emergency rooms, lobbies, hallways, patient rooms, wards, washrooms, scrub stations, mop rooms, morgues, pathology labs, nurse's stations, laboratories, veterinary clinics, and other health care facilities; in schools, hotels, motels, restaurants, offices, homes, and industrial buildings.

For use on pre-cleaned hard non-porous surfaces such as tiled porcelain, waste receptacles, compactors, on, in and around rest room fixtures, toilet seats, locker rooms, telephones, door knobs, hospital apparatus and equipment, furniture, empty containerized storage units. For athletes foot fungus on floors, shower stalls, rest rooms. Follow method of application specified above. Do not use on medical devices.

AS A BACTERIOSTAT, FUNGISTAT, AND DEODORIZING AGENT

Apply Microban Hospital Spray Plus as directed to floors, walls, ceiling that have been damaged by smoke, fire, floods and sewage backups.

cleanup operations in areas such as attics, wall voids, basements, areas above suspended ceilings and similar inaccessible spaces in building. Spray avian, rodent and other animal and human wastes and carcasses as directed above, prior to removal and disposal. Treat spaces and surfaces as specified in method of application.

For use as a bacteriostat, fungistat, and deodorizing agent on carpeting found in schools, hotels, motels, homes, shelters, commercial, industrial and public buildings. After flood or water damage, vacuum or extract as much water as possible before application. Apply full strength with sprayer until moist to carpet nap and rake or brush in. To treat the backing and padding a fogger is recommended. Allow to remain until dry. For best results carpets should be cleaned before application.

TO DISINFECT FARM PREMISES, POULTRY HOUSES, ANIMAL PENS AND VEHICLES

Remove all animals and feed from premises, vehicles, and enclosures. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities and fixtures occupied or traversed by animals. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate all surfaces with the recommended disinfecting solution for a period of 10 minutes. Immerse all halters, ropes, and other types of equipment used in handling and retraining animals, as well as forks, shovels, and scrapers used for removing litter and manure. Ventilate buildings, cars, boats, and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set, or dried. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse. Apply Microban Hospital Spray Plus full strength as directed in general method of application until thoroughly wet. Allow all surfaces to dry thoroughly, ventilate enclosed spaces adequately. Do not house animals in treated areas until thoroughly dry and free from odors. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before reuse.

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: May be harmful if swallowed (inhaled or absorbed through the skin). Do not breathe vapors (dust or spray must). Do not get in eyes, on skin or on clothing. May cause eye (and skin) irritation.

STORAGE AND DISPOSAL

Storage: Keep from freezing. Store in ventilated areas. Do not contaminate water, food or feed by storage and disposal. Dispose left overs in landfill approved for pesticides.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Large containers (55 gallon drums and 5 gallon pails): Triple rinse (or equivalent) all containers and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by procedures approved by state and local authorities. Small container (1 gallon or less): Rinse thoroughly and discard with trash.

TEST FABRIC OR SURFACE IN AN INCONSPICUOUS AREA FOR COLOR FASTNESS OR ADVERSE REACTION

ACCEPTED
1 2 APR 1993
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended for the pesticide registered under:
EPA Reg. No. 4763-17

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Microban Hospital Spray Plus should be applied with hand or power operated spray, foggers, automated metering systems or gravity drip dispensers as a broad spectrum bactericide, mold and mildewcide, fungicide and deodorizing agent to previously cleaned surfaces, spray areas until thoroughly moist, giving special attention to cracks and crevices. Allow ten to twenty minutes for drying. It is not necessary to wipe the sprayed surface. Microban should not stain those materials not stained by water.

Microban Hospital Spray Plus when used as directed is effective in decontaminating bedding, mattresses, textiles, drapes, upholstered furniture, rugs, carpets and in storage areas to control mold, mildew, fungus and odors wherever they are a problem.

- Pest Control Service
- Odor Control Service
- Carpet Cleaning Service
- Water Damage Service
- Restoration Service
- Disinfection Service
- Waste-Removal Service
- Second hand furniture stores
- Trash Chutes and Garbage Rooms
- Waste Compactors
- Storage areas and Basements
- Condensation collection and Drainage areas
- Laundry receiving stations
- Air ducts
- Movie booths
- Bowling alleys
- Antique shops
- Refuse trucks
- Exercise rooms
- Shoes & boots
- Veterinaries
- Upholsterers shops

© 1984 MGC

SHAKE WELL BEFORE AND DURING USE

