

KILLS: Dandelions, Chickweed, Knotweed, Plantains, Henbit, Spruge and many other broadleaf weeds.

HOW TO MIX

1. For use on St. Augustine grass, bentgrass, buffalo grass and grass seedlings, mix 1 ounce in 1 gallon of water and uniformly spray 400 sq. ft., for example an area 10 feet wide and 40 feet long. If necessary, repeat treatment in 2 to 3 weeks, if hard to kill weeds persist.
2. For use on certain other grasses such as Bermuda, bahia, bluegrass, centipede, fescue and rye. Mix 1½ ounce in 1 gallon of water and spray uniformly over 400 sq. ft. On hard to kill species, or under poor growing conditions, or on older weeds, mix 2 ounces in 1 gallon of water and spray on 400 sq. ft. with special attention to thorough coverage of the hard to kill weeds.

DIRECTIONS FOR USE

GENERAL INFORMATION: St. Augustine Grass Broadleaf Herbicide is specially formulated for maximum safety to seedling grasses and 2,4-D sensitive grasses such as St. Augustine grass, bentgrass and buffalo grass. Sprays should be applied when broadleaf weeds are small and tender for best results. If weeds are older and hard to kill, the higher rates recommended should be used. Avoid spraying St. Augustine grass under conditions of high temperature and moisture when the St. Augustine is growing rapidly. Under these conditions, or in cases of accidental overdose, some slight temporary discoloration of the St. Augustine foliage might occur.

To insure weeds are growing and able to absorb the herbicide irrigate the lawn 2 or 3 times prior to spraying. Do not water for 24 hours after treatment. Although spray works faster with temperatures above 70°F, sprays can be applied at any time plants are growing. Spray when air is calm to avoid drift. Spray will injure or kill clover, dichondra and most ornamental and vegetable crops.

Amount of water in spray is not important, except to evenly distribute the chemical. If directions call for 1 ounce in 1 gallon on 400 sq. ft. it may be more convenient to use 1 oz. in 2 gallon on 400 sq. ft. depending on the sprayer. Generally large volume of water result in more uniform coverage.



ST. AUGUSTINE GRASS BROADLEAF HERBICIDE Contains TRIMEC (R) Herbicide

ACTIVE INGREDIENTS:

*Dimethylamine salt of 2,4-dichlorophenoxyacetic acid 3.23%

**Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid 10.59%

***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid) 1.28%

INERT INGREDIENTS 84.90%

TOTAL 100.00%

*2,4-dichlorophenoxyacetic acid equivalent 0.235 pounds per gallon or 2.69%

**2-(2-methyl-4-chlorophenoxy) propionic acid equivalent

0.764 pounds per gallon or 8.76%

***3,6-dichloro-o-anisic acid equivalent 0.094 pounds per gallon or 1.07%

CAUTION: KEEP OUT OF REACH OF CHILDREN

See side panels for additional cautions and information

U. S. Patent No. 3,284,186

EPA Reg. No. 6720-178



MANUFACTURED BY

Southern MILL CREEK PRODUCTS COMPANY, Inc.
TAMPA, FLORIDA

