

**COLD DISINFECTION:** For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into solution containing 1½ ounce of NEO-SAN per gallon of water for a minimum time of ten minutes.

**RESTAURANT AND BAR RINSE:** Dishes, glassware, silverware, cooking utensils, wash with soap or synthetic detergent; rinse thoroughly and immerse in a sanitizing solution containing 1 ounce NEO-SAN to 4 gallons of water. **No terminal potable water rinse required.**

**DANGER: Keep Out of Reach of Children.** Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

**FIRST AID:** In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician immediately.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Rinse empty container thoroughly with water and discard.

**NET CONTENTS:**

# NEO-SAN

DISINFECTANT

DEODORIZER

SANITIZER

## ACTIVE INGREDIENTS:

n-Alkyl (60%C14, 30%C16, 5%C12, 5%C18)  
dimethyl benzyl ammonium chlorides . . . . . 5%  
n-Alkyl (68%C12, 32%C14) dimethyl  
Ethylbenzyl ammonium chlorides . . . . . 5%  
Ethyl Alcohol . . . . . 2.5%

INERT INGREDIENTS: . . . . . 87.5%  
TOTAL . . . . . 100.0%

NEO-SAN is a concentrate and should be diluted before using.

## DANGER

KEEP OUT OF REACH OF CHILDREN

See other precautionary statements on left side panel.

EPA Reg. No. 6811-37

**USE DIRECTIONS: HOUSEHOLD**—For sanitizing and disinfecting floors, walls and inanimate hard surfaces. For Schools, Homes, Locker Rooms, Garbage Pails, Sink Tops, Corridors, Classrooms, Offices and Shower Stalls. Apply solution with mop or cloth. Sanitize with 1 ounce NEO-SAN to 4 gallons of water. Disinfect with 3 ounces NEO-SAN to 5 gallons of water. Kills Staphylococcus Aureus and Salmonella Choleraesuis.

**HOSPITALS:** For floors, walls and other hard surfaces. In Nursing Homes, Institutions, as well as Sink Tops, Garbage Pails, Telephones and Restrooms, use NEO-SAN at 3½ ounces per 5 gallons of water. At this level NEO-SAN will be effective against Pseudomonas Aeruginosa PRD-10.

**FOOD PROCESSING EQUIPMENT:** For sanitization of previously cleaned food processing equipment and food utensils dilute 1 ounce of NEO-SAN per 4 gallons of water to provide 200 ppm of active quaternary. **For this type of use a terminal rinse with potable water is required.**

**DAIRIES:** To sanitize dairy equipment, first clean and rinse the equipment thoroughly. Then apply sanitizing solution containing 1 ounce NEO-SAN to 4 gallons of water (200 ppm). At this level, **No Potable Water Rinse** is required. Follow recommendations of local Health Board.

ACCEPTED  
APRIL 23, 1973  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND ROENTICIDE ACT  
FOR ECONOMIC POISON REGISTRATION  
NO. 6811-37, SUBSISTENT  
TO ATTACHMENTS.

MERRILL

RESEARCH

PRODUCTS CO.

DALLAS, TEXAS

29