

[6811-39]

CAUTION: Harmful if swallowed. Cover or remove fishbowls. Avoid contamination of feed and food-stuffs. Use in food processing plants only when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered or thoroughly cleaned before using. In hospitals, patients should be removed from room prior to treatment. Room should be ventilated for 2 hours after treatment. Do not return patients to room until after ventilation.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

Rinse empty container thoroughly with water and discard it.

DIRECTIONS FOR USE: REPCO-TOX IS A CONCENTRATE AND MUST BE DILUTED AS LISTED BELOW BEFORE USING.

Dilute to 0.5% by adding 1 part by weight of the 2.5% concentrate to 4 parts of petroleum distillate. Dilute to 0.25% by adding 1 part by weight of the 2.5% concentrate to 9 parts of petroleum distillate. Dilute to 0.10% (which would be used for flying insects only) by adding 1 part by weight of the 2.5% concentrate to 24 parts of petroleum distillate.

REPCO-TOX SPACE SPRAY INSECTICIDE CONCENTRATE

6811-39

ACCEPTED

ACT 30 1972

UNDER THE FEDERAL INSECTICIDE AND FUNGICIDE ACT OF 1947
REGISTERED SUBJECT TO ATTACHED COMMENTS

- *BAKERIES* *FOOD PLANTS* *RESTAURANTS*
- *SCHOOLS* *HOTELS* *MOTELS* *OFFICE BUILDINGS*
- *HOSPITALS* *FOOD HANDLING PLANTS*

Active Ingredients:

* (5-Benzyl-3-furyl) methyl 2, 2-dimethyl-3-(2-methylpropenyl) cyclopropanecarboxylate	2.500%
Related compounds	0.341%
Aromatic Petroleum Hydrocarbons	3.309%
Petroleum Distillate	93.750%
Inert Ingredients:	0.100%
	100.000%

* Cis/trans isomers ratio: max. 30% (±) cis and min. 70% (±) trans.
*U.S. Patent No. 3,465,007

CAUTION: Keep Out Of Reach Of Children. Do Not Store Near Heat Or Open Flame. See left side panel for additional cautions.

EPA REG. NO. 6811-39

FOR HOUSEFLIES, MOSQUITOES, AND GNATS:

Close all doors and windows. Point spray nozzle upward, direct the spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then retreat the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

WASPS: Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places, contacting as many insects as possible. Repeat as necessary.

FOR CRAWLING INSECTS: Roaches, earwigs, grain mites, sowbugs, spiders and centipedes: Spray thoroughly into hiding places such as cracks, openings around pipes and sinks and storage areas. Spray directly on insects where possible.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

For most effective results use in a good sprayer adjusted to deliver a fine mist.

NET CONTENTS:

MERRILL

RESEARCH PRODUCTS CO.

DALLAS, TEXAS

29