

OCTAGON PROCESS INC.

596 RIVER ROAD

EDGEWATER, N. J. 07020

6840-530-4762

**RODENTICIDE BAIT
ANTICOAGULANT, FUMARIN**

O-R-500C

NET WEIGHT: 1 LB. 8 OZS. (24 OZS.)

ACTIVE INGREDIENTS:

PERCENT

3-(alpha-acetonylfurfuryl)-4-hydroxycoumarin 0.025

INERT INGREDIENTS: 99.975

ACCEPTED WITH COMMENTS

KEEP OUT OF REACH OF CHILDREN

CAUTION

Keep away from humans, domestic animals, and pets. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses by bishydroxycoumarin.

Baits should be placed in areas inaccessible to children, pet, wildlife and domestic animals or in tamper-proof bait boxes.

Do not reuse container. Destroy when empty.

DIRECTIONS: Rats: Place 4 to 8 ounces of Rodenticide, Bait Anticoagulant in bait stations where rats feed, water or travel. Place baits along walls, in corners, and concealed locations where rats can feed unobserved. Replenish baits as necessary, and maintain stations for not less than 10 days. (Baiting may be continued indefinitely, and increased time of exposure of baits may result in better control of rats.) If bait molds, decomposes, or becomes sour, replace with fresh material.

Mice: Place tablespoonful amounts of bait at 8 to 12 foot intervals where mice feed, water or travel. Replenish baits as necessary, and maintain stations for not less than 15 days. If bait becomes moldy, sour or decomposes, replace with fresh material.

USDA REG. NO. 6830-13