

# INSECTICIDE, CHLORDANE, 72 Percent, Water Emulsifiable Concentrate

OCTAGON C515  
FEDERAL SPECIFICATION O-I-515  
(APPROX 8 LBS CHLORDANE/GALLON)

ACTIVE INGREDIENTS	Percent by weight
Technical Chlordane	72%
Petroleum Hydrocarbons	23%
INERT INGREDIENTS	5%
	100%

Equivalent to 43.2% technical 4,7-methanotetrahydrochlordane  
and 28.8% of related compounds

## WARNING! KEEP OUT OF REACH OF CHILDREN DO NOT USE WITHOUT DILUTING

**DO NOT USE IN FOOD PREPARATION AREAS OR WHERE FOOD IS EXPOSED.**

Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. When extensive use is involved, operators should wear respirators approved by the U. S. Bureau of Mines. See manual "Insect and Rodent Control" (AFM 85-7, TM 5-632, NAVDOCKS IT-Pu-2) for details.

Avoid contamination of feed and foodstuffs. In living quarters, use as strip treatment (not for complete coverage of walls).

To be applied by trained personnel only.

This product is toxic to fish and wildlife. Keep out of any body of water. Apply this product only as specified on this label.

Do not reuse empty containers. Destroy by crushing or perforate and bury in a safe place.

### DIRECTIONS FOR USE

#### RESIDUAL SPRAY IN BUILDINGS

approx. 3 percent; Add 1 part of concentrate, while stirring, to 30 parts of water. About 4 ounces of concentrate makes 1 gallon of emulsion. Apply as a coarse spray to surfaces at the rate of approximately 1 gallon per 1000 or 1 pint per 125 lineal feet to cracks, crevices and other harborages of roaches, and to runways of ants.

#### OUTDOOR SPRAY

approx. 1 percent. Add 1 part of concentrate, while stirring to 90 parts of water (about 7 ounces for 5 gallons of emulsion). Apply as coarse spray. For the control of mosquitoes, flies and sand flies, apply lightly to vegetation, tree trunks, etc. at a rate of 10 to 15 gallons per acre depending on the amount of vegetation and type of terrain to be treated.

For the control of mites (chiggers, and ticks), apply to low vegetation and to ground surface at the rate of approximately 20 gallons per acre depending on the amount of vegetation. Keep children and pets off treated areas until this material is washed into the soil and is dry. For the control of fleas, apply at rate of 10 to 20 gallons per acre around and under buildings. For the control of ants, apply at rate of 10 to 20 gallons per acre.

#### TERMITE CONTROL

Use 1 percent emulsion as prepared and described under outdoor spray. Apply in accordance with current instructions for termite control. See manual "Insect and Rodent Control" (AFM 85-7, TM 5-632, NAVDOCKS IT-Pu-2) for details.  
USDA REG. NO. 6830-15

