

# OCTAGON PROCESS INC.

596 RIVER ROAD EDGEWATER, N. J.

6840 - 530 - 4762 O - R - 500B

## RODENTICIDE, BAIT, ANTICOAGULANT, WARFARIN 0.025% OCTAGON WARFARIN 0.025%

NET WT 24 OUNCES	PERCENT
ACTIVE INGREDIENTS	BY WEIGHT
WARFARIN 3-(alpha-acetylbenzyl)-4-hydroxycoumarin	0.025%
INERT INGREDIENTS	99.975%
Dye	0.050%
Rolled Oats	90.325%
Petrolatum Liquid	4.800%
Sugar Granulated	4.800%

Total 100.000%

### CAUTION

#### KEEP OUT OF REACH OF CHILDREN

**CAUTION:** Keep away from humans, domestic animals and pets. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bis-hydroxycoumarin.

**RAT CONTROL** Place 4 to 8 ounces of Rodenticide, Bait, Anticoagulant in bait stations. Place so as to protect children, pets and livestock in locations where rats or mice feed. Replenish baits as necessary, and maintain stations for not less than 12 days. Baiting may be continued indefinitely as increased time of exposure of baits may result in better rodent control. If bait molds, decomposes, or sours, replace with fresh material.

**MOUSE CONTROL** Tablespoon amounts 1/4 to 1/2 ounce of bait should be placed at 8-12 foot intervals. A continuous supply of bait should be maintained for at least 15 days.

USDA REG NO 6830-25

**ACCEPTED**

3/30/71

UNLAWFUL TO REPRODUCE OR TRANSMIT IN ANY MANNER WITHOUT THE WRITTEN PERMISSION OF THE REGISTERED OWNER. SUBJECT TO ATTACHED COMMENTS.

**ACCEPTED**  
 3/30/71  
 UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND RODENTICIDE ACT  
 FOR ECONOMIC PEST CONTROL REGIST  
 ED UNDER NO. 6830-25  
 SUBJECT TO ATTACHED COMMENTS.

MOUSE CONTROL Tablespoon amounts (1/4 to 1/2 ounce) of bait should be placed at 8-12 foot intervals. A continuous supply of bait should be maintained for at least 15 days.  
 USDA REG NO 6830-25

**OCTAGON PROCESS INC.**  
 596 RIVER ROAD EDGEWATER, N. J.

6840-530-4762 O-R-500B

**RODENTICIDE, BAIT,  
 ANTICOAGULANT, WARFARIN 0.025%  
 OCTAGON WARFARIN 0.025%**

NET WT 24 OUNCES	PERCENT
ACTIVE INGREDIENTS	BY WEIGHT
WARFARIN 3-(alpha-acetylbenzyl)-4-hydroxycoumarin	0.025%
INERT INGREDIENTS	99.975%
Dye	0.050%
Rolled Oats	90.325%
Petrolatum Liquid	4.800%
Sugar Granulated	4.800%
	<hr/>
Total	100.000%

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**

**CAUTION:** Keep away from humans, domestic animals and pets. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bis-hydroxycoumarin.

**RAT CONTROL:** Place 4 to 8 ounces of Rodenticide, Bait, Anticoagulant in bait stations. Place so as to protect children, pets and livestock in locations where rats or mice feed. Replenish baits as necessary, and maintain stations for not less than 12 days. Baiting may be continued indefinitely as increased time of exposure of baits may result in better rodent control. If bait molds, decomposes, or sours, replace with fresh material.

**MOUSE CONTROL:** Tablespoon amounts (1/4 to 1/2 ounce) of bait should be placed at 8-12 foot intervals. A continuous supply of bait should be maintained for at least 15 days.  
 USDA REG NO 6830-25