

# OCTAGON PROCESS INC.

596 RIVER ROAD

EDGEWATER, N. J.

## RODENTICIDE, BAIT, ANTICOAGULANT, WARFARIN 0.025% OCTAGON WARFARIN 0.025%

NET WT 100 LBS. ACTIVE INGREDIENTS	PERCENT BY WEIGHT
WARFARIN 3- alpha-acetonylbenzyl -4-hydroxycoumarin	0.025%
INERT INGREDIENTS	99.975%
Cracked Corn	19.000%
Corn Meal	42.900%
Rolled Oats	28.475%
Petrolatum Liquid	4.800%
Sugar Granulated	4.800%
	<hr/>
	Total 100.000%

### CAUTION

#### KEEP OUT OF REACH OF CHILDREN

**CAUTION:** Keep away from humans, domestic animals and pets. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bis-hydroxycoumarin.

**RAT CONTROL:** Place 4 to 8 ounces of Rodenticide, Bait, Anticoagulant in bait stations. Place so as to protect children, pets and livestock in locations where rats or mice feed. Replenish baits as necessary, and maintain stations for not less than 12 days. Baiting may be continued indefinitely as increased time of exposure of baits may result in better rodent control. If bait molds, decomposes, or sours, replace with fresh material.

**MOUSE CONTROL:** Tablespoon amounts, 1/4 to 1/2 ounce of bait should be placed at 8-12 foot intervals. A continuous supply of bait should be maintained for at least 15 days.

USDA REG NO. 6830-42

MADE IN U.S.A.  
1/1/71