

OCTAGON PROCESS INC., Edgewater, New Jersey 07020
INSECTICIDE, CHLORDANE, 45 PERCENT
WATER EMULSIFIABLE CONCENTRATE
(APPROX. 4 LBS. CHLORDANE GALLON)

Active Ingredients	Percent by Weight
Technical Chlordane	45.0
Emulsifier Hydrocarbons	50.0
Inert Ingredients	
Technical 226 (total conc. 17.4) (total active conc. 17.4)	5.0
17.4 (total inert conc. 82.6)	

CAUTION
KEEP OUT OF REACH OF CHILDREN

DO NOT USE WITHOUT DILUTING

HARMFUL IF SWALLOWED Contact with skin can cause toxic symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. When extensive use is involved, operators should wear respirators approved by the U. S. Bureau of Mines. Avoid contamination of feed and foodstuffs.

TO BE APPLIED BY TRAINED PERSONNEL ONLY

THIS PRODUCT IS TOXIC TO FISH AND WILDLIFE Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DO NOT REUSE EMPTY CONTAINERS Destroy by crushing or perforate and bury in a safe place. "DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME"

MIXING INSTRUCTIONS

This concentrate will not form emulsions that are indefinitely stable. To insure good performance:

1. Prepare the emulsion as soon before actual use as possible.
2. Pour the proper amount of concentrate slowly into the required volume of water with constant stirring and continue agitation for 5 minutes after the addition of the product - see following directions for use for dilutions recommended for control of various pests.
3. Vigorously resuspend immediately before use.

DIRECTIONS FOR USE

RESIDUAL SPRAY IN BUILDINGS (SPOT APPLICATION ONLY)

DO NOT USE IN DAIRY BARN OR POULTRY HOUSES For use in homes around window sills, door sills, stairs, basements, porches, garages, as well as industrial buildings. In living quarters, use as strip treatment only, not for complete coverage of walls. **DO NOT USE IN EDIBLE PRODUCTS AREA OF FOOD PROCESSING PLANTS, RESTAURANTS OR OTHER AREAS WHERE FOOD IS COMMERCIALY PREPARED OR PROCESSED** **DO NOT USE IN SERVING AREAS WHILE FOOD IS EXPOSED**

ROACHES AND ANTS Add 1 part concentrate to 24 parts of water (2 1/2 ounces per gallon of water) and mix thoroughly. As a strip treatment indoors, spray a point of view and apply a 2 inch strip around door frames and sills, window frames and sills, and around the edge of the floor and edge of the ceiling and a strip from floor to ceiling at each corner, and in rooms, at least 20 in. wall space, wall, ceiling, and floor. For roaches, apply as a residual spray to cracks, crevices, under furniture, and the walls of the rooms, at the approximate rate of 1 gallon per 1000 sq. ft. Repeat applications as necessary.

OUTDOOR SPRAY

DO NOT SPRAY VEGETATION USED FOR FOOD, FEED OR FORAGE

MOSQUITOES, FLIES AND SAND FLIES For control of these insects, add 1 part concentrate to 47 parts of water (2 3/4 ounces per gallon of water) and mix thoroughly. Apply only to vegetation at the rate of 5 to 10 gallons per acre, depending on the amount of vegetation to be treated.

MITES (CHIGGERS), TICKS AND FLEAS Add 1 part concentrate to 24 parts of water (2 1/2 ounces) for control of ticks and apply the same concentration at the rate of 15 to 20 gallons per acre, depending on amount of vegetation. For fleas, apply at the rate of 20 to 25 gallons per acre. For chiggers, apply at the rate of 20 to 25 gallons per acre.

EPA REG. NO. 6699-72