

SAMPLE LABEL

Sanitizing of Food Processing Equipment and other Hard Surfaces in Food Contact Locations.

For sanitizing food processing equipment, dairy equipment, food utensils, dishes, silverware, glasses, sink tops, countertops, refrigerated storage and display equipment and other hard surfaces. No potable water rinse is required.

Wash and rinse all articles thoroughly, then apply a solution of 1 oz _____ in 4 gallons of water (150 ppm active). Surfaces should remain wet for at least one minute followed by adequate draining and air drying. Fresh solution should be prepared daily or when use solution becomes visibly dirty. For mechanical application, use solution may not be reused for sanitizing applications.

Apply to sink tops, countertops, refrigerated storage and display equipment and other stationary hard surfaces by cloth or brush. No potable water rinse is required.

Dishes, silverware, glasses, cooking utensils and other similar size food processing equipment can be sanitized by immersion in a 1 oz /4 gallon dilution of _____. No potable water rinse is required.

At 1 oz /4 gallons, _____ fulfills the criteria of appendix F of the Grade "A" Pasteurized Milk Ordinances 1978 Recommendations of the U.S. Public Health Services in waters up to 1000 ppm of hardness calculated as CaCO₃ when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against Escherichia coli and Staphylococcus aureus.

The udders, flanks, and teats of dairy cows can be sanitized by washing with a solution of 1 oz _____ in 4 gallons of warm water. No potable water rinse is required.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towel back into sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.

Precautionary Statements

Hazardous to Humans and Domestic Animals

DANGER

Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

Storage and Disposal

- Do not contaminate water, food, or feed by storage and disposal.
- Do not store on side.
- Avoid creasing or impacting of side walls.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities by burning. If burned, stay out of smoke.
Metal Containers: Triple rinse (or equivalent); then offer for recycling or reconditioning or dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. (If container is 1 gallon or less, use this container disposal statement.)
Do not reuse empty container (bottle, can, bucket). Wrap container and put in trash.

DISINFECTANT-SANITIZER FUNGICIDE DEODORIZER

Disinfectant-Sanitizer-Fungicide Deodorizer for Hospital, Institutional, Industrial, School, Dairy and Other Farm Use

Active Ingredients

Didecyl dimethyl ammonium chloride 7.5%

Inert Ingredients

92.5%

100.0%

KEEP OUT OF REACH OF CHILDREN

DANGER

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Registration No.

EPA Establishment No.

Net Contents

MANUFACTURED BY:

LONZA INC., 22-10 Route 208, Fair Lawn, N.J. 07410

ACCEPTED

08

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg No. 70830-48

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Apply _____ with a cloth, mop or mechanical spray device. When applied with a mechanical spray device, surface must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily or when the use solution becomes visibly dirty.

Disinfection in Hospitals, Nursing Homes and Other Health Care Institutions.

For disinfecting floors, walls, countertops, bathing areas, lavatories, bedframes, tables, chairs, garbage pails and other hard surfaces.

Add 3 1/2 oz. _____ to 4 gallons water. Apply to previously cleaned hard surface with mop or cloth.

At this use-level, _____ is effective against Pseudomonas aeruginosa.

Disinfectant in Institutions, Industry, and Schools

For disinfecting floors, walls, bedframes, countertops, tables, chairs, garbage pails, bathroom fixtures and other hard surfaces.

Add 2 oz. of _____ to 4 gallons of water. Apply to previously cleaned hard surface with mop or cloth.

At 2 oz./4 gallon use-level _____ is effective against Staphylococcus aureus, Salmonella choleraesuis and Trichophyton interdigitale (the athlete's foot fungus).

Disinfection of Barber Tools

Precleaned barber tools (such as combs, brushes, razors, and scissors) can be disinfected by immersing in a 1/2 oz/gallon solution of _____.

Disinfection of Poultry Equipment, Animal Quarters and Kennels.

Poultry brooders, watering founts, feeding equipment and other animal quarters (such as stalls and kennel areas) can be disinfected after thorough cleaning by applying a solution of 2 oz. _____ in 4 gallons of water with a mop, cloth or brush.

Small utensils should be immersed in this solution.

Prior to disinfection, all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and watering appliances. Remove all litter and droppings from floors, walls and other surfaces occupied or traversed by poultry or other animals.

After disinfection, ventilate buildings, coops and other closed spaces. Do not house poultry, or other animals, or employ equipment until treatment has been absorbed, sat or dried.

All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

LD 4

with COMMENTS in EPA Letter Dated

JUN 12 1989

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg No.

6836-48