

Sanitizing of Food Processing Equipment in Food Contact Locations

For sanitizing food processing equipment, dairy equipment, utensils, dishes, silverware, glassware, etc., use 3 oz. of Bardac 20-10 in 4 gallons of water. Equipment should remain wet for at least 10 minutes. No potable water rinse is required.

Wash and rinse all articles thoroughly. Then apply 3 oz. of Bardac 20-10 in 4 gallons of water. Equipment should remain wet for at least 10 minutes. No potable water rinse is required.

Apply to sink tops, refrigerators, and other food processing equipment and the surrounding area. No potable water rinse is required.

Dishes, silverware, glassware, and other articles should be washed with a solution of 3 oz. Bardac 20-10 in 4 gallons of water. No potable water rinse is required.

At 100°F, 3 oz. Bardac 20-10 in 4 gallons of water is the minimum concentration for the Grater-A-Pasteurized Milk and other food processing equipment. The U.S. Public Health Service Water Laboratory has determined that the hardness calculated as CaCO₃ when used at 100°F is 100 grains per gallon. Germicidal and Detergent Sanitizer Method against *Escherichia coli* and *Staphylococcus aureus*.

The udders, flanks, and teats of dairy cows should be washed with a solution of 3 oz. Bardac 20-10 in 4 gallons of water. No potable water rinse is required.

Use a fresh towel for each cow. Avoid contact with the face and solution by dripping. Do not place wet towels in sanitizing solution. When solution becomes very dirty, discard and apply a fresh solution.

Precautionary Statements

Hazards to Humans and domestic animals

DANGER

Keep out of reach of children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

STORAGE AND DISPOSAL

- DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL
- OPEN DUMPING IS PROHIBITED
- DO NOT REUSE EMPTY CONTAINER

PESTICIDE DISPOSAL

PESTICIDE OR RINSATE THAT CANNOT BE USED OR CHEMICALLY REPROCESSED SHOULD BE DISPOSED OF IN A LANDFILL APPROVED FOR PESTICIDES OR BURIED IN A SAFE PLACE AWAY FROM WATER SUPPLIES.

CONTAINER DISPOSAL

TRIPLE RINSE (OR EQUIVALENT) AND DISPOSE IN AN INCINERATOR OR LANDFILL APPROVED FOR PESTICIDE CONTAINERS, OR BURY IN A SAFE PLACE.

GENERAL

CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATIVE PROCEDURES SUCH AS LIMITED OPEN BURNING.

BARDAC 20-10

Disinfectant/Sanitizer

Biocide/Deodorizer

with Organic Tolerance

for Hospital, Institutional, Industrial, School, Dairy, and Other Food Use

Active Ingredients

Octyl decyl dimethyl ammonium chloride	5.0%
Didecyl dimethyl ammonium chloride	2.5%
Dioctyl dimethyl ammonium chloride	2.5%

Inert Ingredients

90.0%
100.0%

KEEP OUT OF REACH OF CHILDREN

DANGER

Statement of Practical Treatment

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

SEE LEFT PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Registration No. 6836-65

EPA Establishment No. 6836-IL-1

Net Contents

Manufactured By:

LONZA INC. • 22-10 Route 208 • Fair Lawn, N.J. 07410

Directions for Use

GENERAL CLASSIFICATION

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Bardac 20-10 is a disinfectant/sanitizer, biocide/deodorizer with organic tolerance for Hospital, Institutional, Industrial, School, Dairy, and Other Food Use.

Bardac 20-10 is a disinfectant/sanitizer, biocide/deodorizer with organic tolerance for Hospital, Institutional, Industrial, School, Dairy, and Other Food Use.

Bardac 20-10 is a disinfectant/sanitizer, biocide/deodorizer with organic tolerance for Hospital, Institutional, Industrial, School, Dairy, and Other Food Use.

Apply 3 oz. Bardac 20-10 in 4 gallons of water. Apply to surfaces, equipment, and other food contact areas.

2 oz. Bardac 20-10 in 4 gallons of water. Apply to surfaces, equipment, and other food contact areas.

Bardac 20-10 is a disinfectant/sanitizer, biocide/deodorizer with organic tolerance for Hospital, Institutional, Industrial, School, Dairy, and Other Food Use.

Bardac 20-10 is a disinfectant/sanitizer, biocide/deodorizer with organic tolerance for Hospital, Institutional, Industrial, School, Dairy, and Other Food Use.

Apply 3 oz. Bardac 20-10 in 4 gallons of water. Apply to surfaces, equipment, and other food contact areas.

Bardac 20-10 is a disinfectant/sanitizer, biocide/deodorizer with organic tolerance for Hospital, Institutional, Industrial, School, Dairy, and Other Food Use.

Disinfection of Poultry Equipment, Animal Quarters and Kennels.

Poultry brooders, watering fountains, feeding equipment and other animal quarters (such as stalls and kennel areas) can be disinfected after thorough cleaning by applying a solution of 3 oz. Bardac 20-10 to 4 gallons of water with a mop, cloth or brush. Small utensils should be immersed in this solution.

Prior to disinfection, all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and watering appliances. Remove all litter and droppings from floors, walls and other surfaces occupied or traversed by poultry or other animals.

After disinfection, ventilate buildings, coops and other closed spaces. Do not house poultry, or other animals, or employ equipment until treatment has been absorbed, set or dried.

All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

11 7/80