

PRODUCT INFORMATION

ACCEPTED
 JUN 30 1987
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg No. **6836-66**

BARDAC 205M
BARDAC 208M

Bardac 205M/208M is a radically new development based upon Lonza Inc. patented "Twin Chain" quaternary ammonium compound technology. Bardac 205M/208M, when evaluated by accepted laboratory procedures, provides superior germicidal and fungicidal activity far beyond that achieved with currently available quaternary ammonium compounds. This provides the formulator with unequalled latitude in the design of biocidal systems.

Chemical Composition

<u>Active Ingredients</u>	<u>Bardac 205M</u>	<u>Bardac 208M</u>
Alkyl (C ₁₄ , 50%; C ₁₂ , 40%; C ₁₆ , 10%) dimethyl benzyl ammonium chloride	20.0%	32.0%
Octyl decyl dimethyl ammonium chloride	15.0%	24.0%
Dioctyl dimethyl ammonium chloride	7.5%	12.0%
Didecyl dimethyl ammonium chloride	7.5%	12.0%
<u>Inert Ingredients</u>	50.0%	20.0%

Physical Properties

Average molecular weight	342	
pH (10% solution)	6.5—9.0	
Physical state	Liquid	
Color	Clear to light amber	
Flash point (Seta Flash)	116°F	118°F
Specific gravity @ 25°C	0.946	0.912
	(7.89 lbs./gal.)	(7.61 lbs./gal.)
<u>EPA Registration No.</u>	6836-66	6836-67
<u>CAS No.</u>	68424-95-5 & 130-08-2	

Summary of the superior performance characteristics of Bardac 205M/208M:

- Better disinfectant performance.
- Broad spectrum biocidal activity against both gram positive and gram negative organisms.
- Increased hard water tolerance for sanitizing activity
- Superior fungicidal performance.
- Substantial organic soil tolerance (in accordance with the latest EPA requirements).

GERMICIDAL ACTIVITY

Important Note: All microbiological evaluations were performed in the presence of an organic soil load represented by 5% blood serum; proposed EPA guidelines — Subpart G. The serum was added to the inoculum prior to the carrier drying step (or other appropriate procedure).

Disinfectant Activity Determined by AOAC Use-Dilution Tests

The minimum concentration of Bardac 205M/208M required for effective disinfection is described in the current procedures of the AOAC; commonly known as the Use Dilution Test.

<u>Test Organism</u>	<u>ATCC #</u>	<u>Minimum Effective Concentration</u>
Staphylococcus aureus	6538	250 ppm active quaternary
Salmonella choleraesuis	10708	250 ppm active quaternary
Pseudomonas aeruginosa	10524	450 ppm active quaternary

The broad spectrum germicidal activity of Bardac 205M/208M was confirmed by the AOAC Use-Dilution evaluations against the following organisms:

<u>Test Organism</u>	<u>ATCC #</u>	<u>Minimum Effective Concentration</u>
Escherichia coli	11229	250 ppm active quaternary
Serratia marcescens	8101	250 ppm active quaternary
Brevibacterium ammoniagenes	6871	250 ppm active quaternary
Salmonella typhi	6539	250 ppm active quaternary
Pseudomonas cepacia	17765 25608 25416	450 ppm active quaternary

**Disinfectant Activity Determined by AOAC Use Dilution tests
In the Presence of Hard Water and Organic Soil**

<u>Test Organism</u>	<u>ATCC #</u>	<u>Hard Water Concentration</u>	<u>Minimum Effective Concentration</u>
Pseudomonas aeruginosa	10524	0 ppm/CaCO ₃	450 ppm active quaternary
		200 ppm/CaCO ₃	850 ppm active quaternary
		400 ppm/CaCO ₃	850 ppm active quaternary
		500 ppm/CaCO ₃	1000 ppm active quaternary
Salmonella choleraesuis	10708	0 ppm/CaCO ₃	250 ppm active quaternary
		300 ppm/CaCO ₃	600 ppm active quaternary
		400 ppm/CaCO ₃	600 ppm active quaternary
		500 ppm/CaCO ₃	700 ppm active quaternary
Pseudomonas cepacia	17765 25416 25608	400 ppm/CaCO ₃	850 ppm active quaternary

ACCEPTED
 JUN 30 1987
 U.S. Environmental Protection Agency
 Research Triangle Park, NC 27711
 as submitted for the pesticide registration under EPA Reg. No. 6536-66

VIRUCIDAL ACTIVITY

The virucidal activity of Bardac 205M/208M was confirmed by current EPA accepted protocol against Influenza A₂ (Hong Kong), Herpes simplex Type 1 and Type 2, and Vaccinia at 450 ppm active quaternary.

Sanitizing Performance/Hard Water Tolerance

Bardac 205M/208M has been cleared by the FDA as an "Indirect Food Additive" under 21CFR178.1010 "Sanitizing Solutions" at a concentration of 150-400 ppm active, and requires no potable water rinse. This clearance covers the usage of Bardac 205M/208M on food processing equipment and utensils and food contact surfaces in public eating places. In addition, the use of sanitizing solutions based on Bardac 205M/208M fulfills the criteria of the Grade "A" Pasteurized Milk Ordinance 1978 Recommendations of the United States Public Health Service.

The hard water tolerance of Bardac 205M/208M is measured by the AOAC Germicidal and Detergent Sanitizer Method, commonly called the hard water tolerance test. Exposure of 100 million organisms of Escherichia coli (#11229) to 150 ppm of Bardac 205M/208M for 30 seconds at 25°C in 800 ppm of water hardness results in the required reduction of 99.999% of the bacteria.

Calcium and magnesium salts are typical hard water components; however, other electrolytes may be present during actual field applications.

<u>Test Organism</u>	<u>Concentration of Bardac 205M/208M Required for 99.999% Reduction</u>	<u>Hard Water Ceiling</u>
Escherichia coli, #11229	150 ppm	800 ppm hard water
	200 ppm	1100 ppm hard water
Staphylococcus aureus, #6538	150 ppm	800 ppm hard water
	200 ppm	1100 ppm hard water

Non-Food Contact Application

The non-food contact surface sanitizing activity of Bardac 205M/208M was determined by the following test methodology: EPA DIS/TSS-10, February 6, 1979 and the Guidelines for Registering Pesticides in the United States, Subpart G - Product Performance, June 22, 1979 Draft, Recommended Method #8 (Sanitizers-Non-Food Contact Surfaces), p 101-102.

<u>Test Organism</u>	<u>Concentration of Bardac 205M/208M Required for 99.999% Reduction</u>	<u>Hard Water Ceiling</u>	<u>Contact Time</u>
Staphylococcus aureus (ATCC #6538)	150 ppm (active)	800 ppm	2 min
	200 ppm (active)	1100 ppm	1 min.
Klebsiella pneumoniae (ATCC #4352)	150 ppm (active)	800 ppm	2 min.
	200 ppm (active)	1100 ppm	1 min.

Fungicidal Performance as Determined by the AOAC Fungicidal Test

Possessing superior fungicidal activity, Bardac 205M/208M effectively passes the AOAC Fungicidal test at one-eighth the concentration required for conventional alkyl benzyl quaternaries.

<u>Test Organism</u>	<u>Ten Minute Killing Dilution (100% active)</u>
Trichophyton mentagrophytes	1:8000 125 ppm

ACCEPTED

JUN 30 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 6836-66

Summary of Applications and Recommended Use-Levels

<u>Application</u>	<u>Recommended Use-Levels on 100% Active Basis</u>
Hospital Disinfection	450 ppm active quaternary
General Disinfection	250 ppm active quaternary
Sanitizing	150 ppm active quaternary

Product Registration

The Lonza Technical Service Department will assist you with Bardac 205M/208M based formulations and EPA registration.

Prototype formulations based on Bardac 205M/208M, their EPA data base references and sample labels are available upon request. For formulations of your own development, you are responsible to provide data currently required by the EPA to support that registration.

Disinfectant and sanitizer products containing Quaternary Ammonium Compounds must be registered with the U.S. Environmental Protection Agency. Applications for registration must be accompanied by two copies of your proposed label and should be sent to Product Manager No. 31, Office of Pesticide Programs, Registration Division (TS-767C), U.S. Environmental Protection Agency, Washington, D.C. 20460. Some state agencies also require registration of your product independent of your EPA registration.

Precautionary Statements


Danger: Keep Out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Do not contaminate water, food or feed by storage or disposal.

Statement Of Practical Treatment: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

Note To Physician: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Container Disposal: Dispose of in an incinerator or landfill approved for pesticide containers. Bury in a safe place or return to drum reconditioners.



The seller makes no warranty, expressed or implied, concerning the accuracy or any results to be obtained from the use of any information and no warranty, expressed or implied, concerning the use of the products other than indicated above. The buyer assumes all risks of use and handling. No statement is intended or shall be construed as a recommendation to infringe any patent.