

Sanitizing of Food Processing Equipment and Other Facilities
Contact Locations

No Potable water rinse is required

205-75

BARDAC

No Potable water rinse is required

BARDAC-205-75 No Potable water rinse is required

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC-205-75

BARDAC 205M-7.5

Disinfectant-Sanitizer
Fungicide-Deodorizer
with Organic Soil Tolerance
for Hospital, Institutional,
Industrial, School, Dairy and Other Facility Use

Active Ingredients

Alkyl (C-12 to C-14) 40% C-10 to C-12 dimethyl	
benzyl dimethylammonium chloride	1.25%
Decyl dimethylammonium chloride	1.25%
Dodecyl dimethylammonium chloride	1.25%

Inert Ingredients

92.500%

100.000%

KEEP OUT OF REACH OF CHILDREN

DANGER

Statement of Practical Treatment

In case of contact immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes call a physician. Remove and wash contaminated clothing before reuse. If swallowed drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

SEE LEFT PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

EPA Registration No

6836-70

EPA Establishment No.

6836-IL-1

Net Contents

MANUFACTURED BY

ION77 INC 22-10 Route 208 Fair Lawn, N.J. 07410

Directions for Use

GENERAL CLASSIFICATION

As a disinfectant Federal Law to use this product in a manner consistent with its labeling.

Apply BARDAC-205-7.5 with a cloth mop or mechanical spray device. When applied with a mechanical spray device, surface must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily or when the use solution becomes cloudy.

Disinfection in Hospitals, Nursing Homes and Other Health Care Institutions. For disinfecting floors, walls, counter tops, bathing areas, lavatories, bed frames, chairs, chairs, garbage pails and other hard surfaces.

Add 1 oz. of BARDAC-205-7.5 to 4.5 gallons water. Apply to above cleaned hard surfaces. At this use level BARDAC-205-7.5 is effective against *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Salmonella choleraesuis* in the presence of 5% blood serum when evaluated by the AOAC Use-Dilution Test.

Disinfection in Institutions, Industry and Schools. Add 1 oz. of BARDAC-205-7.5 to 4.5 gallons of water. 20 oz. of 1:25 gallons of water use level for the purpose of effectiveness of BARDAC-205-7.5 in the presence of 5% blood serum against *Trichophyton mentagrophytes* has been shown utilizing the AOAC Fungus Test.

Disinfection of Barber Tools. Pre-cleaned barber tools such as brushes, razors and scissors can be disinfected by immersing in a 5 oz. gallon solution of BARDAC-205-7.5.

Disinfection of Poultry Equipment, Animal Quarters and Kennels. Poultry brooders, watering fountains, feeders, and other animal quarters (such as stalls and pens) can be disinfected after thorough cleaning by applying a solution of 2 oz. BARDAC-205-7.5 to 4.5 gallons of water. Small animals can be immersed in this solution.

Disinfect the floors of poultry, other animals and kennels with this solution. This solution may be used on floors and other hard surfaces. Apply with a cloth mop or mechanical spray device. Surfaces must remain wet for 10 minutes.

Apply to the floors of poultry, other animals and kennels with this solution. This solution may be used on floors and other hard surfaces. Apply with a cloth mop or mechanical spray device. Surfaces must remain wet for 10 minutes.

Apply to the floors of poultry, other animals and kennels with this solution. This solution may be used on floors and other hard surfaces. Apply with a cloth mop or mechanical spray device. Surfaces must remain wet for 10 minutes.

8