

Cross contamination is of major housekeeping concern not only in hospitals, but in schools, institutions, and industry. LONZA FORMULATION S-37 has been formulated to aid in the reduction of cross contamination in these areas. LONZA FORMULATION S-37 is a one-step disinfectant cleaner that is effective against a broad spectrum of bacteria, is virucidal, fungicidal, and inhibits the growth of mold and mildew and their odors when used as directed. LONZA FORMULATION S-37 is an effective one-step sanitizer-cleaner for use on non-food contact surfaces.

When used as directed, LONZA FORMULATION S-37 will deodorize surfaces in toilet areas, behind and under sinks and counters, garbage cans and garbage storage areas, and other places where bacterial growth can cause mal odors.

Precautionary Statements

Hazard to Humans and Domestic Animals

DANGER

Keep Out of Reach of Children. Corrosive. Causes eye and skin damage. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

Storage and Disposal

- Do not contaminate water, food, or feed by storage or disposal.
- Do not store on side
- Avoid creasing or impacting of side walls.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray or mixture of rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities by burning. If burned, stay out of smoke.
Metal Container: Triple rinse (or equivalent), then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by other procedures approved by state and local authorities (If container is 1 gallon or less, use this container disposal statement.)
Do not reuse empty container (bottle, can, bucket). Wrap container and put in trash.

BEST AVAILABLE COPY

LONZA FORMULATION S-37

Disinfectant-Cleaner-Sanitizer
Fungicide-Mildewstat-Virucide-Deodorizer
with Organic Soil Tolerance
for Hospitals, Institutional and Industrial Use

Active Ingredients

Octyl decyl dimethyl ammonium chloride	2.10%
Dioctyl dimethyl ammonium chloride	0.84%
Didodecyl dimethyl ammonium chloride	1.26%
Alkyl (C ₁₂ , 50%; C ₁₃ , 40%; C ₁₄ , 10%) dimethyl benzyl ammonium chloride	2.80%

Inert Ingredients

93.00%
100.00%

KEEP OUT OF REACH OF CHILDREN

DANGER

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Registration No.

6836-72

EPA Establishment No.

6836-IL-1

Net Contents

MANUFACTURED BY:

LONZA INC., 17-17 Route 208,

NEW YORK, N.Y. 10013

Accepted for the Federal Government
Fungicide and Mildewstat
as amended for the purpose
registered under
EPA Reg. No. 6836-72

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

LONZA FORMULATION S-37 is a proven one step cleaner-disinfectant-sanitizer-fungicide-mildewstat-virucide in the presence of moderate amounts of organic soil.

Apply LONZA FORMULATION S-37 to walls, floors and other hard (inanimate) non porous surfaces such as tables, chairs, countertops, tile, porcelain, sinks and bedframes with a cloth, mop or mechanical spray device so as to thoroughly wet surfaces. For heavily soiled areas, a preliminary cleaning is required. Prepare a fresh solution daily or when use and then becomes visibly dirty.

Disinfection - to disinfect hard, non porous surfaces a solution of 1 oz. per gallon of water. Treated surfaces must remain wet for 10 minutes. The activity of LONZA FORMULATION S-37 has been evaluated in the presence of 5% serum by the APAC use dilution test and found to be effective against a broad spectrum of gram negative and gram positive organisms as represented by:

<i>Pseudomonas aeruginosa</i>	<i>Salmonella schellenghii</i>
<i>Staphylococcus aureus</i>	<i>Streptococcus faecalis</i>
<i>Salmonella choleraesuis</i>	<i>Shigella dysenteriae</i>
<i>Escherichia coli</i>	<i>Brachytherium amnioniae</i>
<i>Streptococcus pyogenes</i>	<i>Serratia marcescens</i>
<i>Proteus mirabilis</i>	<i>Salmonella typhi</i>
<i>Klebsiella pneumoniae</i>	<i>Pseudomonas cepacia</i>
<i>Enterobacter aerogenes</i>	

Fungicidal Performance

LONZA FORMULATION S-37 is an effective fungicide against *Trichophyton mentagrophytes* (the athlete's foot fungus) which is used on surfaces in areas such as locker rooms, dressing rooms, shower and bath areas, exercise facilities, etc. 1 oz. or 1 gallon.

Mold and Mildew Control

At 1 oz. or 1 gallon, LONZA FORMULATION S-37 will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard, non porous surfaces (as indicated in general instructions above). Allow 24 hours to dry and repeat when mildew growth returns.

Virucidal Performance

At 1 oz. per gallon or less LONZA FORMULATION S-37 was evaluated in the presence of 5% serum and found to be effective against the following viruses: Influenza A, B, C, Herpes Simplex Types 1 and 2 and Varicella on both hard and non porous environmental surfaces.

Sanitizing Non Food Contact Surfaces (such as walls, tables, etc.) At 1 oz. per 25 gallons use LONZA FORMULATION S-37 as an effective sanitizer against *Staphylococcus aureus* and *Klebsiella pneumoniae* in the presence of 5% serum on porous and non porous environmental surfaces. Treated surfaces must remain wet for 2 minutes.