

Precautionary Statements
Hazardous to Humans and Domestic
Animals

DANGER

Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed.

Storage and Disposal

- Do not contaminate water, food, or feed by storage disposal.
- Do not store on side.
- Avoid creasing or impacting of side walls.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray or mixture of rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities by burning. If burned, stay out of smoke.


Metal Container: Triple rinse (or equivalent), then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. (If container is 1 gallon or less, use this container disposal statement.)

Do not reuse empty container (bottle, can, bucket). Wrap container and put in trash.

ACCEPTED

SEP 12 1985

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 6836-83



LONZA CS 201

No-Rinse Acid Cleaner Sanitizer For Dairy, Food Processing and Farm Use

Removes Milkstone

Active Ingredients

Octyl decyl dimethyl ammonium chloride	2.2950%
Didecyl dimethyl ammonium chloride	1.1475%
Dioctyl dimethyl ammonium chloride	1.1475%
Alkyl (C ₁₄ , 50%; C ₁₂ , 40%; C ₁₆ , 10%) dimethyl benzyl ammonium chloride	3.0600%

Inert Ingredients

92.3500%
100.0000%

KEEP OUT OF REACH OF CHILDREN

DANGER

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Registration No. 6836-83

EPA Establishment No. 6836-IL-1

Net Contents

MANUFACTURED BY:

LONZA INC., 22-10 Route 208, Fair Lawn, N.J. 07410

For Sale For Use And Storage By
Maintenance Personnel Only

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

LONZA CS 201 is an acid cleaner sanitizer for use on food processing and dairy equipment in water up to 600 ppm hardness (as CaCO₃). Do not rinse the water. For all equipment, a preliminary cleaning is required before sanitizing. Prepare a fresh solution daily or when use solution becomes visibly dirty. For mechanical application, use solution may not be reused for sanitizing applications.

Cleaning Bulk Milk Tanks, Dairy and Food Processing Equipment

Sanitizing—To sanitize hard non-porous food contact surfaces, such as milk tanks and milking equipment, clean with a good detergent and rinse thoroughly, dilute 1 oz. of LONZA CS 201 in 4 gallons of water (150 ppm active quaternary) and apply with a cloth, brush, mechanical spray device (spray surface until thoroughly wetted) or by immersion. Treated surfaces should remain wet for at least 60 seconds followed by adequate draining and air drying. Do not rinse with water.

Grade A Pasteurized Milk Ordinance

At 1 oz./4 gallons LONZA CS 201 fulfills the criteria of Appendix F of the Grade "A" Pasteurized Milk Ordinances 1978 Recommendations of the U.S. Public Health Services in waters up to 600 ppm of hardness calculated as CaCO₃ when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against escherichia coli and Staphylococcus aureus.

The udders, flanks and teats of dairy cows can be sanitized by washing with a solution of 1 oz. LONZA CS 201 in 4 gallons of warm water. No potable water rinse is required. Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towel back into sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.