Cross-contamination is of major housekeeping concern not only in hospitals, but in schools, institutions and industry. LONZA FORMULATION S-18F has been formulated to aid in the reduction of cross-contamination in these

LONZA FORMULATION S-18F is a one-step disinfectant cleaner that is effective against a broad spectrum of bacteria, viruses (including EIV), fungi and inhibits the growth of mold and milder and their odors when used as directed.

LONZA FORMULATION S-18F is an effective one-step sanitizer-cleaner for use on food contact, as well as non-food contact surfaces. When used as directed, LONZA FORMULATION S-18F will deodorize surfaces. in toilet areas, behind and under sinks and counters, garbage cans and garbage storage areas and other places where bacterial growth can cause mal odors.

PRECAUTICMARY STATEMENTS HAZARD TO BUHANS AND DOMESTIC ANTIMALS

DANGER

Keep out of reach of children. Corrosive. Causes eye and skin damage. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed or absorbed through skin. Avoid contamination of food.

STURAGE AND DISPOSAL

- Do not contaminate water, food, or feed by storage disposal.
- Do not store on side
- Avoid creasing or impacting of side walls.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide. spray or mixture of rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Festicide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance,

CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Hetal Container: Triple rinse (or equivalent), then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

(If container is 1 gallon or less, use this container disposal statement.) Do not rouse empty container (bottle, can, bucket). Wrap container and put in

LONZA FORMULATION S-18F

DISINFECTANT, CLEANER, SANITIZER, FUNGICIDE MILDEWSTAT VIRUCIDE, DEODORIZER FOR BOSPITALS, INSTITUTIONAL AND INDUSTRIAL USE

Active Ingredients

Octyl decyl dimethyl ammonium chloride Dioctyl dimethyl ammonium chlorida Didecyl dimethyl ammonium chlorida	3.255% 1.302% 1.953%		
		Alkyl (C ₁₄ , 50%; C ₁₂ , 40%; C ₁₆ , 10%)	
		Alkyl (C ₁₄ , 50%; C ₁₂ , 40%; C ₁₆ , 10%) dimethyl benzyl ammonium chloride	4.340%

Inert Ingredients Total Ingredients

KEEP OUT OF REACH OF CRITISHEN

DANGER

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a phys cian immediately.

NOTE TO PHYSICIAM: Probable mucosal damage may contraindicate the use of gastric lavage.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Registration No.

6836-136

EPA Establ:shment No.

6836-IL-1

Net Contents

ACCEPTED OCT 20 1992

U For the Foder I Insecticide, I am de and Endentride Act. cam nied, for the posticide EPA Reg. No. 6836-136

danufactured by: L/N/A IN', 17-17 Route 208, Fair Lawn, NJ 07410 DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

LONZA FORMULATION S-18F is a proven one-step disinfectant-cleaner-sanitizer-fungicide-mildewstatvarueide which is effective in water up to 400 ppm hardness in the presence of 5% serum contamination. Apply LONZA FORMULATION S-18F to walls, floors and other hard (inenimate) non-porous surfaces, such as tables, chairs, countertops, sinks, porcelain, tile and bedframes with a cloth, mop or mechanical spray device so as to thoroughly wet so faces. For heavily soiled areas, a preliminary cleaning is required. Prepare a fresh solution daily or when use solution becomes visibly dirty.

DISTRICTION: To disinfect hard, non-porous surfaces, add 1 oz. per gallon of water. Treated surfaces must remain wet for 10 minutes. The activity of LONZA FORMULATION S-18F has been evaluated in the presence of 5% serum and 400 ppm hard water by the ACAC Use-Dilution test and found to effective against a broad spectrum of gram negative and gram positive organisms as represented

Staphylococcus aureus Salmonella choleraesuis Shigella dysenteriae Escherichia coli Streptococcus pyogenes Salmonella typhi Klubsiella pneumoniae Enterobacter aerogenes Pseudomonas cepacia

111-1

Pseudomonas seruginosa Salmonella schottmuelleri Streptococcus faecalis Brevibacterium ammoniagenes Serratia marcescens

VIRICIDAL PERFURMANCE: At 1 oz. per gallon uselevel LONZA FORMULATION S-18F was evaluated in the presence of 5% serum and 400 ppm hard water against the following viruses found on insmimate, hard, nonporous, environmental surfaces;

HIV-1 (AIDS virus) Influenza A/Brazil Vaccinia

Herpes Simplex Type 1 and Type 2

FUNGICIDAL CONTROL: LONZA FORMULATION S-18F is an effective fungicide against Trichophyton mentagrophytes (the athlete's foot fungus) when used on .urfaces in areas such as locker rooms, dressing rooms, shower and bath areas, exercise facilities at 1 oz. per gallon.

MOLD AND MILDEN CONTROL: At 1 oz. per gallon, LONZA FORMULATION S-18F will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard, non-porous surfaces (as indicated in general instructions above). Allow to dry on surface and repeat when mildew growth returns

SANITIZING FOOD CONTACT SURFACES: At 1 oz. per 5 1/2 gallons water use LONZA FORMULATION S-18F for sanitizing and cleaning of equipment and utensils in food processing, dairy industry, bars, restaurants, institutional kitchens, meet and poultry processing plant and farms. Do not rinse with water. For heavily soiled equipment or utensils, pre-wash with LONZA FORMULATION S-18F. Rinse with clean water, then apply recommended dilution of LONZA FORMULATION S-18F with a cloth, mechanical spray device (spray surface until thoroughly wetted) or immersion. Treated surfaces must remain wet for one minute. Prepare a fresh solution for every use. DO NOT reuse mechanically applied use solution for senitizing. LONZA FORMULATION S-18F can be used in accordance with the U.S. Public Health Service food service sanitation recommendation.

KILLS BIY OF FRE-CLEARED ENVIRONMENTAL. STREPACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD FLUIDS in health care settings or other setting in which there i an expected likelihood of soiling of inamimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV-1) (associated

1 4 1 1

"SPECIAL DESTRUCTIONS FOR CLEANING AND DECOSTANDRATION AGAINST HIV-1 ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS":

CONTACT TIME: Leave surface wet for 10 minutes with 1 oz. per gallon use solution.

CLEANING PROCEDURE: Blood and other body fluid containing HIV must be thoroughly cleaned fr surfaces and objects before application of LOW. FORMULATION 5-18F.

PERSONAL PROTECTION: Clean up should always be done wearing protective latex gloves, gowns, masks and eye protection.

DISPOSAL OF DEFECTIONS MATERIAL: Blood, body fluids, cleaning materials and clothing should be autoclayed and disposed of according to local regulations for infectious waste disposal.

U.S. PUBLIC REALTH SERVICE FOOD SERVICE SANITIZATION RECOMMENDATIONS

CIPARING AND SANITIZING: Equipment and utensils shall be thoroughly pre-flushed or pre-scraped and when necessary, pre-scaked to remove gross food particles and soil.

- 1. Thoroughly wash equipment and utensils in a hot detergent solution.
- 2. Rinse utensils and equipment thoroughly with clean water.
- 3. Sanitize equipment and utensils by immersi in a 1 oz. per 5 1/2 gallon solution of LONZ FORMULATION S-18F (150 ppm active) for at least 60 seconds at a temperature of 75°F. The Children of the control of the c
- 4. For equipment and utensils too large to sanitize by immersion, apply a 1 oz. per 5 1/2 gallon solution of LONZA FORMULATION S-18F (150 ppm active) by rinsing, spraying or swatting until thoroughly wetted.
- 5 Allow sanitized surface to drain and air dry Do not rinse.

Wash thoroughly with soap and *ater afte. handling.

02/03/92