

6836-204

12/28/2000

1/5

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**DANGER.** Corrosive. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin or on clothing. Wear protective eyewear (goggles, face shield or safety glasses) and rubber gloves when handling. May be fatal if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse.

**STATEMENT OF PRACTICAL TREATMENT (FIRST AID)**

**IF IN EYES:** Hold eyelids open and flush with a steady gentle stream of water for 15 minutes. Get medical attention.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention.

**IF SWALLOWED:** Call a doctor or get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person. Drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

*(If container size is 5 gallons or greater, use the following Environmental Hazards statements.)*

**ENVIRONMENTAL HAZARDS**

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

*(If container size is one gallon or less, use the following Storage and Disposal statements.)*

**STORAGE AND DISPOSAL**

Store in original container in areas inaccessible to children. Do not reuse empty container. Wrap and discard in trash.

*(If container size is greater than one gallon, use the following Storage and Disposal statements.)*

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE**

pen dumping is prohibited. Store in original container in areas inaccessible to children.

**PESTICIDE DISPOSAL**

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray or mixture of rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL**

Do not reuse empty container.

*(For Metal containers:)* Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose in a sanitary landfill, or by other procedures approved by state and local authorities.

*(For Plastic containers:)* Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**LONZA FORMULATION DC-110N**

Disinfectant – Cleaner – Sanitizer – Fungicide – Mildewstat -  
Virucide\* - Deodorizer  
for Hospitals, Institutional and Industrial Use

Effective in hard water up to 400 ppm hardness (Calculated as CaCO<sub>3</sub>) in the presence of 5% serum contamination

Fresh clean scent / Lemon scent / Pine Scent / Mint scent /  
Floral scent / Citrus scent

|                                    |         |
|------------------------------------|---------|
| Active Ingredient:                 |         |
| Didecyl dimethyl ammonium chloride | 3.85%   |
| Inert Ingredients:                 | 96.15%  |
| Total:                             | 100.00% |

**KEEP OUT OF REACH OF CHILDREN**

**DANGER PELIGRO**

SEE SIDE / LEFT / RIGHT / BACK PANEL FOR  
PRECAUTIONARY STATEMENTS AND FIRST AID

EPA Reg. No. 6836-204  
EPA Est. No. (as noted on the container)  
Net Contents (as noted on the container)

Manufactured by:  
LONZA INC., 17-17 Route 208, Fair Lawn, NJ 07410

**ACCEPTED**

DEC 28 2000

Under the Federal Insecticide, Fungicide, and  
Rodenticide Act as amended, and the  
pesticide, registered under  
EPA Reg. No. 6836-204

Cross-contamination is of major housekeeping concern not only in hospitals, but also in schools, institutions and industry. LONZA FORMULATION DC-110N has been formulated to aid in the reduction of cross-contamination from treated surfaces in these areas.

LONZA FORMULATION DC-110N a one-step disinfectant cleaner that is effective against a broad spectrum of bacteria, is virucidal\* (including HIV-1), fungicidal, and inhibits the growth of mold and mildew and their odors when used as directed.

LONZA FORMULATION DC-110N is an effective one-step sanitizer-cleaner for use on non-food, contact surfaces.

When used as directed, LONZA FORMULATION DC-110N will deodorize surfaces in toilet areas, behind and under sinks and counters, garbage cans and garbage storage areas, and other places where bacterial growth can cause mal odors.

LONZA FORMULATION DC-110N is an effective fungicide against Trichophyton mentagrophytes (the athlete's foot fungus) when used on surfaces in areas such as locker rooms, dressing rooms, shower and bath areas, exercise facilities.


LONZA FORMULATION DC-110N is recommended for use in hospitals, medical and dental offices, nursing homes, public rest rooms, institutions, schools, colleges, athletic facilities, food processing plants, food storage areas, kitchens, restaurants and bars, airports, transportation terminals, hotels and motels. For use in households. This product contains no phosphorous.

LONZA FORMULATION DC-110N is a no-rinse neutral pH disinfectant cleaner that disinfects, cleans and deodorizes in one labor-saving step.

LONZA FORMULATION DC-110N can be used with a mop and bucket, trigger sprayers, sponge or by soaking.

LONZA FORMULATION DC-110N provides effective cleaning length that will not dull most metal-interlock floor finishes, and does not require a rinse prior to re-coat.

LONZA FORMULATION DC-110N improves labor results by effectively controlling odors.



LONZA FORMULATION DC-110N Cleans, shines, deodorizes and disinfects hard, nonporous household surfaces. Inhibits the growth mold and mildew, leaving bathrooms & kitchens clean and smelling fresh.

LONZA FORMULATION DC-110N is recommended for non-scratch cleaning of showers and tubs, shower doors and curtains, fixtures and toilet bowls.

LONZA FORMULATION DC-110N is for use in kitchens, bathrooms and other household areas.

LONZA FORMULATION DC-110N kills household germs in 10 minutes; effective against harmful Pseudomonas aeruginosa (pseudomonas), Staphylococcus aureus (staph) and Streptococcus pyogenes (strep), Influenza A/PR and Trichophyton mentagrophytes (athlete's foot fungus).

LONZA FORMULATION DC-110N is a neutral pH multi-surface cleaner, deodorizer and disinfectant. Use on windows, mirrors and other glass surfaces.

Use LONZA FORMULATION DC-110N on the multi-touch surfaces responsible for cross-contamination.

Use LONZA FORMULATION DC-110N on finished floors, high speed burnished floors, conductive flooring, washable walls, tables, chairs, bathroom bowls, sinks, basins, shower stalls, tubs, tiles, toilets, countertops, stove tops, appliances, refrigerators, cabinets, garbage cans, tables, outdoor furniture, chairs, desks, telephones, door knobs and handles, and glass surfaces including windows and mirrors.

LONZA FORMULATION DC-110N is recommended for Poultry Premise Sanitation (Hatcheries):

- Egg Receiving Area    Tray Dumping Area    Chick Processing Area
- Egg Holding Area    Chick Holding Room    Chick Loading Area
- Setter Room    Hatchery Room    Poultry Buildings

LONZA FORMULATION DC-110N is recommended for Swine Premise Sanitation:

- Farrowing Barns & Areas    Dressing Plants    Blocks
- Waterers and Feeders    Loading Equipment    Creep Area
- Hauling Equipment    Nursery    Chutes

3/5

**LONZA FORMULATION DC-110N** is a Hospital Disinfectant at 2 ounces per gallon (16 milliliters per liter, 1:64 dilution), modified in the presence of 250 ppm hard water, and at 3 ounces per gallon (24 milliliters per liter, 1:42 dilution), modified in the presence of 400 ppm hard water.

**LONZA FORMULATION DC-110N** is Bactericidal according to the AOAC Use-Dilution Test method, Virucidal\* according to the virucidal qualification on hard, inanimate surfaces, Fungicidal according to the AOAC fungicidal test, modified in the presence of 5% organic serum against:

|                                       |                            |
|---------------------------------------|----------------------------|
| Pseudomonas aeruginosa                | Proteus vulgaris           |
| Salmonella choleraesuis               | Salmonella schottmuelleri  |
| Staphylococcus aureus                 | Salmonella typhi           |
| Acinetobacter baumannii               | Salmonella typhimurium     |
| Brevibacterium ammoniagenes           | Serratia marcescens        |
| Chlamydia psittaci                    | Shigella dysenteriae       |
| Enterobacter aerogenes                | Shigella flexneri          |
| Enterobacter cloacae                  | Shigella sonnei            |
| Enterococcus (Streptococcus) faecalis | Staphylococcus epidermidis |
| Escherichia coli                      | Streptococcus pyogenes     |
| Klebsiella pneumoniae                 |                            |

Escherichia coli - Methicillin resistant  
Enterococcus (Streptococcus) faecalis - Vancomycin-resistant (VRE)

Enterococcus (Streptococcus) faecalis-(Methicillin-resistant)  
Klebsiella pneumoniae - Methicillin resistant  
Staphylococcus aureus - Methicillin resistant (MRSA)  
Staphylococcus aureus - Vancomycin Intermediate Resistant (VISA)

Aspergillus niger  
Trichophyton mentagrophytes

|                        |                 |
|------------------------|-----------------|
| *HIV-1 (AIDS virus)    | *influenza A/PR |
| *Herpes Simplex Type 1 | *Rubella        |
| *Herpes Simplex Type 2 | *Vaccinia       |

**LONZA FORMULATION DC-110N** is a sanitizer for hard, nonporous, non-food, contact surfaces at 2 - 3 ounces per gallon (16 - 24 milliliters per liter) against:

Enterobacter aerogenes                      Staphylococcus aureus

**ACCEPTED**

DEC 28 2000

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide, registered under EPA Reg. No. 6836-204

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant / high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

**Preparation of use-solution:**

For water hardness up to 250 ppm, add 2 ounces per gallon (16 milliliters per liter) of water (1:64).  
For water hardness up to 400 ppm, add 3 ounces per gallon (24 milliliters per liter) of water (1:42), to disinfect hard, nonporous surfaces.  
For heavy duty use, add 8 ounces per gallon (64 milliliters per liter) of water (1:16), to disinfect hard, nonporous surfaces.

**Disinfection / \*Virucidal / Fungicidal directions:**

Apply use-solution to hard, (inanimate), nonporous surfaces thoroughly wetting surfaces with a cloth, mop, sponge or sprayer.

For heavily soiled areas, a preliminary cleaning is required.

For sprayer applications, use a coarse spray device. Spray 6-8 inches from surface, rub with brush, sponge or cloth. Do not breathe spray.

Treated surfaces must remain wet for 10 minutes (2 minutes for disinfection against all viruses claimed). Wipe dry with a clean cloth or sponge or allow to air dry. Prepare a fresh solution daily or when use solution becomes diluted or soiled.

Rinse all surfaces, which come in contact with food, such as countertops, appliances, tables and stovetops, with potable water before reuse. This product should not be used to clean utensils, glassware and dishes.

**Mold and Mildew Control directions:**

To control the growth of mold and mildew and their odors on hard, nonporous surfaces, apply use-solution to thoroughly wet all treated surfaces completely. Let air dry. Repeat application weekly or when growth or odor reappears.

4/5

**Sanitizing Non-Food, Contact Surface directions:**

Add 2 - 3 ounces per gallon (16 - 24 milliliters per liter) of water, to sanitize hard, nonporous, environmental surfaces. Treated surfaces must remain wet for 60 seconds. Wipe dry with a clean cloth or sponge or allow to air dry.

**\*KILLS HIV ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces / objects with blood or body fluids and in which the surfaces / objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 ON SURFACES /OBJECTS SOILED WITH BLOOD / BODY FLUIDS.**

**Personal protection:** Clean-up should always be done wearing protective latex gloves, gowns, masks and eye protection.

**Cleaning procedure:** blood and other body fluids containing HIV must be thoroughly cleaned from surfaces and objects before application of LONZA FORMULTION DC-110N.

**Contact time:** Leave surface wet for 2 minutes with 2 ounces per gallon (16 milliliters per liter) use-solution. Use a 10-minute contact time for disinfection against all other bacteria and fungi claimed.

**DISPOSAL OF INFECTIOUS MATERIALS:** Blood, body fluids, cleaning materials and clothing should be autoclaved and disposed of according to local regulations for infectious waste disposal.

**NON-ACID TOILET BOWL DISINFECTANT / CLEANER DIRECTIONS:**

**From concentrate:**

Add 2 - 3 ounces (60-90 milliliters) to the toilet bowl and mix; brush thoroughly over exposed surfaces and under the rim; allow to stand for 10 minutes and flush.

**From use-solution:**

Empty toilet bowl or urinal and apply 2 - 3 ounces per gallon (16 - 24 milliliters per liter) use-solution to exposed surfaces including under the rim with a cloth, mop, sponge or sprayer; allow to stand for 10 minutes and flush.

**For Heavy-duty use:**

Empty toilet bowl or urinal and apply 8 ounces per gallon (64 milliliters per liter) use-solution to exposed surfaces, including under the rim with a cloth, mop, sponge or sprayer; allow to stand for 10 minutes and flush.

**Glass Cleaning/Deodorizing Directions:**

Add 1 - 4 ounce per gallon (8 - 32 milliliters per liter) of water to clean and deodorize windows, mirrors and glass surfaces.


Use a coarse spray device. Spray 6-8 inches from surface; rub with sponge or cloth. Do not breathe spray.

**Directions for Food Processing Plants:**

For use on non-food, contact surfaces such as floors, walls, tables, garbage cans and disposal areas. Follow disinfection directions.

**Directions for Food Storage Areas:**

For use on non-food, contact surfaces such as shelves, floors, walls and tables: Before using this product, food products and packaging materials must be removed from the room or carefully protected, then follow disinfection directions.



5/5

**POULTRY and SWINE PREMISE SANITATION**

**Site preparation**

The first step in any on-going sanitation program should be the removal of gross contamination and debris. This may be accomplished using a shovel, broom, or vacuum depending on the area to be disinfected. The efficacy of even the most efficient germicidal cleaner is reduced in the presence of heavy organic matter. Once the heavy debris is eliminated, disinfection can be accomplished in one easy step.

**Cleaning**

For all general cleaning, use 2 - 3 ounces of **LONZA FORMULATION DC-110N** per gallon of water. Apply **LONZA FORMULATION DC-110N** using a cloth, mop or pressure sprayer so as to thoroughly wet surface to be disinfected. Allow to remain wet for 10 minutes and then let air dry. For heavily soiled areas, a pre-cleaning step is required. Prepare a fresh solution after each use.

**DISINFECTION**

**POULTRY and SWINE PREMISE**

Remove all animals and feeds from premises, trucks, coops and crates. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Use 2 - 3 ounces of **LONZA FORMULATION DC-110N** per gallon of water. Saturate surfaces with the recommended disinfecting solution for a period of 10 minutes. Wipe dry with a clean cloth or sponge or allow to air dry. Ventilate buildings, coops, and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels and scrapers used for removing litter and manure.

**Hatcheries:** Use 2 - 3 ounces of **LONZA FORMULATION DC-110N** per gallon of water to treat hatchers, setters, trays, racks, carts, sexing tables, delivery trucks and other hard surfaces. Leave all treated surfaces exposed to disinfectant solution wet for 10 minutes or more. Wipe dry with a clean cloth or sponge or allow to air dry.

**Vehicles:** Clean all vehicles including mats, crates, cabs and wheels with high pressure water and **LONZA FORMULATION DC-110N**. Use 2 - 3 ounces per gallon to treat all vehicles. Leave all treated surfaces exposed to disinfectant solution wet for 10 minutes or more. Wipe dry with a clean cloth or sponge or allow to air dry.

**Shoe Baths:** Shoe baths containing one inch of freshly made solution should be placed at all entrances to buildings and hatcheries. Scrape waterproof work boots and place in a 2 - 3 oz. per gallon of water solution for 60 seconds prior to entering the area. Change the solution in the bath daily or when solution appears dirty.

**Dressing Plant Use:** Disinfect equipment, utensils, walls and floors in poultry and animal dressing plants. Disinfect offal rooms, exterior walls and loading platforms of dressing plants. Cover or remove all food and packaging materials. Remove all gross soils. Saturate all surfaces with the recommended use solution of 2 ounces per gallon. Scrub to loosen all soils. Allow to soak for 10 minutes and thoroughly rinse all wetted and cleaned surfaces with potable water. Wipe dry with a clean cloth or sponge or allow to air dry.

**SANITIZING HATCHERY ROOMS USING FOGGING DEVICES**

Remove all animals and feed from premises, vehicles and enclosures. Remove all litter and manure from floors, walls and surfaces of the room to be treated. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Close room off so fog is confined to room to be treated. Mix four parts **LONZA FORMULATION DC-110N** to five parts water (2 gallons of **LONZA FORMULATION DC-110N** to 2.5 gallons water). Insert the nozzle of the fogger through a suitable opening into the room. With the setting in maximum output, fog for one minute for each 4000 cubic feet of space in the room. When fogging is completed, ventilate buildings, and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed or dried.

**SANITIZING INCUBATORS AND HATCHERS USING FOGGING DEVICES**

Mix 24 ounces **LONZA FORMULATION DC-110N** to 122 ounces water. Fog 3 - 8 ounces of this into setters and hatchers immediately after transfer. Repeat daily in setters and every 12 hours in hatchers. Discontinue hatcher treatments at least 24 hours prior to pulling the hatch. Do not allow people to contact or breathe this fog and do not enter until the fog has settled (30 - 60 minutes after fogging is completed). It is acceptable to fog setters and hatchers with a 2 - 3 ounces per gallon solution of **LONZA FORMULATION DC-110N** on an hourly or every other hour basis. If this is done, fog for 30 - 90 seconds once per hour or once every two hours. When fogging is completed, ventilate buildings, and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed or dried. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.

**NOTE:** The fog generated is irritating to the eyes, skin and mucous membranes. Under no circumstances should a room or building be entered by anyone within two hours of the actual fogging. If the building must be entered, then the individuals entering the building must wear a self contained respirator approved by NIOSH/MSHA, goggles, long sleeves and long pants.

**FOGGING IS TO BE USED AS AN ADJUNCT TO ACCEPTABLE MANUAL CLEANING AND DISINFECTING OF ROOM AND MACHINE SURFACES.**

**ACCEPTED**  
DEC 28 2008  
Under the Federal Insecticide, Fungicide, and  
Rodenticide Act as amended, for the  
pesticide, registered under  
EPA Reg. No. 6836-204