

6836-206

1/6/2010

10/11

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

JAN -6 2010

Joanna Holcombe
Sr. Commercial Regulatory Services Associate
Lonza Inc.
90 Boroline Road
Allendale, NJ 07401

Subject: Lonza Formulation DC-109N
EPA Reg. No. 6836-206
Application Dated: December 03, 2009
Receipt Date: December 07, 2009

Dear Joanna Holcombe

The following notification submitted in connection with registration under the provisions of PR Notice 98-10, Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) section 3(c)9. is acceptable.

Proposed Amendment

Adding Marketing language for the Pandemic 2009 H1N1

Comments

Based on a review of the material submitted, the following comments apply:
This application for notification to add marketing language for the Pandemic 2009 H1N1 is acceptable. A copy has been placed in our records for future reference.

Should you have any questions or comments concerning this letter, please contact Velma Noble at (703) 308-6233.

Sincerely,

Velma Noble
Product Manager (31)
Regulatory Management Branch1 Antimicrobia
Division (7510P)

CONCURRENCES

SYMBOL	SURNAME	DATE					

Via FedEx

Lonza Inc
Allendale
90 Boroline Road
Allendale, NJ 07401, USA

Ms. Velma Noble, PM-31
Document Processing Desk (NOTIFY)
Office of Pesticide Programs (7504P)
US Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 S. Crystal Drive
Arlington, VA 22202

Joanna Holcombe
Lonza Life Science Ingredients
Commercial Regulatory Services

Tel 201-316-9297
Fax 201 696-3470
joanna.holcombe@lonza.com

December 3, 2009

**SUBJECT: Lonza Formulation DC-109N, EPA Registration No. 6836-206
Application for Label Notification**

Dear Ms. Noble:

The purpose of this notification is to add the following:

**Add optional marketing language per EPA Guidance for Testing and Labeling Claims
against Pandemic 2009 H1N1.**


Please find enclosed the following documents in support of this label notification.

- Application for Pesticide Notification
- One copy of the label with changes highlighted

Please provide your decision electronically to me at joanna.holcombe@lonza.com and subsequently send me a letter confirming the Agency's decision. If you have any questions or need any additional information, please feel free to contact me at 201-316-9297.

Sincerely,
Lonza Inc.

Joanna Holcombe
Sr. Commercial Regulatory Services Associate



4 of 11

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER. Corrosive. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin or on clothing. Wear protective clothing, eyewear (goggles, face shield or safety glasses) and protective (rubber, chemical resistant) gloves when handling. May be fatal if swallowed. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

Call a poison control center or doctor for treatment advice. Have the container or label with you when calling a poison control center or doctor, or going for treatment.

(Note to reviewer: If the container size is 5 gallons or greater, the following Environmental Hazards statements will be used:)

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

**LONZA FORMULATION
DC-109N**

Disinfectant - Cleaner - Sanitizer - Fungicide -
Mildewstat - Virucide* - Deodorizer
for Hospitals, Institutional and Industrial Use

Effective in hard water up to 400 ppm hardness
(Calculated as CaCO₃) in the presence of 5% serum
contamination

Fresh / Clean / Lemon / Pine / Mint / Floral / Citrus
(scent) (fragrance) (scented) (odor)

Formulated for effective poultry premise sanitation
Formulated for effective swine premise sanitation

Active Ingredient:	
Didecyl dimethyl ammonium chloride.....	7.68%
Other Ingredients:.....	92.32%
Total:.....	100.00%


KEEP OUT OF REACH OF CHILDREN

DANGER PELIGRO

See (side) (left) (right) (back) panel for Precautionary
Statements and First Aid.

EPA Reg. No. 6836-206
EPA Est. No. (as noted on the container)
Net Contents (as noted on the container)
[Country of origin (insert country)]
[Manufactured in (insert country)]
[Barcode]

Manufactured by:
LONZA INC.
90 Boroline Road, Allendale, NJ 07401



Note to reviewer: The following is considered optional marketing language:

Concentrate (Concentrated)
Disinfects (Disinfectant)
Cleans (Cleaner)
Deodorizes (Deodorant)

Cross-contamination is of major housekeeping concern not only in hospitals, but also in schools, institutions and industry. **LONZA FORMULATION DC-109N** has been formulated to aid in the reduction of cross-contamination from treated surfaces in these areas.

LONZA FORMULATION DC-109N:

- Is a one-step disinfectant cleaner that is effective against a broad spectrum of bacteria, is virucidal* (including HIV-1 and HBV), fungicidal, and inhibits the growth of mold and mildew and their odors when used as directed.
- is an effective one-step sanitizer-cleaner for use on non-food contact surfaces.
- will deodorize surfaces in restrooms and toilet areas, behind and under sinks and counters, garbage cans and garbage storage areas, and other places where bacterial growth can cause mal odors.
- Is an effective fungicide against Trichophyton mentagrophytes (the athlete's foot fungus) when used on surfaces in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities.
- contains no (phosphates or) phosphorous (compounds).
- is a no-rinse neutral pH disinfectant cleaner that disinfects, cleans and deodorizes in one labor-saving step.
- can be used with a mop and bucket, trigger sprayers, sponge, or by soaking.
- provides effective cleaning strength that will not dull most metal-interlock floor finishes, and does not require a rinse prior to recoat.
- improves labor results by effectively controlling odors.
- cleans, shines, deodorizes and disinfects hard nonporous household surfaces. Inhibits the growth mold and mildew, leaving bathrooms & kitchens clean and smelling fresh.
- kills household germs; effective against Pseudomonas aeruginosa (Pseudomonas), Staphylococcus aureus (Staph) and Streptococcus faecalis (Strep), Influenza A/PR Virus and Trichophyton mentagrophytes (Athlete's foot fungus) (Cause of Ringworm).
- is a neutral pH multi-surface cleaner, deodorizer, and disinfectant. Use on windows, mirrors, and other glass surfaces.

Use LONZA FORMULATION DC-109N


- for non-scratch cleaning of showers and tubs, shower doors and curtains, fixtures, and toilet bowls.
- in kitchens, bathrooms and other household areas.
- on the multi-touch surfaces responsible for cross-contamination.
- on glass surfaces including windows and mirrors.

LONZA FORMULATION DC-109N is recommended for Poultry Premise Sanitation (Hatcheries):

Chick Processing Area	Chick Holding Room
Egg Receiving Area	Egg Holding Area
Chick Loading Area	Tray Dumping Area
Setter Room	Hatchery Room
Poultry Buildings	

LONZA FORMULATION DC-109N is recommended for Swine Premise Sanitation:

Farrowing Barns & Areas	Dressing Plants
Blocks	Waterers and Feeders
Loading Equipment	Creep Area
Hauling Equipment	Nursery
Chutes	



(Note to reviewer: The following is considered optional marketing language.)

(AREAS OF USE:)

Use Lonza Formulation DC-109N in:

Bathrooms
Homes (households)
Kitchens

Dental offices
Hospitals
Medical Offices
Nursing homes

Day care centers
Nurseries

Bars
Cafeterias
Convenience stores
Food processing plants
Food storage areas
Institutional kitchens
Restaurants
USDA inspected food processing facilities

Athletic facilities
Barber shops
Business and office buildings
Colleges
Correctional facilities
Dressing rooms
Exercise facilities
Factories
Hotels
Institutional facilities
Institutions
Locker rooms
Motels
Prisons
Public facilities
Public rest rooms
Schools
Shower and bath areas
Salons

Camp grounds
Play ground equipment

Animal laboratories
Dairy farms
Hog farms
Kennels
Poultry farms
Pet animal quarters
Pet shops
Turkey farms
Zoos

Airports
Airplanes
Boats
Buses
Camps
Cars
Emergency vehicles
Mobile homes
Ships
Taxis
Trailers
Trains
Transportation terminals

(TYPES OF SURFACES:)

Use Lonza Formulation DC-109N on washable hard, nonporous surfaces of:


Appliances
Bathroom bowls
Bathroom basins
Bathroom fixtures
Bathtubs
Cabinets
Cages
Chairs
Coils and drain pans of air conditioning and refrigeration equipment and heat pumps
Conductive flooring
Counters (countertops)
Countertop laminates
Desks
Doorknobs
Door handles
Fiberglass bathtubs
Fiberglass sinks
Floors
Floors, finished
Floors, high speed burnished
Garbage cans
Highchairs
Kennel runs
Microwave ovens (Exterior Surfaces)
Outdoor furniture

Refrigerators (Exterior Surfaces)
Refrigerated storage and display equipment

Showers
Shower stalls
Sinks (bathroom, kitchen)
Stoves (stovetops)
Tables, (picnic tables)
Telephones
Toilets
Toilet bowls
Toilet bowl surfaces
Tubs
Tiles, glazed
Ultrasonic baths
Urinals
Walls
Whirlpool bathtubs

Other hard nonporous surfaces made of:

Formica®
Glazed ceramic
Glazed enameled surfaces
Glazed porcelain
Metal
Plastic (such as polystyrene or polypropylene)
Sealed stone
Stainless steel
Upholstery, vinyl and plastic
Woodwork, finished



LONZA FORMULATION DC-109N is a Hospital Disinfectant at 1 oz. per gallon (8 mL per liter, 1:128 dilution), modified in the presence of 250 ppm hard water, and at 1-1/3 oz. per gallon (10 mL per liter, 1:100 dilution), modified in the presence of 400 ppm hard water.

LONZA FORMULATION DC-109N is Bactericidal according to the AOAC Use-Dilution Test method, Virucidal* according to the virucidal qualification on hard inanimate surfaces, Fungicidal according to the AOAC fungicidal test, modified in the presence of 5% organic serum against:

(Bacteria:)

- Pseudomonas aeruginosa (Pseudomonas)
- Salmonella enterica (Salmonella)
- Staphylococcus aureus (Staph)

- Acinetobacter baumannii
- Brevibacterium ammoniagenes
- Chlamydia psittaci
- Enterobacter aerogenes
- Enterobacter cloacae
- Enterococcus (Streptococcus) faecalis-Methicillin-resistant
- Enterococcus faecalis - Vancomycin-resistant (VRE)
- Escherichia coli (E. coli)
- Escherichia coli - Methicillin resistant
- Enterococcus (Streptococcus) faecalis
- Klebsiella pneumoniae
- Klebsiella pneumoniae - Methicillin resistant)
- Proteus vulgaris
- Salmonella schottmuelleri
- Salmonella typhi
- Salmonella typhimurium
- Serratia marcescens
- Shigella dysenteriae
- Shigella flexneri
- Shigella sonnei
- Staphylococcus aureus - Methicillin resistant (MRSA)
- Staphylococcus aureus - Vancomycin Intermediate Resistant (VISA)

- Staphylococcus epidermidis
- Streptococcus pyogenes (Strep)

(Viruses:)

- *Hepatitis B Virus
- *Herpes Simplex Virus Type 1
- *Herpes Simplex Virus Type 2
- *HIV-1 (AIDS virus)
- *Influenza A/PR Virus (Influenza)
- *Rubella (Causative agent of German measles)
- *Vaccinia Virus

(Fungi:)

- Aspergillus niger
- Candida albicans
- Trichophyton mentagrophytes (Athlete's Foot Fungus)
(Cause of Ringworm)

LONZA FORMULATION DC-109N is a sanitizer for hard, nonporous, non-food, contact surfaces at 1 - 1 1/3 oz. per gallon (8 - 10 mL per liter) against:

- Enterobacter aerogenes
- Staphylococcus aureus

(Note to reviewer: the following is considered optional marketing language:)

- Respiratory illnesses attributable to Pandemic 2009 H1N1 are caused by influenza A virus. This product (Product Name) is a broad-spectrum hard surface disinfectant that has been shown to be effective against (influenza A virus tested and listed on the label) and is expected to inactivate all influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).
- This product has demonstrated effectiveness against influenza A virus and is expected to inactivate all influenza A viruses including Pandemic 2009 H1N1 influenza A virus.
- This product has demonstrated effectiveness against (influenza A virus tested and listed on the label) and is expected to inactivate all influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).
- Kills Pandemic 2009 H1N1 influenza A virus (formerly called swine flu).
- Kills Pandemic 2009 H1N1 influenza A virus.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Preparation of use-solution:

- For water hardness up to 250 ppm, add 1 oz. per gallon (8 mL per liter) of water (1:128).
- For water hardness up to 400 ppm, add 1 1/3 oz. per gallon (10 mL per liter) of water (1:100).
- For heavy duty use, add 4 oz. per gallon (32 mL per liter) of water (1:32).

DISINFECTION / *VIRUCIDAL / FUNGICIDAL DIRECTIONS:


Apply use-solution to hard nonporous surfaces thoroughly wetting the surfaces with a cloth, mop, sponge or sprayer.

Treated surfaces must remain wet for 10 minutes (2 minutes for disinfection against all viruses claimed except HBV). Wipe dry with a clean cloth or sponge or allow to air dry. For heavily soiled areas, a preliminary cleaning is required.

For sprayer applications, use a coarse spray device. Spray 6 - 8 inches from surface; rub with brush, sponge or cloth. Do not breathe spray.

Prepare a fresh solution daily or more often if the use solution becomes diluted or soiled.

Rinse all surfaces, which come in contact with food, such as countertops, appliances, tables, and stovetops, with potable water before reuse. Do not use on utensils, glassware and dishes.



(Note to reviewer: For labels that list medical premises and metal and / or stainless steel surfaces, one of the following statements must be used.):

This product is not to be used as a terminal sterilant / high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

(or)

This product is not for use on medical device surfaces.

MOLD AND MILDEW CONTROL DIRECTIONS:

To control the growth of mold and mildew and their odors on hard, nonporous surfaces, apply use-solution to thoroughly wet all treated surfaces completely. Let air dry. Repeat application weekly or when growth or odor reappears.

***KILLS HIV AND HBV ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces / objects with blood or body fluids and in which the surfaces / objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) or Hepatitis B Virus (HBV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 OR HBV ON SURFACES / OBJECTS SOILED WITH BLOOD / BODY FLUIDS.

Personal protection: Clean-up should always be done wearing protective gloves, gowns, masks and eye protection.

Cleaning procedure: Blood and other body fluids containing HIV or HBV must be thoroughly cleaned from surfaces and objects before application of Lonza Formulation DC-109N.

Contact time:

For HIV: Use 1 oz. of Lonza Formulation DC-109N per gallon of water (8 mL per liter). Leave surfaces wet for 2 minutes

For HBV: Use 1 1/3 oz. of Lonza Formulation DC-109N per gallon of water (10 mL per liter). Leave surfaces wet for 10 minutes.

For disinfection against all other bacteria and fungi claimed: Use a 10 minute contact time.

Disposal of infectious materials: Blood, body fluids, cleaning materials and clothing should be autoclaved and disposed of according to local regulations for infectious waste disposal.

NON-ACID TOILET BOWL AND URINAL DISINFECTANT / CLEANER DIRECTIONS:

Remove heavy soil prior to disinfection.

From concentrate:

Add 1 - 1 1/3 oz. (30 - 40 mL) to the toilet bowl and mix; brush thoroughly over exposed surfaces and under the rim with a cloth, mop or sponge. Allow to stand for 10 minutes and flush.

From use-solution:

Empty toilet bowl or urinal and apply 1 - 1 1/3 oz. per gallon (8 - 10 mL per liter) use-solution to exposed surfaces including under the rim with a cloth, mop, sponge or sprayer.

Brush or swab thoroughly. Allow to stand for 10 minutes and flush.

For Heavy-duty use:

Empty toilet bowl or urinal and apply a 4 oz. per gallon (32 mL per liter) use-solution to exposed surfaces, including under the rim with a cloth, mop, sponge or sprayer. Brush or swab thoroughly. Allow to stand for 10 minutes and flush.

GLASS CLEANING/DEODORIZING DIRECTIONS:

Add 1/2 - 2 oz. per gallon (4 - 16 mL per liter) of water to clean and deodorize windows, mirrors and glass surfaces.

For sprayer applications, use a coarse spray device. Spray 6 - 8 inches from surface; rub with sponge or cloth. Do not breathe spray.

SALON (BARBER) INSTRUMENTS AND TOOLS DISINFECTION (BACTERICIDE) / VIRUCIDE / FUNGICIDE DIRECTIONS:


Mix 1 - 1 1/3 oz. of Lonza Formulation DC-109N per gallon of water (or equivalent dilution) to disinfect hard, nonporous instruments and tools. Completely immerse combs, brushes, rollers, razors, scissors, blades, manicure and other salon instruments and tools for 10 minutes. For heavily soiled instruments and tools, a preliminary cleaning is required. Rinse thoroughly and dry before use. Prepare a fresh solution daily or more often if the solution becomes visibly diluted, cloudy or soiled.

Note: Plastics may remain immersed until ready to use. Stainless steel shears and instruments must be removed after 10 minutes, rinsed, dried and kept in a clean, non-contaminated receptacle. Prolonged soaking may cause damage to metal instruments.

ULTRASONIC BATH DISINFECTANT DIRECTIONS:

Use Lonza Formulation DC-109N to disinfect hard nonporous non-critical instruments / objects compatible with Ultrasonic cleaning units.

Pour fresh use-solution of 1 - 1 1/3 oz. of Lonza Formulation DC-109N per gallon of water (or equivalent dilution) directly into bath chamber. Preclean soiled objects. Place objects into unit and operate for a minimum of 10 minutes, according to manufacturer's use directions. Remove objects and rinse with sterile water (sterile water for injection), or allow to air dry. Prepare a fresh solution for each use.



WHIRLPOOL BATH DISINFECTION DIRECTIONS:

After using whirlpool unit, drain and fill with a use solution of 1 - 1 1/3 oz. of **Lonza Formulation DC-109N** per gallon of water to just cover the intake valve. Start the pump to circulate the solution. Wash down the unit sides, seat of the chair lift, and any related equipment with a clean swab, brush or sponge. Treated surfaces must remain wet for 10 minutes. After the unit has been thoroughly disinfected, drain the solution from the unit and rinse disinfected surfaces with fresh water.

Repeat for heavily soiled units.

DIRECTIONS FOR FOOD PROCESSING PLANTS:

For use on non-food, contact surfaces such as floors, walls, tables, garbage cans and disposal areas. Before using this product, food products and packaging materials must be removed or carefully protected. Follow disinfection directions.

DIRECTIONS FOR FOOD STORAGE AREAS:

For use on non-food, contact surfaces such as shelves, floors, walls and tables: Before using this product, food products and packaging materials must be removed from the room or carefully protected, then follow disinfection directions.

DIRECTIONS FOR SANITIZING NON-FOOD CONTACT SURFACES:

Add 1 - 1 1/3 oz. per gallon (8 - 10 mL per liter) of water, to sanitize hard, nonporous, environmental surfaces. Treated surfaces must remain wet for 60 seconds. Wipe dry with a clean cloth or sponge or allow to air dry.

POULTRY AND SWINE PREMISE DISINFECTION DIRECTIONS

SITE PREPARATION:

The first step in any on-going sanitation program should be the removal of gross contamination and debris. This may be accomplished using a shovel, broom, or vacuum depending on the area to be disinfected. The efficacy of even the most efficient germicidal cleaner is reduced in the presence of heavy organic matter. Once the heavy debris is eliminated, disinfection can be ACCOMPLISHED IN ONE EASY STEP.


CLEANING:

For all general cleaning, use 1 - 1 1/3 oz. of **LONZA FORMULATION DC-109N** per gallon of water (1:128 solution). Apply **LONZA FORMULATION DC-109N** using a cloth, mop, or pressure sprayer so as to thoroughly wet surfaces. Wipe dry with a clean cloth or sponge or allow to air dry.

DISINFECTION:

Remove all animals and feeds from premises, vehicles, trucks, and enclosures such as coops and crates. Remove all litter, droppings and manure from floors, walls and surfaces of barns, pens, stall, chutes and other facilities occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Use 1 - 1 1/3 oz. of **LONZA FORMULATION DC-109N** per gallon of water. Saturate surfaces with the recommended disinfecting solution for a period of 10 minutes. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels and scrapers used for removing litter and manure. Wipe dry with a clean cloth or sponge or allow to air dry. Ventilate buildings, coops and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.

HATCHERIES: Use 1 - 1 1/3 oz. of **LONZA FORMULATION DC-109N** per gallon of water to treat hatchers, setters, trays, racks, carts, sexing tables, delivery trucks and other hard surfaces. Leave all treated surfaces wet for 10 minutes or more. Wipe dry with a clean cloth or sponge or allow to air dry.



VEHICLES: Clean all vehicles including mats, crates, cabs, and wheels with high pressure water and **LONZA FORMULATION DC-109N**. Use 1 - 1 1/3 oz. per gallon to treat all vehicles. Leave all treated surfaces wet for 10 minutes or more. Wipe dry with a clean cloth or sponge or allow to air dry.

SHOE BATH SANITIZER DIRECTIONS: Shoe baths containing one inch of freshly made solution should be placed at all entrances to buildings and hatcheries. Scrape waterproof work boots and place in a 1 to 1 1/3 oz. per gallon of water solution for 60 seconds prior to entering the area. Change the solution in the bath daily or more often if the solution appears diluted or soiled.

DRESSING PLANT DISINFECTION DIRECTIONS: Disinfect equipment, utensils, walls and floors in poultry and animal dressing plants. Disinfect offal rooms, exterior walls and loading platforms of dressing plants. Cover or remove all food and packaging materials. Remove all gross soils. Saturate all surfaces with the recommended use solution of 1 oz. per gallon. Scrub to loosen all soils. Allow to soak for 10 minutes and thoroughly rinse all wetted and cleaned surfaces with potable water. Wipe dry with a clean cloth or sponge or allow to air dry.

SANITIZING HATCHERY ROOMS USING FOGGING DEVICES:

Remove all animals and feed from premises, vehicles and enclosures. Remove all litter and manure from floors, walls and surfaces of the room to be treated. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Close room off so fog is confined to room to be treated. Wear a dust mist respirator when mixing the use solution and pouring it into the fogging apparatus. Mix two parts **LONZA FORMULATION DC-109N** to five parts water (1 gallon of **LONZA FORMULATION DC-109N** to 2.5 gallons water). Insert the nozzle of the fogger through a suitable opening into the room. With the setting in maximum output, fog for one minute for each 4000 cubic feet of space in the room. When fogging is completed, ventilate buildings, and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed or dried. Thoroughly scrub all treated feed racks, troughs, automatic feeders, fountains and waters with soap or detergent, and rinse with potable water before reuse.

SANITIZING INCUBATORS AND HATCHERS USING FOGGING DEVICES


Wear a dust mist respirator when mixing the use solution and pouring it into the fogging apparatus. Mix 12 oz. of **LONZA FORMULATION DC-109N** to 122 oz. water. Fog 3 - 8 oz. of this into setters and hatchers immediately after transfer. Repeat daily in setters and every 12 hours in hatchers. Discontinue hatcher treatments at least 24 hours prior to pulling the hatch. Do not allow people to contact or breathe this fog and do not enter until the fog has settled (30 - 60 minutes after fogging is completed). It is acceptable to fog setters and hatchers with a 1 - 1 1/3 oz. per gallon solution of **LONZA FORMULATION DC-109N** on an hourly or every other hour basis. If this is done, fog for 30 - 90 seconds once per hour or once every two hours. When fogging is completed, ventilate buildings, and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed or dried. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.

NOTE: The fog generated is irritating to the eyes, skin and mucous membranes. Under no circumstances must a room or building be entered by anyone within two hours of the actual fogging and a minimum of 4 air exchanges (ACH) per hour in the facility. If the building must be entered, then the individuals entering the building must wear a self contained respirator approved by NIOSH/MSHA, goggles, long sleeves and long pants.

FOGGING IS TO BE USED AS AN ADJUNCT TO ACCEPTABLE MANUAL CLEANING AND DISINFECTING OR ROOM AND MACHINE SURFACES.

VETERINARY PRACTICE / ANIMAL CARE / ANIMAL LABORATORY / ZOOS / PET SHOP / KENNELS DISINFECTION DIRECTIONS:

For cleaning and disinfecting the following hard nonporous surfaces: equipment used for animal food or water, utensils, instruments, cages, kennels, stables and catteries. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use-solution of 1 to 1 1/3 oz. of **LONZA FORMULATION DC-109N** per gallon of water (or equivalent dilution) for a period of 10 minutes. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals as well as forks, shovels, and scrapers used for removing litter and manure. Wipe or allow to air dry. Rinse all surfaces that come in contact with food, including equipment used for feeding or watering, with potable water before reuse. Ventilade buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried.



FARM PREMISE USE:

Remove all animals and feed from premises, vehicles and enclosures. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean surfaces with soap or detergent and rinse with water. Saturate surfaces with a use-solution of 1 to 1 1/3 oz. of **LONZA FORMULATION DC-109N** per gallon of water (or equivalent dilution) for a period of 10 minutes. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels and scrapers used for removing litter and manure. Ventilate buildings, cars, trucks, boats and other closed spaces.

Do not house livestock or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.

(Note to reviewer: For Nonrefillable Containers for commercial, industrial, and institutional uses – all sizes – No Reuse Rinsate Statement for Public Health Use products)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Open dumping is prohibited. Store in original container in areas inaccessible to children.

Pesticide Disposal:

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal:

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying.

(Plastic and Metal Containers:) Triple rinse as follows: Fill container 1/4 full with water and reclose the container. Agitate vigorously, and dispose of rinsate consistent with pesticide disposal instructions. Repeat two more times. Then offer for recycling if available or puncture and dispose in sanitary landfill or by other procedures approved by state and local authorities. Follow pesticide disposal instructions for rinsate. If not triple rinsed, these containers are acute hazardous wastes and must be disposed in accordance with local, state, and federal regulations.

(For metal containers only: DO NOT cut or weld metal containers.)

(Bag in Box Containers:) Completely empty bag into application equipment. Then offer for recycling if available or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

