

NET WEIGHT 10 LBS

BES-TEX

GOLD ★ STOCK

DUSTING SULPHUR

CAUTION: KEEP OUT OF REACH OF CHILDREN

DO NOT use sulphur within 2 weeks of an oil spray. **DO NOT** use sulphur on apples or other sensitive crops when temperatures exceed or are likely to exceed 90 degrees F. Sulphur may cause severe fruit and foliage injury to certain crops. **DO NOT** use sulphur on apricots, Anjou and Comice pears, cranberries, cucurbits (cucumbers, squash, and some varieties of melons), filberts, spinach, tung trees, walnuts and other sensitive plants. When growing crops for processing, consult the processor before applying sulphur. **DO NOT** apply when weather conditions favor drift from areas treated. **DO NOT** contaminate water by cleaning of equipment or disposal of wastes.

USE ONLY FOR USE

BEANS AND PEAS

BES-TEX INSECTICIDES CO., INC

NET WEIGHT 50 LBS.

ALL L. A. A. U.
+ B. C. S. 12X
G. L. D.



GOLD ★ STOCK

ACTIVE INGREDIENT:
Sulphur 93.00%
INERT INGREDIENTS 7.00%
TOTAL 100.00%

DUSTING SULPHUR

CAUTION: KEEP OUT OF REACH OF CHILDREN



DIRECTIONS FOR USE

Apply this product only as specified on this label. Gold Stock Dusting Sulphur is formulated for use in dust mixtures or alone as a dust when a 325 mesh dust is required. For best results apply when wind is calm. Dust often enough to keep foliage and fruit covered with a light uniform coating.

APPLES: To control Scab and powdery mildew, apply 30 to 40 lbs. per acre in the green tip to pink bud stage. Use on sulphur tolerant varieties only.

PEACH AND NECTARINE: To control Brown rot, coryneum blight, rust, peach silver mite and powdery mildew apply 50 to 60 lbs. per acre from popcorn stage to harvest.

PLUM AND PRUNE: To control Brown rot, leaf spot, and prune rust apply 35 to 50 lbs. per acre starting 5 weeks before harvest and continuing at weekly intervals.

BEANS AND PEAS: To control Powdery mildew, rust, and spider mites apply 15 to 30 lbs. per acre. For rust control start applications as soon as the first true leaves form and continue at weekly intervals until blossom time. NOTE: May cause injury to some varieties of beans.

COTTON: To control Atlantic mite and cotton rust apply 45 to 50 lbs. per acre. For rust control, start applications as soon as the first true leaves form and continue at weekly intervals until blossom time.

MELON (HONEYDEW AND SULPHUR-RESISTANT MELONS ONLY): To control Powdery mildew and spider mites apply 20 to 25 lbs. per acre.

PEANUTS: To control Rust and leaf spot apply 15 to 30 lbs. per acre.

CITRUS: To control Six spotted and citrus rust mites apply 80 lbs. per acre. 1/2 to 1 1/2

lbs. per tree, depending on size, for citrus rust mite in Florida.) To control Citrus flat and Yuma spider mites apply 100 lbs. per acre. To control Citrus thrips apply 100 lbs. per acre in 3 applications — mid-March — petal fall — and 2 to 4 weeks later.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

WARRANTY: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

Manufactured for
BES-TEX INSECTICIDES CO., INC.

San Angelo, Texas — EPA Est. No. 6853
EPA Registration No. 6853-19

Consult with your State Agricultural Experiment Station or your State Agricultural Extension Service for additional information consistent with this label cautions and directions.

White
B.A.A.H.

X