

READ THOROUGHLY AND UNDERSTAND ENTIRE LABEL BEFORE USING THIS PRODUCT

**DIRECTIONS FOR USE**

**COTTON:** Drill Kleen is useful as a spray application to weeds in cotton for post-emergent control of weeds and grasses, including Johnsongrass, nutgrass, watergrass, goosegrass, dallisgrass, cockspur, pigweed and renostergrass.

Mix at a rate of 1/2 gallon in 50 gallons of water. Apply this 50 gallons of solution on 3 to 4 acres on a band width of 12 to 15 inches on cotton grown on 40 inch rows. A second application may be required about 1 to 3 weeks after the first application. Do not make more than two applications per crop season.

APPLY ONLY WHEN COTTON IS 6 INCHES HIGH TO FIRST BLOOM. DO NOT APPLY AFTER FIRST BLOOM. DO NOT FEED TREATED FOLIAGE TO LIVESTOCK, OR GRAZE TREATED AREAS.

**MIXING INSTRUCTIONS**

THE CONTENTS OF THIS CONTAINER MUST BE SHAKEN THOROUGHLY BEFORE EACH USE UNTIL A UNIFORM MIXTURE IS OBTAINED. The thoroughly shaken Drill Kleen must be mixed with water until a uniform solution is obtained. Fill the spray equipment reservoir about half full with water and add the required amount of Drill Kleen with agitation. Finish filling the reservoir with water and apply. Continue to agitate thoroughly during application.

If the spray equipment is shut off when it still contains some Drill Kleen solution, agitate thoroughly before applying again. Do not store spray solution in reservoir or other tank for a prolonged period and clean equipment thoroughly after use by flushing with water.

**APPLICATION METHODS**

Drill Kleen spray solutions should be applied with a low volume sprayer having satisfactory pumping and bypass action. Nozzles should be placed so as to avoid spraying the cotton, but also so as to give good coverage of the weeds and grasses. Proper coverage is very important.

ADJUST NOZZLES TO DIRECT THE SPRAY MIST IN A MANNER TO ALLOW MAXIMUM COVERAGE OF WEEDS. KEEP SPRAY OFF COTTON FOLIAGE.

NET CONTENTS 1 GAL.

# DRILL KLEEN

For Post-Emergent Weed Control in Cotton

**INGREDIENTS**

Active Ingredient:		
Monosodium Acid Methanesulfonate	27.0%	
3-(3,4-dichlorophenyl)-1,1-dimethylurea (Diuron)	8.4%	
Inert Ingredients:	64.6%	
Total	100.0%	
Total arsenic, all in water soluble form, expressed as elemental	12.3%	

Contains 3.3 lbs. Monosodium Acid Methanesulfonate (Equiv. to 6 lbs. DSMA) and 0.9 lbs. Diuron with Surfactant per gallon.

## WARNING

**KEEP OUT OF REACH OF CHILDREN**

(See right panel for further warning statements.)

MANUFACTURED FOR

**farm chemical services**  
MONROE, LOUISIANA

**PRECAUTIONS**

Do not contaminate waters used by wildlife, aquatic life, domestic animals or for irrigation or domestic water supplies. Do not graze treated areas or feed livestock treated foliage.

**WARNING**

Harmful if taken internally. Avoid contact with skin. Avoid breathing dust or spray mist. Wash thoroughly after using. Store in a safe place away from feed and food products. Keep children and domestic animals off treated areas until this product has been washed into the soil.

**ANTIDOTE: IF TAKEN INTERNALLY, INDUCE VOMITING AND CALL PHYSICIAN AT ONCE. DO NOT CONTAMINATE WATERS USED FOR DOMESTIC CONSUMPTION OR BY ANIMALS, WILDLIFE AND AQUATIC LIFE, OR WATER USED FOR IRRIGATION PURPOSES.**

Use of this product in a manner, or at any time other than in accordance with the direction may produce plant injury, excessive residues or other undesirable results.

**NOTICE**

Seller's only guarantee, warranty, claim or representation is that this material conforms in content to that specified on the label. Climatic, soil and other conditions beyond Seller's control, necessarily affect results. Therefore, since this product will be used without knowledge or control of Seller, Seller shall not be responsible or liable for injuries to crops, persons or property which arise from use of or handling of this product, whether used in accordance with directions or not. Buyers accept and use this product under these conditions.

3

5