SANTEE

CLEANER · DISINFECTANT · DEODORIZER · FUNGICIDE

SANTEE is a heavy duty cleaner and disinfectant designed for general purpose use where heavy duty cleaning and sanitizing is of prime importance. When used as directed, SANTEE will remove most household dirt and grime in a single, simple application.

ACTIVE INGREDIENTS:

n-Alkyl (60% C_{14} , 30% C_{16} , 5% C_{12} , 5% C_{18}) dimethyl	
benzyl ammonium chlorides	0.8%
n-Alkyl (68% C ₁₂ , 32% C ₁₁) dimethyl ethylbenzyl	
ammonium chlorides	0.8%
Sodium Metasilicate	2.4%
Tetrasodium ethylenediamine tetraacetate	1.0%
INERT INGREDIENTS:	95.0%
	100.0%

PHENOL COEFFICIENTS: GUARANTEED MINIMUM

S. aureus 15 S. typhosa '9

DIRECTIONS FOR USE:

For light duty cleaning, sanitizing and deodorizing, use 2 ounces per gallon.

For heavy duty cleaning and disinfecting and fungicidal activity against pathogenic fungi, use $4\frac{1}{2}$ ounces per gallon.

For best results, use SANTEE with a cloth or mop and apply to walls, floors and other hard surfaces. If used for sanitization of food processing equipment or food contact items, limit the level of active quaternary to 200 ppm (8 ounces per 5 gallons). Food contact surfaces must be rinsed with potable water prior to reuse.

For extra heavy industrial cleaning and sanitizing, use a second application.

Do not mix with soap or anionic solutions.

SANTEE is non-corrosive in use dilutions on inanimate hard surfaces.

Not recommended for Pseudomonas aeruginosa.

DANGER: KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling concentrate. Harmful if swallowed. Avoid contamination of food.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

EPA Est 6869-NC-1 EPA Reg. No 6869-9 Do not reuse empty container. Return drums to drum reconditioner or destroy by perforating or crushing and burying in a safe place away fom water supplies.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg white, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.