


REVISED LABEL

S P E R L O X - S Z *
INSTANTLY DISPERSIBLE FUNGICIDE



Active Ingredient

Sulfur
Zineb (Zinc ethylenebisdithiocarbamate) 8%
(Zinc equivalent as metallic . . . (1.89%)

Inert Ingredients 52%

CAUTION: Keep out of reach of children . . . May cause irritation of eyes, nose, throat, and skin . . . May be harmful if inhaled or swallowed . . . Avoid breathing dust or spray mist . . . Avoid contact with skin, eyes, and clothing . . . Keep away from fire and sparks.

In case of contact, flush with plenty of water; for eyes, get medical attention.

Do not use treated peanut hay for feed.

IMPORTANT: Never allow Sperlox-SZ to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of Sperlox-SZ as a fungicide and create vapors which may be flammable. Keep container closed when not in use.

DIRECTIONS FOR USE:

Pour recommended amount of this material on surface of water in nearly filled tank with the agitator or by-pass running. Add balance of water. This material is compatible with Sperlox Brand insecticides and fungicides. Do not combine with wettable powders and emulsifiable liquids unless previous use has proved them to be physically compatible. Use screen not finer than 50 mesh in the sprayer.

RECOMMENDATIONS:

PEANUTS: Cercospora Leaf Spot - Apply 2 1/2 lbs. (early season) - 5 lbs. (late season) in 10 gallons of water per acre. Spray to give thorough coverage of the foliage. Begin application at first appearance of leaf spot and repeat at 10 to 14 day intervals until harvest.

NOTICE:

Use of this material other than in accordance with directions on this label may produce plant injury, excessive residues, or other undesirable results. Dawson Chemical Company makes no express or implied warranties other than specified on this label.

USDA Reg. No. 6874-5

Net Weight - 25 Lbs.

DAWSON CHEMICAL COMPANY
1525 N. Post Oak Road
Houston, Texas 77055

* Trademark

Produced in Mexico

Patent Pending