

Directions for Use

— RATS —

Place 4 to 6 ounces of bait in dry containers next to burrows, in runways or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 12 days and continue baiting until all signs of feeding have stopped. Replace stale, damp or moldy baits with fresh bait.

— MICE —

- Tablespoon amounts (1/4 to 1/2 ounce) of bait should be placed at 8 - 12 foot intervals.
- Bait should be placed in locations not accessible to children, pets, and domestic animals, or in tamper-proof bait boxes.
- An uninterrupted supply of bait should be maintained for at least 15 days.

ABSOLUTELY GUARANTEED:

If you are not entirely satisfied after using this package, return to dealer for full refund of purchase price.

USDA Registration Number - 6877-3



KILL RATS and MICE
with New Improved

NU-BRO RAT BAIT

Contains **PROlin**

Anticoagulant rodenticide

CAUTION

Keep out of reach of children.

See Right Panel for Additional Cautions

NET WEIGHT LBS.

B & N PRODUCTS CO.

HARLAN, IOWA

ACTIVE INGREDIENTS:

Warfarin 3-(α -acetylbenzyl)-
4-hydroxycoumarin 0.025%

"N"-(2-quinoxaliny) sulfanilamide
(sulfaquinoxaline)." 0.025%

INERT INGREDIENTS 99.950%

CAUTION

Keep out of reach of children

Protect children, pets and domestic animals from bait. If swallowed by humans, domestic animals, or pets this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

This rodenticide contains warfarin, an anticoagulant chemical that reduces the clotting ability of the blood, and, upon repeated feedings, causes fatal hemorrhage in rats. It also contains an antibacterial agent, sulfaquinoxaline, which inhibits bacteria that produce vitamin K, an antidote for warfarin. It thereby increases effectiveness against hard-to-kill rats and gives better and more thorough control.