

(CENTER PANEL)

**DILL**

TURF HERBICIDE MCPP

Especially useful for the control of surface-creeping broadleaf weeds such as clovers, chickweeds, stitchwort, ground ivy and knotweed on sports and ornamental turf, including golf greens, fairways and tees; as well as home lawns and other non-crop turf areas.

ACTIVE INGREDIENT:

Diethanolamine Salt of 2-(2-methyl-4-chloro-  
phenoxy) propionic acid\* . . . . . 32.0%

INERT INGREDIENTS . . . . . 67.2%

\* Equivalent to 22.0% 2-(2-methyl-4-chlorophenoxy) propionic acid or 2.0 lbs. MCPP acid per gallon.

C A U T I O N

Keep Out of Reach of Children

Avoid contact with skin, eyes or clothing. Avoid inhaling spray mist. Keep out of reach of children. Avoid spray drift to susceptible plants as this product may injure tomatoes, beans, peas, ornamentals, etc. (coarse sprays are less likely to drift). Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes (or do not use the same equipment for other purposes). Do not transport or store near seeds, fertilizers, insecticides or fungicides. Do not use in or near greenhouses. Avoid contamination of water intended for irrigation of crops or domestic purposes. Protect from freezing. If allowed to freeze, thaw and re-mix before using. Do not use on St. Augustine Grass.

Do NOT Store with Foodstuffs

Manufactured For

J. J. DILL COMPANY

ACCEPTED  
JUN 9 1967  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND ROENTGENIC ACT  
FOR ECONOMIC POISON REGISTRY  
ED UNDER NO. 100-215-11111  
TO ATTACHED COMPLAINT

GENERAL INFORMATION: MCPP has proven to be a very useful herbicide for control of the surface-creeping or prostrate broadleaf weeds listed below. Although relatively slow in action, one of the advantages of Turf Herbicide MCPP is that it is safer to fine turf grasses, than 2,4-D, silvex or 2,4,5-T. Can be used safely on most bent varieties, but use caution on Seaside, Arlington and Congressional bents which may be injured.

WEEDS CONTROLLED

Common Chickweed	(Stellaria media)
Mouse-ear chickweed	(Cerastium vulgatum)
Red Clover	(Trifolium pratense)
White Clover	(Trifolium repens)
Ground Ivy	(Glechoma hederacea)
Stitchwort	(Stellaria graminea)
Knotweed	(Polygonum aviculare)

DIRECTIONS FOR USE

APPLICATION RATE ON:	PER 1,000 SQ. FT.	PER ACRE
Established greens and fine turf lawns	1½ - 2 fl. ozs.	4 - 5 pints
Other established turf and fairways	2½ - 3 fl. ozs.	6 - 7 pints

Better weed control, especially of knotweed has been obtained with weekly applications at half recommended rate. This is preferred, especially on bent greens and aprons where there has been no previous experience.

USDA REG. NO. \_\_\_\_\_