MIXING INSTRUCTIONS

Mix the oughly one part of FUMARIN[®]-22 CON-CENTRATE with nineteen parts of a suitable bait material such as corn meal, rolled oats, and stock food

A small amount of an edible oil such as mineral, corn or peanut oil may be used up to 5% to prevent dusting and to increase acceptability.

When meat or fish is added to baits, dogs and cats will actively search out the bait placements. Therefore, baits must be placed so as to positively prevent feeding by these animals.

CAUTION

KEEP OUT OF REACH OF CHILDREN

See side panels for additional cautions.



USDA Reg. No. 6900-81

GENERAL DIRECTIONS

Place bait in locations where rats and mice feed, water, or travel.

Bait placements along walls, in corners, and in concealed locations where rats and mice can feed unobserved are desirable.

Bait should be placed in locations not accessible to children, pets, and domestic animals or in tamper-proof bait boxes.

DIRECTIONS FOR USE

FOR RATS: Use in amounts of 1/2 to 1 lb. per bait station.

FOR MICE: Use 1 to 2 oz. per bait station and use more frequent placement (every 8 to 10 feet)

Baits should be examined regularly and replenished before completely eaten.

For effective control baiting should be maintained continuously as long as feeding is observed. Average kill times are from 6 to 14 days but some feeding may continue for 3 to 4 weeks. Where reinfestation is likely, permanent bait stations should be used and bait placed in them as often as necessary.

*FUMARIN® -22 CONCENTRATE

KILLS RATS AND MICE

An Anti-coagulant Concentrate for preparing baits to kill rats and mice.

CONTAINS

ACTIVE INGREDIENT	By Wt.
3-(alpha-acetonylfurfuryl)-4-hydroxycoumarin*	$.5^{\circ}$
INERT INGREDIENTS	99.5°
Total	100 0%

MANUFACTURED FOR THE J. J. DILL COMPANY
1407 RAVINE ROAD, KALAMAZOO, MICHIGAN . P. O. BOX 788

MIXING INSTRUCTIONS

Mix the bugn'y report of FUMARIN⁸-22 CON-CENTRATE with mineteen parts of a suitable balt material such as corn meal, rolled cuts, and stack food

A small amount of an edible oil such as mineral, corn or peanut oil may be used up to 5% to prevent dusting and to increase acceptability.

When meat or fish is added to baits, dogs and cats will actively search out the bait placements. Therefore, baits must be placed so as to positively prevent feeding by these animals.

GENERAL DIRECTIONS

Place bait in locations where rats and mice feed, water, or travel.

Bait placements along walls, in corners, and in concealed locations where rats and mice can feed unobserved are desirable.

Bait should be placed in locations not accessible to children, pets, and domestic animals or in tamper-proof bait boxes.

DIRECTIONS FOR USE

FOR RATS: Use in amounts of 1 2 to 1 lb. per bait station.

FOR MICE: Use 1 to 2 oz. per bait station and use more frequent placement (every 8 to 10 feet)

Baits should be examined regularly and replenished before completely eaten.

For effective control baiting should be maintained continuously as long as feeding is observed. Average kill times are from 6 to 14 days but some feeding may continue for 3 to 4 weeks. Where reinfestation is likely, permanent bait stations should be used and bait placed in them as often as necessary.

CAUTION

KEEP OUT OF REACH OF CHILDREN

See side panels for additional cautions.



USDA Reg. No. 6900-81

*FUMARIN® -22 CONCENTRATE

KILLS RATS AND MICE

An Anti-coagulant Concentrate for preparing baits to kill rate and mice

CONTAINS

ACTIVE INGREDIENT		.5, 41
3 alpha acetonyifurfuryi 4 hydroxy comarini		<u> </u>
INERT INGREDIENTS		.0 5
	Tota.	100 0

MANUFACTURED FOR THE J. J. DILL COMPANY
1407 RAVINE ROAD, KALAMAZOO, MICHIGAN . DO BOX 788

CAUTION

KEEP OUT OF REACH OF CHILDREN

See side panels for additional cautions.



USDA Reg. No. 6900-81

In case baits are eaten, give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is alear. Call a physician immediately

Baits that become sour, moldy, spoiled, or contaminated in any way should be replaced

FUMARIN®-22 CONCENTRATE contains as its

active ingredient an anti-coagulant chemical.

Therefore, FUMARIN = 22 must be protected from

or placed where humans, livestock, poultry, or

NOTE FOR PHYSICIANS:

pets will not eat them.

Intravenous injections of Viramin K1 is indicated together with administration of blood transfusions.

LOT NO.

IMPORTANT

promptly.

CAUTION

NOTICE

THE J. J. DILL COMPANY

MANUFACTURERS AND DISTRIBUTORS

OFFICE AND WAREHOUSE -

1407 RAVINE ROAD • (616) 349-7755

MAIL ADDRESS -

P.O. BOX 788, KALAMAZOO, MICH. 49005

*FUMARIN® -22 CONCENTRATE

KILLS RATS AND MICE

An Anti-coagulant Concentrate for preparing baits to kill rats and mice.

CONTAINS

ACTIVE INGRED ET. I		By W
3 ampha acetony (forfor, 1997) to the speciment		i.
INERT INGREDIENTS		49.5
	Tatal	100.0

MANUFACTURED FOR THE J. J. DILL COMPANY
1407 RAVINE ROAD, KALAMAZOO, MICHIGAN . . . O. BOX 788