


MIXING INSTRUCTIONS—DRY BAITS

Mix thoroughly 1 part DIPHACINONE-WS CONCENTRATE with 19 parts of a suitable bait material such as corn meal, rolled oats, and stock food.

A small amount of an edible oil such as mineral, corn, or peanut oil may be used up to 5% to prevent dusting and to increase acceptability.


When meat or fish is added to baits, dogs and cats will actively search out the bait placements. Therefore, baits must be placed so as to positively prevent feeding by these animals.

by these animals.



With Cornmeal

DILL



EPA Reg. No. 6900-84

For ef
tinuou
times
contin
perma
in the

DIPH/
active
DIPH/
tected
try, p

GENERAL DIRECTIONS

MIXING INSTRUCTIONS—LIQUID BAITS

Dissolve 1 1/2 oz. (by wt.) of DIPHACINONE WS CONCENTRATE in 1 quart of water. Expose in water fountains in protected stations.

GENERAL DIRECTIONS

Place bait in locations where rats and mice feed, water, or travel.

Bait placements along walls, in corners and in concealed locations where rats and mice can feed unobserved are desirable.

Bait should be placed in locations not accessible to children, pets, domestic animals, and wildlife, or in tamper proof boxes.

DIRECTIONS FOR USE—LIQUID BAITS

For Rats: Pint liquid dispenser per location should be the minimum amount placed. An uninterrupted supply of bait should be maintained for at least 10 days.

For Mice: Pint liquid dispenser should be placed at 8 to 10 foot intervals. An uninterrupted supply of bait should be maintained for at least 15 days.

DIRECTIONS FOR USE—DRY BAITS

For Rats: Use in amounts of 1/2 to 1 lb. per bait station. An uninterrupted supply of bait should be maintained for at least 15 days.

For Mice: Use 1 to 2 oz. per bait station and use more frequent placement (every 8 to 10 feet). An uninterrupted supply of bait should be maintained for at least 15 days.

in them as often as necessary.

UNIVERSAL CONCENTRATE — USE WET OR DRY

DIPHACINONE WS CONCENTRATE

KILLS RATS AND MICE

An Anti-coagulant Concentrate for preparing baits to kill rats and mice.

CONTAINS

ACTIVE INGREDIENTS	By Wt.
Sodium Salt of Diphacinone (2 Diphenylacetyl-1, 3-indandione)	0.106%
INERT INGREDIENTS (sugar)	99.894%
Total	100.000%

MANUFACTURED FOR THE J. J. DILL COMPANY
1407 RAVINE ROAD, KALAMAZOO, MICHIGAN . P O BOX 788

In cas
glass
clear.

Intrav
togeth

Baits
ated i
reuse
crushi

LOT

T H
OFFI

MAIL

NET


MIXING INSTRUCTIONS—DRY BAITS

Mix thoroughly 1 part DIPHACINONE WS CONCENTRATE with 19 parts of a suitable bait material such as corn meal, rolled oats, and stock food.

A small amount of an edible oil such as mineral, corn, or peanut oil may be used up to 5% to prevent dusting and to increase acceptability.

When meat or fish is added to baits, dogs and cats will actively search out the bait placements. Therefore, baits must be placed so as to positively prevent feeding by these animals.

by these animals.



EPA Reg. No. 6900-84

GENERAL DIRECTIONS

MIXING INSTRUCTIONS—LIQUID BAITS

Dissolve 1 oz. (by wt.) of DIPHACINONE WS CONCENTRATE in 1 quart of water. Expose in water troughs in protected stations.

GENERAL DIRECTIONS

Place bait in locations where rats and mice feed, water or travel.

Bait placements along walls, in corners and in concealed locations where rats and mice can feed undisturbed are desirable.

Bait should be placed in locations not accessible to children, pets, domestic animals, and wildlife, or in tamper proof boxes.

DIRECTIONS FOR USE—LIQUID BAITS

For Rats: Pint liquid dispenser per location should be the minimum amount placed. An uninterrupted supply of bait should be maintained for at least 10 days.

For Mice: Pint liquid dispenser should be placed at 8 to 10 foot intervals. An uninterrupted supply of bait should be maintained for at least 15 days.

DIRECTIONS FOR USE—DRY BAITS

For Rats: Use in amounts of 1/2 to 1 lb. per bait station. An uninterrupted supply of bait should be maintained for at least 15 days.

For Mice: Use 1 to 2 oz. per bait station and use more frequent placement (every 8 to 10 feet). An uninterrupted supply of bait should be maintained for at least 15 days.

in them as often as necessary

UNIVERSAL CONCENTRATE — USE WET OR DRY

DIPHACINONE WS CONCENTRATE

KILLS RATS AND MICE

An Anti-coagulant Concentrate for preparing baits to kill rats and mice.

CONTAINS

ACTIVE INGREDIENTS	3.7%
Sodium Salt of Diphacinone (2 Diphenylacetyl-1,3-indandione)	0.106
INERT INGREDIENTS (sugar)	99.894
Total	100.000

MANUFACTURED FOR THE J. J. DILL COMPANY
1107 RAVINE ROAD, KALAMAZOO, MICHIGAN 49001, P.O. BOX 788

For eff
tinuous
times
contin
permal
in ther

DIPHA
active
DIPHA
tected
try, pe

In case
glass o
clear. C

Intrave
togethe

Baits t
ated in
reuse
crushin



LOT I

T H
M
OFFIC

MAIL

NET V

BAITS
CONCENTR-
terial such
heral, corn,
ent dusting
nd cats will
Therefore,
ent feeding



EPA Reg. No. 6900-84

For effective control baiting should be maintained continuously as long as feeding is observed. Average kill times are from 6 to 14 days but some feeding may continue for 3 or 4 weeks. Where reinfestation is likely, permanent bait stations should be used and bait placed in them as often as necessary.

CAUTION

DIPHACINONE-WS CONCENTRATE contains as its active ingredient an anti-coagulant chemical. Therefore, DIPHACINONE-WS CONCENTRATE must be protected from or placed where humans, livestock, poultry, pets, or wildlife will not eat them.

BAITS
E WS CON
e in water
eed, water
nd in con
feed unob
ccessible to
dlife, or in
BAITS
h should be
ted supply
t 10 days
placed at 8
ply of bait
ITS
or bait sta
d be main
on and use
l feet). An
ntained for

UNIVERSAL CONCENTRATE -- USE WET OR DRY

DIPHACINONE WS CONCENTRATE

KILLS RATS AND MICE

An Anti coagulant Concentrate for preparing baits to kill rats and mice.

CONTAINS

ACTIVE INGREDIENTS	By Wt.
Sodium Salt of Diphacinone (2 Diphenylacetyl 1, 3 indandione)	0.106
INERT INGREDIENTS (sugar)	99.894
Total	100.000

MANUFACTURED FOR THE J. J. DILL COMPANY
1407 RAVINE ROAD, KALAMAZOO, MICHIGAN P. O. BOX 788

NOTICE

In case baits are eaten, give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately.

NOTE FOR PHYSICIANS

Intravenous injections of Vitamin K1 is indicated together with administration of blood transfusions.

IMPORTANT

Baits that become sour, moldy, spoiled, or contaminated in any way should be replaced promptly. Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

LOT No.

THE J. J. DILL COMPANY
MANUFACTURERS AND DISTRIBUTORS
OFFICE AND WAREHOUSE
1407 RAVINE RD (616) 349-7755
MAIL ADDRESS
P.O. BOX 788, KALAMAZOO, MICH 49005

NET WEIGHT LBS.