
DIRECTIONS FOR USE

D-Lete is a nonselective herbicide that can be applied before or after plant growth begins. Although D-Lete has considerable activity through foliar contact, much of its activity is through roots, therefore, its effectiveness is dependent on rainfall to move it into the root zone. Very dry soil conditions and lack of sufficient rainfall may result in poor weed control. D-Lete can be applied prior to or up to 3 months after weed emergence. Application rates vary from 21 to 125 gallons of D-Lete per acre, depending on climatic conditions, soil type, the weeds present and the stage of growth of the weeds.



CAUTION

KEEP OUT OF THE REACH OF CHILDREN

See side panels for additional cautions.

EPA Reg. No. 6900-134

In the following rate recommendations, the higher rates are intended for use on heavier soils, where weed growth is heavy, and where rainfall can be expected to be relatively high. The higher rates should also be used where longer residual control is desired in regions with a long growing season.

For best results apply D-Lete at rates recommended below prior to weed emergence or when weeds are young and actively growing.

For control of annual and susceptible perennial weeds (such as downy bromegrass, oatgrass, goosegrass, quackgrass, puncturevine, goldenrod, and plantain): Broadcast 21 to 31 gallons D-Lete in 50-100 gallons of water per acre. For small areas, apply 4 1/4-6 1/4 pints D-Lete per 1,000 square feet in sufficient water to give thorough and uniform coverage.

For control of hard-to-kill perennial weeds and grasses (such as Johnson grass, bindweed, and wild carrot): Broadcast 84-125 gallons D-Lete in 50-100 gallons of water per acre. For small areas, use 2-3 gallons D-Lete per 1,000 square feet in sufficient water to give thorough and uniform coverage.

For faster top-kill of existing vegetation, apply D-Lete in diesel oil, fuel oil, or weed oil. Use oils at rates of 100-200 gallons per acre. Tall, dense vegetation such as Johnsongrass will generally require 150 or more gallons of oil per acre for thorough coverage.

D-LETE LIQUID

A LIQUID CONCENTRATE CONTAINING PRAMITOL* FOR KILLING WEEDS AND GRASSES WHERE LONG LASTING RESIDUAL EFFECT IS DESIRED.

CONTAINS


ACTIVE INGREDIENTS		By Wt.
2,4-bis (isopropylamino)-6-methoxy-s-triazine		6.0%
INERT INGREDIENTS		94.0%
	TOTAL	100.0%

D-Lete contains .472 lbs. Pramitol* per gallon. *Reg. Trade Mark CIBA-Geigy

MANUFACTURED BY THE J. J. DILL COMPANY
1407 RAVINE ROAD, KALAMAZOO, MICHIGAN . P. O. BOX 788

DIRECTIONS FOR USE

D-Lete is a nonselective herbicide that can be applied before or after plant growth begins. Although D-Lete has considerable activity through foliar contact, much of its activity is through roots, therefore, its effectiveness is dependent on rainfall to move it into the root zone. Very dry soil conditions and lack of sufficient rainfall may result in poor weed control. D-Lete can be applied prior to or up to 3 months after weed emergence. Application rates vary from 21 to 125 gallons of D-Lete per acre, depending on climatic conditions, soil type, the weeds present and the stage of growth of the weeds.



In the following rate recommendations, the higher rates are intended for use on heavier soils, where weed growth is heavy, and where rainfall can be expected to be relatively high. The higher rates should also be used where longer residual control is desired in regions with a long growing season.

For best results apply D-Lete at rates recommended below prior to weed emergence or when weeds are young and actively growing.

For control of annual and susceptible perennial weeds (such as downy bromegrass, oatgrass, goosegrass, quackgrass, puncturevine, goldenrod, and plantain): Broadcast 21 to 31 gallons D-Lete in 50-100 gallons of water per acre. For small areas, apply 4 1/4-6 1/4 pints D-Lete per 1,000 square feet in sufficient water to give thorough and uniform coverage.

For control of hard-to-kill perennial weeds and grasses (such as Johnson grass, bindweed, and wild carrot): Broadcast 84-125 gallons D-Lete in 50-100 gallons of water per acre. For small areas, use 2-3 gallons D-Lete per 1,000 square feet in sufficient water to give thorough and uniform coverage.

For faster top-kill of existing vegetation, apply D-Lete in diesel oil, fuel oil, or weed oil. Use oils at rates of 100-200 gallons per acre. Tall, dense vegetation such as Johnsongrass will generally require 150 or more gallons of oil per acre for thorough coverage.

D-LETE® LIQUID


A LIQUID CONCENTRATE CONTAINING PRAMITOL* FOR KILLING WEEDS AND GRASSES WHERE LONG LASTING RESIDUAL EFFECT IS DESIRED.

CONTAINS

ACTIVE INGREDIENTS	By Wt.
2,4-bis (isopropylamino)-6-methoxy-s-triazine	6.0%
INERT INGREDIENTS	
TOTAL	100.0%

D-Lete contains .472 lbs. Pramitol* per gallon. *Reg. Trade Mark C1BA-Geigy

MANUFACTURED BY THE J. J. DILL COMPANY
1407 RAVINE ROAD, KALAMAZOO, MICHIGAN . P. O. BOX 788



CAUTION

KEEP OUT OF THE REACH OF CHILDREN

See side panels for additional cautions.

EPA Reg. No. 6900-134

IMPORTANT

Use only in areas where complete control of all vegetation is desired, such as industrial sites, rights-of-way, lumberyards, petroleum tank farms, around farm buildings, along fence lines, etc. When applied to the soil, this product usually inhibits plant growth for a year or more. It should not be used on land to be cropped, or near adjacent desirable trees, shrubs, or plants, or injury may occur.

It is noncorrosive to equipment.

D-LETE LIQUID

A LIQUID CONCENTRATE CONTAINING PRAMITOL* FOR KILLING WEEDS AND GRASSES WHERE LONG LASTING RESIDUAL EFFECT IS DESIRED.

CONTAINS

ACTIVE INGREDIENTS		By Wt.
2,4-bis (isopropylamino) 6-methoxy-s-triazine		6.0%
INERT INGREDIENTS		94.0%
	TOTAL	100.0%

D-Lete contains .472 lbs. Pramitol* per gallon.

*Reg. Trade Mark CIBA-Geigy

MANUFACTURED BY THE J. J. DILL COMPANY
1407 RAVINE ROAD, KALAMAZOO, MICHIGAN . P. O. BOX 788

CAUTION

Harmful if swallowed. Avoid contact with eyes and skin. In case of skin contact, wash with soap and water. Avoid inhalation of spray mist. Avoid contamination of food and feed.

CAUTION

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place. Return empty drum to drum reconditioner.

Do not use, pour, spill or store near heat or open flame. Store at temperatures above 32° Fahrenheit.

LOT No.

THE J. J. DILL COMPANY

MANUFACTURERS AND DISTRIBUTORS

OFFICE AND WAREHOUSE -

1407 RAVINE RD. (616) 349-7755

MAIL ADDRESS -

P.O. BOX 788, KALAMAZOO, MICH. 49005

NET CONTENTS

GALLONS