[6902-7]

ingredient statement: This formulation is an effective insecticide aerosol designed to be used in automatic dispensing units in dairy barns, horse barns, livestock quarters, veterinary clinics, feed stores, hospitals and other areas where flies, mosquitoes, and gnats are an annoyance. Do not use in food service areas or food preparation areas.

USE ONLY WHEN INSECT INFESTATION EXISTS.

CAUTION: Do not use in nurseries or rooms where infants, ill or aged persons are confined. Large drafts of forced ventilation may reduce or eliminate the insecticidal officery of these installations. Do not install forced y over any food handling or dispensing area.

warning: Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130° F. may cause bursting. Never throw container into fire or inuncharator.

FUNGION AND A TOP OF TOP OF THE PORT OF TH

EPA Reg. No. 6902-7



VETSECT

METERED INSECTICIDE For Time Release

ACTIVE INGREDIENTS:

Pyrethrin	Š			0.9%
Technical	Piper	onyl	Butoxide*	9.0%
Petroleun				
Equivalen	t to	7.2%	(butylcarbityl)	(6-propyl-
piperonyl	ether	and 1	.8% of related c	ompounds.
NERT ING	REDIE	NTS:	***************************************	80.00 %

100.00%

CAUTION: KEEP FROM REACH OF CHILDREN Contents under pressure. See other precautions on the side panel.

SALES TO VETERINARIANS ONLY

MED WITHHI 12 OUNCES

Trade Mark

Mfg. For



OSBORN LABORATORIES, INC INTERNATIONAL MULTIFOODS CORPORATION LE SUEUR MINNESOTA 56068

DIRECTIONS FOR USE

FOR USE IN AUTOMATIC AEROSOL DISPENSER: One unit should be installed for each 6,000 cubic feet of space to be treated. The Dispenser activates the 100 mg. valve once every 15 minutes; after two hours of operation, enough insecticide is suspended in the air to kill a substantial proportion of the flies, mosquitoes and gnats present. Usually the effectiveness of the insecticide will be noticeable within one hour after the dispenser is activated. If less than 24 hour control is required, the dispenser should be set to start one hour before control is desired and run one hour after control is no longer needed. Keep in mind that such factors as air flow rate, size of space treated, humidity and room temperature all affect insect activity and control. Therefore. one unit per 6,000 cubic feet should be considered as an average need with adjustments made depending on specific conditions at place of usage. As with any method of insect control best results are achieved when sanitation and other positive insectprevention practices are followed. This 12 oz. can of insecticide contains approximately 3,300 100 mg. doses, sufficient to sustain operation for a total of 825 hours (34 days on around the clock basis) or 68 days on a 12-hour basis. When used on a 24-hour basis, it is suggested that the container be changed once a month (once every 2 months on a 12-hour basis.)

Aug. 24-1923 6901-7

RECEIVED AUG 10 973