

Sanitizing of Food Processing Equipment and other Hard Surfaces in Food Contact Locations.

For sanitizing food processing equipment, dairy equipment, food utensils, dishes, silverware, glasses, sink tops, countertops, refrigerated storage and display equipment and other hard non-porous surfaces. No potable water rinse is required.

Wash and rinse all articles thoroughly, then apply a solution of 1 oz. SANTEC in 2 gallons of water. (150 ppm active). Surfaces should remain wet for at least one minute followed by adequate draining and air drying. Fresh solution should be prepared daily or when use solution becomes visibly dirty. For mechanical applications, use solution may not be reused for sanitizing applications.

Apply to sink tops, countertops, refrigerated storage and display equipment and other stationary hard surfaces by cloth or brush. No potable water rinse is required.

Dishes, silverware, glasses, cooking utensils and other similar sized food processing equipment can be sanitized by immersion in a 1 oz./2 gallon dilution of SANTEC. No potable water rinse is required.

Precautionary Statements

Hazards to Humans and domestic animals

DANGER

Keep out of reach of children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

STORAGE AND DISPOSAL

- Do not contaminate water, food, or feed by storage disposal.
- Do not store on side.
- Avoid creasing or impacting of side walls.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray or mixture of rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Triple Rinse (or equivalent): Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities by burning. If burned, stay out of smoke.

Metal Container: Triple rinse (or equivalent), then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. (If container is 1 gallon or less, use this container disposal statement.)

Do not reuse empty container (bottle, can, bucket). Wrap container and put in trash.

ACCEPTED
JUN 12 1985
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 6921-8

SANTEC

**DISINFECTANT-SANITIZER
FUNGICIDE DEODORIZER**

Disinfectant-Sanitizer-Fungicide Deodorizer for Hospital, Institutional, Industrial & School Use

Active Ingredient

Didecyl dimethyl ammonium chloride 3.75%

Inert Ingredients 96.25%

100.00%

KEEP OUT OF REACH OF CHILDREN

DANGER

Statement of Practical Treatment

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

SEE LEFT PANEL FOR

ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Registration No. 6921-8

EPA Establishment No. 6921-W11

Manufactured By:

TESCH CHEMICAL CO., INC.

E. Midway Road, Appleton, WI 54912

FOR SALE FOR USE AND STORAGE BY
MAINTENANCE PERSONNEL ONLY

**DIRECTIONS FOR USE
GENERAL CLASSIFICATION**

It is a violation of Federal Law to use this product in a manner inconsistent with its labelling.

Apply SANTEC with a cloth, mop, or mechanical spray device; surface must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily or when the use solution becomes visibly dirty.

Disinfection in Hospitals, Nursing Homes and Other Health Care Institutions.

For disinfecting floors, walls, countertops, bathing areas, lavatories, bedframes, tables, chairs, garbage pails and other hard non-porous surfaces.

Add 2 oz. SANTEC to 1 gallon water. Apply to previously cleaned hard surface with mop or cloth.

At this use-level, SANTEC is effective against *Pseudomonas aeruginosa*.

Disinfectant in Institutions, Industry, and Schools.

For disinfecting floors, walls, bedframes, countertops, tables, chairs, garbage pails, bathroom fixtures and other hard non-porous surfaces.

Add 1 oz. of SANTEC to 1 gallon of water. Apply to previously cleaned hard surface with mop or cloth.

At 1 oz./gallon use-level, SANTEC is effective against *Staphylococcus aureus*, *Salmonella choleraesuis* and *Trichophyton interdigitale* (the athlete's foot fungus).

Net Contents

BEST AVAILABLE COPY