

IONS

lon water.

ons water.

Use a test
dosage as
n required
hlorine.

ght. Keep
s light and
length.

um hypo-
n 6 months
anufacture.

CHLORICIDE-12

Formula S-12

Active Ingredient:

Sodium Hypochlorite	10.5%
Inert Ingredient	89.5%
TOTAL	100.00%

NET CONTENTS WARNING—Keep out of the reach of children

ONE U. S. GALLON See right panel for additional precautions

ANTIDOTE EXTERNAL

Wash with Water
INTERNAL

Drink large quantities of mu-
cilage, raw egg white, milk or
rice gruel.

CAUTION

Do not mix Chloricide-12 with
acid cleaners.

Harmful if swallowed. Contact
may cause severe irritation
and eye damage. Avoid con-
tact with skin and eyes.

Note To Physician: Probable mu-
cosal damage may contraindicate
the use of gastric lavage.

Printed in U.S.A.

DE-12

**ANTIDOTE
EXTERNAL**

**Wash with Water
INTERNAL**

Drink large quantities of mucilage, raw egg white, milk or rice gruel.

CAUTION

Do not mix Chloricide-12 with acid cleaners.

Harmful if swallowed. Contact may cause severe irritation and eye damage. Avoid contact with skin and eyes.

Note To Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

Printed in U.S.A.

**For Food Processing And
Milk Equipment**

Wash all surfaces thoroughly, rinse with potable water and sanitize as follows: Dairy Equipment — Sanitize with 200 PPM available Chlorine. Apply with a Merit San Spray, Lonn Sprayer or brush. Tables, Meat Blocks, Walls, Floors — Sanitize with 800 PPM by brush, mop or San Spray. Thoroughly wet all surfaces. Rinse with a potable water.

Note: All food contact surfaces, treated with concentration in excess of 200 PPM available chlorine, must be rinsed with potable water before rinse.


Local, State Regulations

Where local or state regulations are in effect, consult authorities for proper sanitizing concentrations and procedures. Rinse empty container thoroughly with water and discard it.

EPA Registration No. 6931-8

..... 10.5%
..... 89.5%
..... 100.00%

**of the reach of children
r additional precautions**



**CHEMICAL
TOOLS
FOR
INDUSTRY™**

TOWERFUL

**Multi-Function
Cooling Water Treatment**

APPLICATION

TOWERFUL is used to control algae, bacteria, and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algae growth, microbiological slime, and other deposits. An initial slug addition of 4.0 to 10.0 fluid ounces of TOWERFUL per 1000 gallons of water to provide a concentration of 22 to 80 parts per million of TOWERFUL based on the total amount of water in the system is recommended. Repeat at the dosage until control is evident; subsequent slug additions of 1.0 to 10.0 fluid ounces of TOWERFUL per 1000 gallons of water (8 to 80 parts per million of TOWERFUL) should be considered every 2 to 6 weeks as needed. For the long-term control of algae, bacteria, and fungi, the dosage should be maintained at 2 to 6 parts per million of TOWERFUL. For the control of microbiological slime, the dosage should be maintained at 2 to 6 parts per million of TOWERFUL.

ACTIVE INGREDIENTS 15.0

Poly(oxyethylene(dimethylamino)ethylene(dimethylamino)ethylene dichloride)


INERT INGREDIENTS 85.0

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Do not swallow. Avoid contact with eyes. If contact occurs, flush thoroughly with water. If swallowed, drink plenty of water. If inhaled, get fresh air. If on skin, wash with soap and water.

For product information, contact the manufacturer. Do not use if the container is damaged. Do not use if the product is discolored or contains foreign matter. Do not use if the product is past its expiration date. Do not use if the product is not clearly labeled. Do not use if the product is not clearly labeled.

net contents U.S. gallons



- ALGAECIDE
- FUNGICIDE
- BACTERICIDE
- ANTIFOULANT

- KILLS AND CONTROLS ALGAE
- MINIMIZES MAINTENANCE PROBLEMS
- CONTAINS NO FREE ACIDS OR ALKALIS

E.P.A. REG. NO. 6962-29-AA

MADE IN U.S.A.

