

[6959-37]



INGREDIENTS

Active ingredient: ...
Inert ingredients: ...

TECHNICAL DATA-FLY CONTROL


Application: ...
Treatment per: ...
Cost of treatment: ...
Delivery rate: ...
Cylinder: ...
Excellent Gas: ...
Coverage: ...

CAUTION

Hold dispenser three feet away from object being sprayed to avoid possible damage.

Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when plant is not in operation. Do not remain in treated areas. Ventilate the area after treatment is completed. This cylinder is not refillable. Do not attempt to recharge the empty cylinder. When cylinder is exhausted, discard in safe place.

Federal law forbids transportation if refilled —penalty up to \$10,000 fine and 10 years imprisonment (18 U. S. C. 831-835).



To KILL flying insects: face flies, house flies, stable flies, horn flies, gnats, fruit flies.

To KILL crawling insects, wasps, roaches, spiders, waterbugs, ants, silverfish.

To KILL fleas

CAUTION:

CONTENTS UNDER PRESSURE
Keep out of reach of children. See side panel for additional cautions.

NET WEIGHT 40 OZS. (2 1/2 lbs.)

Part 39NRC M.I. 11
Service Test Pressure
P.E.A. 30.0 N. 10.0
M.I. 11

DIRECTIONS FOR

MILKING PARLOR AND MILK ROOM: Close all windows and doors. Spray at the rate of 1/2 ounce per 100 square feet. If rooms cannot be closed tight, double the dosage. Moving from the furthest point from the exit toward the exit, direct aerosol upward and spray for 10 to 15 seconds. May be counted off by 10 seconds and two thousand (2,000) cycles. Repeat 1/2 hour, 5 minutes after treatment. Example: A room 20 feet long, 12 feet wide at 10 feet high requires 2 seconds of spray at 4 cycles if not. Area treatment is to be made when animals are in the building and in milking pens or holding pens. For wasps, roaches, spiders, etc., spray insects in their hiding and nesting places. For roaches, spiders, sinks, cabinets, etc., spray for 10 to 15 seconds. When necessary, repeat treatment.

FLEAS: Close all windows and doors. Hold dispenser 3 feet above floor and direct spray toward floor and lower walls at a rate of 1/2 ounce per 100 square feet, making sure that all floor and wall surfaces are sprayed. Also treat the sleeping quarters of patients and animals. Avoid keeping patients in the area for at least 12 hours from the time of treatment. For best results, quarters should be treated with insecticide immediately after treatment. After 2 days, if more than 10 fleas are seen, repeat treatment.

WARNING

Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 140°F may cause contents to vent through container into the neighborhood.

Manufactured by GESSCO, INC., Box 12452, Charlotte, N. C. 28205