

PRECAUTIONARY STATEMENTS:

HAZARD TO HUMANS & DOMESTIC ANIMALS:

CAUTION: Keep away from humans, domestic animals and pets. If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of Vitamin K, combined with blood transfusion are indicated, as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

ENVIRONMENTAL HAZARD: Keep out of lakes, streams or ponds.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS: For control of Norway rats, Roof rats and House mice in and around homes, industrial and agricultural buildings, and similar man-made structures. Treated baits must be placed in tamper proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS: Determine areas where rats are most likely to be present and place the bait. Generally, these areas are along walls, around openings in or beside masonry, in corners, and in creviced places between floors and walls, etc., where rodents or their signs have been observed. Exclude bait from animal food.

Remove as much food as possible.

BAIT-TOX®

Ready-Mixed RAT BAIT (Meal Form)

Kills Norway Rats, Roof Rats and House Mice.

ACTIVE INGREDIENT(S):

2-Pivalyl-1, 3-Indandione 0.025%

INERT INGREDIENTS: 99.975%

TOTAL: 100.000%

CAUTION KEEP OUT OF REACH OF CHILDREN

See side panel for additional
precautionary statements.

manufactured by



1109 Central Ave.
Charlotte, NC 28204

EPA REG. NO. 1-6942
EPA EST. NO. 1-59 NC1

NET CONTENTS: 1 lb.

APPLICATION DIRECTIONS

RATS: Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Apply 1/4 to 1/2 ounce (1-2 level tablespoons) of bait at 8 to 12 foot intervals. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS & MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

DISPOSAL

DO NOT REUSE EMPTY CONTAINER.

WRAP CONTAINER AND PUT IN TRASH COLLECTION

ACCEPTED

SEP 29 1988

6959-42