

Precautionary Statement - Hazard to Humans & Domestic Animals
CAUTION

Harmful if swallowed. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes, or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Avoid contamination of food, utensils, and food preparation areas.

Statement of Practical Treatment

If swallowed: Call a physician or Poison Control Center. Do not induce vomiting.
If inhaled: Remove victim to fresh air. Apply artificial respiration if indicated.
If on skin: Remove contaminated clothing; wash skin with soap and water. Get medical attention if irritation persists.
If in eyes: Flush with plenty of water. Get medical attention if irritation persists.

Environmental Hazards

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning or disposal of equipment.

Physical or Chemical Hazards

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperature above 130°F may cause bursting. Do not spray on plastic, painted, or varnished surfaces or directly into electronic equipment such as radios, televisions, computers, etc.

Applications in Food Areas of Food Handling Establishments

Limit to Crack and Crevice Treatment only- apply a small amount of material directly into cracks and crevices such as expansion joints between different elements of construction or between equipment bases and the floor, wall voids, motor housing, junction boxes, conduits, or hollow equipment legs where insects hide. Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces. Application of this product in the food areas of food handling establishments, other than as a crack and crevice treatment, are not permitted.

Storage and Disposal

Storage: Do not store near heat or open flame.
Disposal: Do not reuse empty container. Replace cap, wrap and discard in trash collection.

Important note: Buyer assumes all risks of use, storage, or handling of this product not in strict accordance with directions given hereon.

C H T 1 2 7



1D RESIDUAL INSECTICIDE

CRACK AND CREVICE USE ONLY

ACTIVE INGREDIENTS:
Chlorpyrifos (o, O-diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate).....1.00%
****INERT INGREDIENTS.....99.00%**
100.00%

**Contains Petroleum Distillates

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

AEROSOL INSECTICIDE TO KILL:

Cockroaches, Waterbugs, Spiders, Ticks, Earwigs, Crickets, Sowbugs, Millipedes, Centipedes, Booklice, silverfish, Ants, Grain Weevils, Flour Beetles, Chocolate Moths, Drug Store Beetles, Boxelder Bugs, Elmleaf Beetles, Clover Mites, Cluster Flies, Carpenter Ants, Carpenter Beetles, and Termites.

ESPECIALLY FORMULATED FOR USE IN:

Hospitals, supermarkets, motels, hotels, homes, transportation equipment (buses, boats, ships, trains, planes), utilities, food processing plants, food storage areas, and restaurants and other food handling establishments.

NET WEIGHT : 16.5 oz. (468 grams)

EPA Reg. No 6959-73
EPA Est No 6959-SC-1

MANUFACTURED BY

CESSCO INC.

6695 A RIVER RD
JOHN'S ISLAND, SC 29455

1-800-476-3666

ACCEPTED
AUG 11 1994
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 6959-73

Directions For Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CESSCO'S 1D ACCUDOSE Insecticide is for use only with injection tube and for injecting into cracks and crevices where insects may be hiding, living and breeding. Place injector tip 1/2" into cracks, crevices, holes, and other small openings where insects may be a problem. Release approx. 1 second spray per shot. In serious infestation, if possible spots approx. 12" apart. In light infestation or preventative treatment place spots approx. 3 feet apart. See the additional directions below for effective control of specified insects.

CRAWLING INSECTS: COCKROACHES, WATER-BUGS, SPIDERS, TICKS, EARWIGS, CRICKETS, SOWBUGS, MILLIPEDES, CENTIPEDES. Spray into hiding places such as cracks and crevices, moist areas, openings around pipes and sinks, under refrigerators, behind baseboards, coffee makers, meter boxes and manholes. Spray directly on insects in these locations where possible. Repeat as necessary once every seven days.

BOOKLOUSE & SILVERFISH: Spray cracks and crevices in baseboards, door frames, bookcase and storage areas. Spray directly on insects in these location where possible. Repeat as necessary once every seven days.

ANTS: Spray into trails, nests, and points of entry around doors, windows, structural cracks and crevices. Spray directly on insect in these places if possible. Repeat as necessary once every seven days.

GRAIN WEEVILS, FLOUR BEETLES, CHOCOLATE MOTHS, DRUG STORE BEETLES (in exposed adult and larvae stages) and hibernating stages of: **BOXELDER BUGS, ELMLEAF BEETLES, CLUSTER FLIES and CLOVER MITES.** Contact as many as possible when treating infested areas such as cracks and crevices, in wall voids, around windows and door frames, under siding, in machinery and in dark secluded areas. Repeat as necessary once every seven days.

Localized treatment of wood-destroying insects: **CARPENTER ANTS, CARPENTER BEES and TERMITES.** For control of carpenter ants, locate source of infestation. Inject a 5- to 10 second spray through openings or small newly drilled holes into wall voids where these ants or their nests are present. For control of carpenter bees, locate source of infestation. Inject a 5- to 10-second spray into bee tunnels and cavities. Seal entrance openings with small pieces of stainless steel wool after spray application. For control of termites, inject a 5- to 10-second spray into voids and channels in damaged wood and in spaces between wooden chambers of a structure. Application may be made to inaccessible areas by drilling then injecting the spray. Treatment of localized areas is intended to kill workers and winged reproductive forms of termites in the treated areas and to prevent infestations for a temporary period. This type of application is not intended to be a substitute for soil treatment or mechanical alteration to control subterranean termites.

BEST AVAILABLE COPY