

JUN 09 1994

Reg # 6959-87

PM-13

1/2/1

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 6959-97

**CESSCO III P**  
INSECT KILLER

- \*CONTAINS PYRETHROID AND PERMANENT
- \*SEEKS OUT BUGS WHERE THEY HIDE
- \*KILLS ON CONTACT
- \*CONTROLS INDUSTRIAL INSECTS
- \*FOR COMMERCIAL/AGRICULTURAL USE
- \*A BROAD SPECTRUM INSECTICIDE HIGHLY EFFECTIVE AGAINST ANTS, (INCLUDING CARPENTER ANTS), BEES, BEETLES, (INCLUDING CARPET, FLOUR, GRAIN, AND GROUND BEETLES), CENTIPEDES, CLOVER MITER, COCKROACHES, CRICKETS (INCLUDING MOLE CRICKETS), EARWIGS, FIREBRATS, FLEAS, MILLIPEDES, MOTHS, PILLBUGS, SCORPIONS, SILVERFISH, SOWBUGS, SPIDERS, TICKS, WASPS, AND OTHER PESTS LISTED ON THE LABEL.
- \*KILLS LICE AND THEIR EGGS
- \*FOR USE ON BEDDING, FURNITURE AND OTHER INANIMATE OBJECTS INFESTED WITH LICE
- \*NON-STAINING
- \*NO UNPLEASANT ODOR
- \*WATER-BASED FORMULA
- \*FOR USE ON DOGS AND ON PREMISES TO CONTROL FLEAS AND TICKS
- \*CONTROLS FLEAS AND TICKS ON DOGS FOR 14 DAYS

ACTIVE INGREDIENTS:

Pyrethrin.....	0.15%
*Piperonyl Butoxide, Technical.....	0.75%
*Permethrin.....	0.30%
INERT INGREDIENTS:..... 98.80%	
100.00%	

\*Equivalent to max. 0.6% (butylcarbonyl)(6-propylpiperonyl) ether and 0.15% related compound. \*\*-(3-phenoxyphenyl) methyl (+/-) cis/trans 3-(2,4-dichloroethoxy)2,2-dimethylcyclopropanecarboxylate. Cis trans ratio: max. 35% (+/-) cis and max. 65% (+/-) trans.

PYRETHROID AND PERMANENT- Registered Trademarks of Roux et Uclaf Corporation.  
EPA REG. NO 5969- EPA EST NO 5969-SC-1

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**  
STATEMENT OF PRACTICAL TREATMENT

IF INHALED: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth to mouth. Get medical attention.  
See Side Panel For Use Cautions

NET CONTENTS: \_\_\_\_\_

**PRECAUTIONARY STATEMENTS**  
Hazardous To Humans & Domestic Animals

**CAUTION**  
Hazardal if inhaled. Avoid breathing vapor or spray mist. Remove contaminated clothing and wash before reuse.

Avoid contamination of food and foodstuffs. Remove pets and birds and cover fish aquaria before spraying. Vacuum room after treatment and ventilate before reoccupying. Do not allow children or pets to contact treated areas until surfaces are dry.

**ENVIRONMENTAL HAZARDS**  
This product is toxic to fish. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwater.  
**Physical Or Chemical Hazards**  
Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130 F may cause bursting.

**DIRECTIONS FOR USE**  
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**  
Store in a cool, dry area. Do not transport or store below 32 degrees F. Wrap container in several layers of newspaper and dispose of in trash. Do not incinerate or puncture.

Do not apply directly to food.  
Do not use in food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas where food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where foods may be prepared or held.  
Nonfood areas are areas such as garbage rooms, lavatories, floor drains (to sewers), drains and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after cleaning and bottling).  
Commercial food/feed areas where the product cannot be used include areas for receiving, serving storage, (Dry, Cold, Frozen, Raw) packaging, edible, waste storage, enclosed processing systems. Not for use in USDA Meat and Poultry Plants.  
Do not use in rooms/areas while occupied by patients, the elderly or infirm. Do not apply to classrooms when in use.

**SHAKE WELL BEFORE USING**  
Remove protective cap, hold container upright and spray from a distance of 12 to 15 inches INDOORS.

Chamber Flies, Houseflies, Fruit Flies, Mosquitoes, Gnats, Wasps, Hornets, Bees, Black Flies, Skipper Flies, Blow Flies and Small Flying Moths: Close all doors and windows and

direct spray upward into center of room with a slow sweeping motion. Spray 5-10 seconds for average room, moving toward the door. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Leave room and keep area closed for 15 minutes after spraying. Ventilate room thoroughly before re-entry. Sweep up and discard fallen insects. Repeat treatment as necessary.  
Cockroaches, Waterbugs, Palmate Bugs, Silverfish, Firebrats, Crickets, Scorpions, Spiders, Millipedes, Centipedes, Sowbugs and Pillbugs: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and doors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatments as necessary.  
Ants: Spray trails, nests and point of entry. Spray on ants where possible. Repeat as necessary.

Indian Meal Moths, Angoumois Grain Moths: Close doors and windows, direct spray into all parts of room especially around stored food product containers, pallets and diked areas. Spray 7-10 seconds for each 1000 cubic feet of space, moving toward the door. Repeat application as necessary.

Exposed stages of Confused Flour Beetles, Cigarette Beetles, Drug Store Beetles, Saw-Toothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites in exposed adult and larval stages: Spray into cracks and crevices, processing and handling equipment, around cartons or containers or other areas where these insects tend to congregate. Contact as many insects as possible. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 min. to 1 hour after treatment. Space spray: Spray 20 seconds per 1000 cu. ft. of space, moving toward the door. Leave room and keep area closed for 15 min. following treatment. Repeat application weekly or as necessary. Infested stored food products should be disposed of, or treated using an approved product intended for this purpose.

Carpet Beetles: Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment as necessary.

Bedbugs: Spray mattresses particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Repeat treatment as necessary. Allow all sprayed articles to dry thoroughly before use.

To control Chastee Moths and Carpet Beetles on wool and other proteinaceous fibers. Remove infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storage direction spray into cracks, joints and crevices. After using infested articles may be treated by spraying lightly, keeping nozzle at least three feet from fabric to avoid staining.

Flies in buildings-Space spray: Hold can 36 inches above floor and direct spray toward floor and lower walls at a rate of 10 seconds per 100 sq. ft. making sure that all floor, sofa and chair surfaces are contacted. Leave room and keep area closed for 15 minutes. Open and ventilate before reoccupying. Repeat treatment after 7 days or as needed.

Flies & Ticks, including ticks which may cause Lyme disease (Adult and Larvae): Remove and thoroughly clean or destroy pet bedding. Spray pet sleeping quarters, bed entire floor and floor covering, with special attention to cracks and crevices around baseboards, and windows and door frames. Infested upholstered furniture and mattresses may also be treated with this product. A test application should be made to upholstery or drapery fabrics in an inconspicuous place before use.

**TO CONTROL Flies, Ticks and Lice on dogs.** Starting at the tail, stroke against the lay of the hair spraying the parted fur in short bursts, directly behind the hand to insure penetrating the coat. Dampen the fur up to the head. Repeat application weekly if necessary to control infestations. NOTE: Not recommended for use on puppies less than four weeks old.

**TO KILL Lice and Lice eggs:** Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Hold container upright with nozzle away from you. Depress valve and spray from a distance of 8-10 inches. Spray each square foot for three seconds. Spray only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all sprayed articles to dry thoroughly before use.

**IN ANIMAL QUARTERS AND MILK ROOMS.** For rapid control of House Flies, Mosquitoes, Gnats, Wasps, Hornets, Bees and Small-Flying Moths, close windows and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

**TO TREAT LIVESTOCK:** To protect beef cattle, dairy cattle, goats, sheep, hogs and horses from attacks of Stable Flies, Horse Flies, Ear Flies, Face Flies, House Flies, Horn Flies, Mosquitoes, and Gnats: Spray about 3 seconds on each side being careful, to spray back, withers and forelegs thoroughly. To protect from Face Flies, spray the face and head, but do not spray into eyes. Repeat treatment when flies are troublesome.

**FOR SWINE.** Do not ship animals for slaughter within 5 days of last treatment.

**TO CONTROL Bloodsucking Lice:** Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

**TO CONTROL Poultry Lice:** spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the bird with a fine mist.

**FOR CONTROL OF Bedbugs and Mites in poultry houses,** spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

**OUTDOORS:**  
Point spray nozzle away from face. Hold can on slight downward angle. For best results, spray when air is calm. Allow a few minutes for product to take effect. Spray with wind L breeze is blowing.

**FLYING INSECTS** Flies, Mosquitoes, Gnats, Small Flying Moths, Hornets and Yellow Jackets. Allow fog to drift over area. Direct spray toward bushes, or grass from at least 4-5 feet away. Give special attention to breeding areas such as garbage cans and compost piles. Repeat as necessary.

**Mud Daubers and Wasp Nests.** Spray mist directly on insects and their nests from approximately 2 feet away. Contacted insects will fly away from fog. Applications should be made in late evening when insects are at rest. Spray into hiding and breeding places or nesting as many insects as possible.

**CRAWLING INSECTS** Ants, Cockroaches, including Asian Cockroaches, Crickets, Mole Crickets, Centipedes, and Silverfish. Spray infested areas (top or bottom area, hitting as many insects as possible. Also legs of tables and chairs to repel ants. Spray ant hills for best control.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with all directions given hereon.

**BEST AVAILABLE COPY**