

6959-95

10/04/2001

1/2



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
6959-95

Date of Issuance:
OCT 4 2001

NOTICE OF PESTICIDE:
 x Registration
 Reregistration

Term of Issuance:
Conditional

Name of Pesticide Product:
CESSCO AGRO-5

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

CESSCO Inc.
3609-A River Road
Johns Island, SC 29455

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Change the label by revising the EPA Registration Number to read, "EPA Reg. No.6959-95".
3. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Signature of Approving Official:

Marion J. Johnson, Jr.
Marion J. Johnson, Jr.

Date:

October 4, 2001

JF

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
& DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Cover or remove food before treatment. All food processing surfaces should be covered during treatment or thoroughly washed and cleaned before using. Do not apply when food processing is in operation. Remove pets, birds, and cover fish aquariums before spraying.

STATEMENT OF

PRATICAL TREATMENTS

- If Swallowed: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor.
- If Inhaled: Move victim to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
- If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for further treatment advice.
- If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
- Have the product container or label with you when calling a poison control center or doctor or going for treatment.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130° F.

STORAGE AND DISPOSAL

Storage: Do not store near heat or open flame. Do not store at temperatures above 130°F.

DISPOSAL: DO NOT ATTEMPT TO REFILL CYLINDER. The cylinder is refillable only by Manufacturer. When empty, close valves, replace safety cap, and return cylinder to manufacturer or distributor.

Important note: Buyer assumes all risks of use, storage, or handling of this product not in accordance with directions given herewith.

Cessco

AGRO-5

ACCEPTED
with COMMENTS
In EPA Letter Dated:
OCT 4 2001

Under the Federal Insecticide, Fungicide, and Rodenticide Act, registered under EPA Reg. No. 6959-97

**AGRICULTURAL
INSECTICIDE
AEROSOL**

used only with Cessco's Automatic
Accudose Aerosol Piping System.

**FOR CONTROL OF
FLYING INSECTS**

INGREDIENTS

ACTIVE INGREDIENTS:

Pyrethrins.....	0.50%
*Piperonyl Butoxide, Technical.....	5.00%
**OTHER INGREDIENTS.....	94.50%
TOTAL.....	100.00%
*Equivalent to 4.00% (butylcarbityl)(6-propylpiperonyl) ether and 1.00% related compounds.	
**Contains Petroleum Distillates	

**KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS**

EPA. EST. #8959-SC-1
EPA. REG. #8959-04
MANUFACTURED BY:
Cessco, Inc.
3609-A River Road
Johns Island, SC 29455
1-800-476-3666

NET WEIGHT: _____ Lbs.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Flies, Horseflies, Deer Flies, Black Flies, Mosquitoes, Gnats, Sandflies, Midges, Wasps, and Small Flying Moths in horsebarns, milking parlors, hog parlors, farrowing pens and holding pens for horses, cows or hogs. Calculate total cubic feet to be treated. Spray at a rate of 1 second per 1000 cubic feet per nozzle tip. Treatments can be made when animals are present. To avoid contamination, do not place a nozzle within 36 inches of food, water, or animal. Dosage may have to be doubled for open barns or holding pens. Repeat treatments as necessary. Optimum results are obtained when daily applications are made.

Out of Doors: For temporary protection around pools, decks, and terraces. Vacate the area prior to treatment. Do not use when people or domestic animals are present. Allow area to ventilate 15 minutes before reentry. Spray 10-15 seconds per treatment depending on wind drift. Repeated treatments may be necessary. Nozzle tips should be directed away from ornamental shrubbery, pools, and any surface or object that is susceptible to staining.

Flies, Gnats, Fruit Flies, Wasps, Small Flying Moths, Accessible stages of Yellow Mealworms, Indian Meal Moths (Chocolate Moths), Angoumois Grain Moths, Almond Moths, Foreign Grain Beetles, Cigarette Beetles, Saw-toothed Grain Beetles, Confused Flour Beetles, and Rice Weevils in food processing plants, canneries, food storage warehouses and plants containing non-perishable packaged, bagged raw, or processed food commodities, as well as bulk grains and animal feed. Processed foods include flour, sugar, cereals, packaged cookies and crackers. Raw commodities include bagged or bulk field dried beans and peas, cocoa beans, nuts grains, and sweet potatoes. Calculate total cubic feet to be treated. Close all doors and windows. Spray at a rate of 1 second per 1000 cubic feet per nozzle tip. Do not treat when food processing is in operation. All food processing surfaces should be covered or thoroughly washed and cleaned before using. Do not remain in treated area. Ventilate area when treatment is completed. Sweep up and destroy fallen insects. Repeat treatments as necessary. Optimum results are obtained when daily treatments are made. The insecticide does not penetrate bagged or packaged containers and infested products should be treated by other effective and approved methods.

Flies and other Flying Insects in chicken houses and caged laying houses: Treatment should not be made more often than twice daily. Calculate total cubic feet to be treated. Spray at a rate of 1 second per 1000 cubic feet per nozzle tip. To avoid contamination, do not place a nozzle within 36 inches of food, water, or fowl. Optimum results are obtained when daily treatments are made.

a/e