

S O I L S E R V

Avoid drift of this product to any crop for which this chemical is **not** registered. Avoid drift to any crop for which use **may be** registered but closer to harvest than the recommended time limits for this chemical. Avoid drift to any feed or forage crop.

DYLOX* 5 PELLETS

COMPOSITION

Active Ingredients:
 Dimethyl 2,2,2-Trichloro-1-hydroxyethylphosphonate** 5.00%
 Inert Ingredients 95.00%

* DYLOX — Registered trademark Chemagro Corp.
 ** U.S. Pat. 2,701,225

CAUTIONS

KEEP OUT OF REACH OF CHILDREN

May be harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with the skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling this product. Wash contaminated clothing in soap and water before re-use.

Do not contaminate feed or food.

If poisoning occurs, obtain prompt medical aid. This material will cause cholinesterase depression if handled carelessly.

Rinse equipment and containers and dispose of wastes by burning in non-crop lands away from water supplies.

Containers (empty bags) should be disposed of by burning (keep out of smoke).

Atropine is antidotal.

Directions for Use

USE AT 20 POUNDS PER ACRE ON THE FOLLOWING:

CROP	PEST	DAYS BEFORE HARVEST
Sugar Beets.....	Surface feeding Cutworms	14 days for beets or 28 days if tops are used for feed.
Alfalfa.....	Armyworms	One application per cutting. 14 days
Borders, Vegetables.....	Salt Marsh Caterpillar	15 lbs. per 100 linear feet, around field to be protected. Application should be made to the soil away from food crops and not in contact with food crops.

6973-6
ACCEPTED
 JUN 3 / 1967
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. SUBJECT
 TO ATTACHED COMMENTS.

* DYLOX — Registered trademark Chemagro Corp.
** U.S. Pat. 2,701,225

CAUTIONS

KEEP OUT OF REACH OF CHILDREN

May be harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with the skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling this product. Wash contaminated clothing in soap and water before re-use.

Do not contaminate feed or food.

If poisoning occurs, obtain prompt medical aid. This material will cause cholinesterase depression if handled carelessly.

Rinse equipment and containers and dispose of wastes by burning in non-crop lands away from water supplies.

Containers (empty bags) should be disposed of by burning (keep out of smoke).

Atropine is antidotal.

Directions for Use

USE AT 20 POUNDS PER ACRE ON THE FOLLOWING:


CROP	PEST	DAYS BEFORE HARVEST
Sugar Beets.....	Surface feeding Cutworms	14 days for beets or 28 days if tops are used for feed.
Alfalfa.....	Armyworms	One application per cutting. 14 days
Borders, Vegetables.....	Salt Marsh Caterpillar	15 lbs. per 100 linear feet, around field to be protected. Application should be made to the soil away from food crops and not in contact with food crops.

DO NOT APPLY TO CROPS OTHER THAN AS PROVIDED FOR
ON THE LABEL

NET CONTENTS 50 POUNDS

FORM 124-A - REV. FEB '67

USDA REG NO 6973-6



1427 ABBOTT STREET

SALINAS, CALIFORNIA

Notice of Warranty: Soilserv, Inc., warrants (its liability under any warranty or representation being limited to the replacement of the product named on this label) that said product as sold by it consists of the ingredients specified, any kind, express or implied, including any warranty or representation regarding the

6973-6
ACCEPTED
 mar 31 1967
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. SUBJECT
 TO APPLICABLE COMMENTS.