



# KEEP OUT OF REACH OF CHILDREN

May be fatal or harmful if swallowed. Avoid breathing of dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid storage near feed or food products. NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

AVOID CONTAMINATION OF FOOD, FEED, WATER SUPPLIES, STREAMS AND PONDS.

KEEP PETS OUT OF TREATED AREAS.

BIRDS FEEDING ON TREATED AREAS MAY BE KILLED.

Soilserv, Inc. warrants that its liability is limited to the replacement of the product named on this label and that said product is guaranteed to contain, in the amounts specified, the ingredients specified on the label.

## DIRECTIONS FOR USE

The bait may be used to protect flowers, ornamentals, shrubs, citrus, deciduous fruits and the vegetables named below. Apply on the ground around and adjacent to plants. Broadcast applications may be used when there is a heavy infestation. Repeat applications as necessary. Control will be obtained with dosage from 25 to 100 pounds per acre of bait.

## AVOID CONTAMINATING EDIBLE PARTS OF PLANTS WHEN APPLYING THIS BAIT

CROP	Dosage Limitations to meet Residue Tolerances	Days Between Final Application and Harvest
Strawberries	50 lbs.	6
Asparagus, cabbage, broccoli, cauliflower, lettuce (head), beets (roots)	50 lbs.	3
Lettuce (leaf), spinach, beets (tops)	50 lbs.	14
Beans, carrots, tomatoes, peppers, potatoes	50 lbs.	No time limitations
Cucumbers, melons, squash	25 lbs.	No time limitations
Peas	35 lbs.	No time limitations
Citrus, apples, apricots, cherries, grapes, nectarines, peaches, pears, plums	50-60 lbs.	No time limitations
Ornamentals, borders	50-100 lbs.	No time limitations

Special instructions for improved performance are:


1. Slugs and snails favor dark, damp places. Bait where these pests stay during daylight hours.
2. Be sure area to be baited is damp. In dry weather, water area well before baiting.
3. Apply bait in evening as these pests feed mainly at night.
4. Until under control, bait should be renewed at least every 7-10 days or more frequently if necessary.
5. Since snails and slugs migrate extensively, when invasions from adjacent areas are anticipated, a barrier of the bait will aid in control.

U. S. DAIRY NO. 6975-10  
FORM 160 - JULY 67

6975-10

ACCEPTED  
NOV 28 1967

Net Weight 50 lbs.



1427 ABBOTT STREET

INC.

AND RODENTICIDE ACT.  
NOMIC POISON REGISTER-  
SERIAL NO. 5411 N.A.S.  
SUBJECT: ATTACHED COMMENTS.

CALIFORNIA

Notice of Warranty: Soilserv, Inc. warrants its liability under any warranty or representation being limited to the replacement of the product named on this label that contains defect or which does not contain the ingredients specified.