

SOILSERV DIBROM® 8 E

Avoid drift of this product to any crop for which this chemical is not registered. Avoid drift to any crop for which use may be registered but closer to harvest than the recommended time limits for this chemical. Avoid drift to an feed or forage crop.

DANGER

Corrosive, causes eye and skin damage. May be fatal or harmful if swallowed. Do not get on skin, eyes, or clothing. Use water proof gloves and shoes. Avoid contact with hands, face, or clothing. Avoid breathing spray mist. In case of contact with skin, remove contaminated clothing and flush skin with plenty of water. In case of eye contact, flush with water. Avoid contamination of food and feed. Do not use in areas where food is produced. **POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.**

FIRST AID

If swallowed, do not induce vomiting. Give 1-2 glasses of water. If inhaled, get fresh air. If on skin, wash with plenty of water. If in eyes, flush with plenty of water. Avoid further contact with the chemical.

NOTE TO PHYSICIAN:

Positively toxic to bees. Do not apply to crops where bees are present. Do not apply to crops where bees are present. Do not apply to crops where bees are present.

BEES

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not use on helianthus, ornamental cherries, Liquidambar or chrysanthemums as injury may occur. Do not store diluted spray. Make new dilution for each use. Wash equipment thoroughly after each use.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, ponds, marshes, and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Snail and crab may be killed at application rates recommended on this label. Do not apply where there are important resources. Apply this product only as specified on this label.

STORE IN A COOL, DRY PLACE

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. TIMING AND DOSAGE MAY VARY AS A CONSEQUENCE OF LOCAL WEATHER OR CONDITIONS. WE RECOMMEND REFERENCE TO AGRICULTURAL EXPERIMENT STATIONS ON SPECIFIC USAGE.

SOILSERV DIBROM 8 E is a registered trademark of the manufacturer. It is a powerful insecticide and acaricide. It is effective against a wide range of insects and mites. It is used for the control of insects and mites on a wide variety of crops.

USE DESIGNATED RATE BY GROUND EQUIPMENT AS A THOROUGH COVERAGE SPRAY. May be used as a foliar spray or as a soil drench. Do not use as a soil drench on crops where the soil is to be planted. Do not use as a soil drench on crops where the soil is to be planted. Do not use as a soil drench on crops where the soil is to be planted.

BEANS (Dry or Field), SOYBEANS, LIMABEANS AND PEAS for Processing: Alfalfa Looper, Aphids, Spider Mites, Lygus Leaf Miner, GREENBEANS: Aphids, Spider Mites, Loopers, Nitidulids, Leafhoppers, etc. Do not apply to crops where the soil is to be planted.

BROCCOLI, CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, KALE AND COLLARDS: Imported Cabbageworm, Diamond Back Caterpillars, Aphids, etc. Do not apply to crops where the soil is to be planted.

COTTON: Lygus Leaf Perforator, Fleahoppers, Armyworms, Bollworms, Loopers, etc. Do not apply to crops where the soil is to be planted.

CUCUMBERS, SUMMER SQUASH: Cucumber Beetles, Aphids, Spider Mites, Loopers, Thrips, Armyworms, etc. Do not apply to crops where the soil is to be planted.

EGGPLANT, PEPPERS: Aphids, Spider Mites, Blister Beetles, Flea Beetles, Leaf Miners, etc. Do not apply to crops where the soil is to be planted.

WALNUTS: Aphids, etc. Do not apply to crops where the soil is to be planted.

MELONS (Cantaloupe, Honeydew, Muskmelons), WATERMELONS, PUMPKINS, WINTER SQUASH: Loopers, Armyworms, Leafhoppers, Spider Mites, Leaf Miners, etc. Do not apply to crops where the soil is to be planted.

ORANGES, LEMONS, GRAPEFRUIT, TANGENINES: Aphids, Salt Brown Scale, Crawlers, Mites, Citrus Cutworms, Leaf Roller, etc. Do not apply to crops where the soil is to be planted.

PEACHES (Western States Only), LYGUS: Suppression of Mites, Peach Twig Borer, Oriental Fruit Moth (California Only), etc. Do not apply to crops where the soil is to be planted.

RICE: Grasshoppers, etc. Do not apply to crops where the soil is to be planted.

STRAWBERRIES: Leaf Roller, Spider Mites, Aphids, Omnivorous Leaf Tier, Spittlebugs (Northwest), etc. Do not apply to crops where the soil is to be planted.

SUGAR BEETS: Spider Mites, Leafhoppers, etc. Do not apply to crops where the soil is to be planted.

TOBACCO: Hornworm, etc. Do not apply to crops where the soil is to be planted.

TOMATOES (Field): Fruit Flies (Drosophila), etc. Do not apply to crops where the soil is to be planted.

GREENHOUSE VAPOR TREATMENT OF ROSES, OTHER ORNAMENTAL PLANTS, TOMATOES, CUCUMBERS: Spider Mites, Adult Whiteflies, Aphids, Leafrollers, Mealybugs, etc. Do not apply to crops where the soil is to be planted.

WARNING: Do not use in areas where food is produced. Do not use in areas where food is produced. Do not use in areas where food is produced.

FRUIT FLIES (Drosophila sp.) in and around Food Processing, Cull Piles and Refuse Areas, Cigar Mills and Wineries, etc. Do not use in areas where food is produced.

ADULT MOSQUITO, GNAT, STABLEFLY (Dog Flies), AND OTHER RESIDENTIAL AREAS: CROPS LISTED ON LABEL. Do not use in areas where food is produced.

GRAPES: Drosophila, Leafhoppers, etc. Do not use in areas where food is produced.

HOPS: Aphids, Spider Mites, Corn Earworm, Hop Looper, etc. Do not use in areas where food is produced.

CELERY, SPINACH, CHARD, TURNIP GREENS: Aphids, Salt Marsh Caterpillars, Leaf Miners, etc. Do not use in areas where food is produced.

Seller makes no warranty, except other than indicated on the label material when such use and or

BROM[®] 8 EC

ACTIVE INGREDIENTS:

NALED (1, 2-dibromo 2, 2-dichloroethyl dimethyl phosphate).....	58.00 %
Xylene	22.30 %
INERT INGREDIENTS:	19.70 %

WARNING: Avoid over-treatment and direct application to plants as injury may result. White, Emerald, and Golden Rapture roses, Pink Champagne Chrysanthemums, Green Wandering Jew, Poinsettias and Dutchman's Pipe may be injured by DIBROM vapor treatment.

FRUIT FLIES (Drosophila sp.) in and around Food Processing Plants, Loading Docks, Cull Piles and Refuse Areas, Cider Mills and Wineries: Use 2 to 4 quarts of diluted spray per 100 pails of water. Apply as a coarse spray to walls, floors, doors, windows, refuse and cull piles where flies congregate. Apply every 5 to 7 days, as long as necessary, for maintenance control. Do not apply to food or refuse piles to be fed to livestock. Avoid contamination of feeds, foodstuffs and food processing machinery. Do not apply when plants are in operation or when foods are present or packaged. Do not spray surfaces which will come into contact with foods. Cover food containers during spraying periods.

ADULT MOSQUITO, GNAT, STABLEFLY (Dog Flies), AND HOUSEFLY CONTROL - RESIDENTIAL AREAS, CROPS LISTED ON LABEL, WOODLAND: Make applications during peak of infestation and repeat as necessary. Do not apply to rip areas within 4 days of harvest of any crop. Poisonous to fish and shell fish. Avoid use where they will be endangered. **ADULT MOSQUITOES:** Treat shrubbery and vegetation where mosquitoes may rest. Shrubbery and vegetation around stagnant pools, marshy areas, swamps, intermittently flooded areas and shorelines may be treated. Application of this product to any body of water is prohibited. **AIRCRAFT:** East: Apply 0.8 to 4 fluid ounces per acre diluted with water. West: Apply 1.6 to 4 fluid ounces per acre diluted with water. Apply 2 to 8 quarts of diluted spray per acre. **MIST BLOWER:** Use 3 to 5 gals. per 100 pails water. Calibrate equipment (rate of travel and output) to apply 1 to 0.25 lb. technical Naled per acre.

GRAPES: Drosophila, Leaf Hoppers. Use 2 to 4 quarts per 100 pails water. Apply as a diluted spray per acre depending on size of vine. Do not apply to fruit in ripening stage. **MUSTARD VARIETIES:** Do not use on mustard varieties as fruit in ripening stage. Water spray every 24 hours.

HOPS: Aphids, Spider Mites, Corn Earworm, Hop Looper, Cutworm. Apply 1 to 2 quarts per 100 pails water.

CELERY, SPINACH, CHARD, TURNIP GREENS: Aphids, Looper, Armyworms, Salt Marsh Caterpillars, Leaf Miners. Apply 2 to 4 quarts per 100 pails water. Do not spray on plants in fruiting stage. Do not apply to plants in fruiting stage. Water spray every 24 hours.

Contains 8 pounds technical Naled per gallon equivalent to 7.2 pounds Naled.
U. S. Pat. No. 2,971,882

READ THE LABEL

DANGER: KEEP OUT OF REACH OF CHILDREN

SEE BACK PANEL FOR ADDITIONAL CAUTIONS.

WHEN CONTAINER IS EMPTY, DO NOT RINSE CONTAINER, OR IF CONTAINER IS RINSED - DISPOSAL OF THE RINSINGS SHOULD BE CONTROLLED ACCORDING TO THE ENVIRONMENTAL CAUTIONS. BURY USED CONTAINER IN APPROVED DISPOSAL SITE IN ACCORD WITH 40CFR Part 165.

®DIBROM is a registered TM of Chevron Chemical Co.

MANUFACTURED FOR



EPA Est. No. 6973-CA-1
CA-2

Net Contents
1 Gallon

ACCEPTED

1427

APR 07 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, California State Department of Pesticide Regulation, Salinas, California, has approved this product for sale in California. EPA Reg. No. 6973-14

ABBOTT STREET SALINAS, CALIFORNIA

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions.

EPA REG. NO. 6973-14 AA
FORM 2 - REVISED MAR 1975

DIBROM[®] 8 E C

ACTIVE INGREDIENTS:

NALED (1, 2-dibromo-2, 2-dichloroethyl dimethyl phosphate).....	58.00 %
Xylene	22.30 %
INERT INGREDIENTS:	19.70 %

Spider Mites, Ckeworms, Leaf Use minimum of

WARNING Avoid overtreatment and direct application to plants as injury may result. White Butterfly and Golden Rapture roses, Pink Champagne Chrysanthemums, Green Wandering Jew, Poinsettias and Dutchman's Pipe may be injured by DIBROM vapor treatment.

ea Beetles, Leaf 5 before harvest

FRUIT FLIES (Drosophila sp.) in and around Food Processing Plants, Loading Docks, Cull Piles and Refuse Areas, Cider Mills and Wineries Use 2 ounces in 2 1/2 gals. Water (5 pints to 100 gals.) Apply as a coarse spray to walls, floors, doorways, windows, refuse and cull piles where insects congregate. Apply every 5 to 7 days as long as necessary to maintain control. Do not apply to cull fruit or refuse piles to be fed to livestock. Avoid contamination of feeds, foodstuffs and food processing machinery. Do not apply when plants are in operation or when foods are present or exposed. Do not spray surfaces which will come into contact with foods. Cover food containers during spraying periods.

within 10 days of

PUMPKINS, Leaf Miners. Do not apply to skin melons. Use

ADULT MOSQUITO, GNAT, STABLEFLY (Dog Flies), AND HOUSEFLY CONTROL — **RESIDENTIAL AREAS, CROPS LISTED ON LABEL, WOODLAND.** Make applications during peak of infestation and repeat as necessary. Do not apply to crop areas within 4 days of harvest of any crop. Poisonous to fish and shellfish. Avoid use where they will be endangered. **ADULT MOSQUITOES** — Treat shrubbery and vegetation where mosquitoes may rest. Shrubby and vegetation around stagnant pools, marshy areas, swamps, intermittently flooded areas and shorelines may be treated. Application of this product to any body of water is prohibited. **AIRCRAFT:** East — Apply 0.8 to 4 fluid ozs. per acre diluted with water. West — Apply 1.6 to 4 fluid ozs. per acre diluted with water. Apply 2 to 8 quarts of diluted spray per acre. **MIST BLOWER:** Use 3 to 5 qts. per 100 gals. water. Calibrate equipment (rate of travel and output) to apply 0.1 to 0.25 lb. technical Naled per acre.

ft Brown Scale 100 gals. water days of harvest

GRAPES: Drosophila, Leaf Hoppers — 1/2 to 2/3 pt. per 100 gals. water. 100 to 150 gal. diluted spray per acre depending on size of vine. Use only 1 pt. per 100 gal. water on Mexican varieties. Do not use on Italian varieties as fruit injury may occur. Worker re-entry, 24 hour.

ach Twig Borer. To avoid fruit damage use 90

HOPS: Aphids, Spider Mites, Corn Earworm, Hop Looper, Cutworm — 1 pt. per 100 gals. water.

of insects. Do not

Leaf Tier, Spitt

CELERY, SPINACH, CHARD, TURNIP GREENS: Aphids, Loopers, Armyworms, Salt Marsh Caterpillars, Leaf Miners. Use ground application only with 1 inch of spray. 1 pt. in 10 to 25 gals. water per acre on plants less than 1/2 grown. On larger plants use 1 1/2 pt. in 100 to 150 gals. water per acre.

1 gals. water. Do

diluted spray per signs of insects

3 gals. water per to 7 days before id Greenhouse. 1 pt. in 100 gals. spray per acre

ORNAMENTAL leafhoppers, Aphids are dry. Apply user. Application more of the pipes should be ed houses. Use ose house for a ment. When op Vent late houses to ventilate lastic or rubber to tomatoes or stant and Non necessary. Adult sary. Aphids

Contains 8 pounds technical naled per gallon equivalent to 7.2 pounds naled. U. S. Pat. No. 2,971,892

READ THE LABEL
DANGER: KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS.

WHEN CONTAINER IS EMPTY, DO NOT RINSE CONTAINER, OR IF CONTAINER IS RINSED — DISPOSAL OF THE RINSINGS SHOULD BE CONTROLLED ACCORDING TO THE ENVIRONMENTAL CAUTIONS. BURY USED CONTAINER IN APPROVED DISPOSAL SITE IN ACCORD WITH 40CFR Part 165.

[®]DIBROM is a registered TM of Chevron Chemical Co.

MANUFACTURED FOR



EPA Est. No. 6973-CA-1 ← CA-2

Net Contents 1 Gallon

ACCEPTED
1427
APR 07 1975
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, this pesticide registered under EPA Reg. No. 6973-14

1427 ABBOTT STREET SALINAS, CALIFORNIA

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA REG. NO. 6973 - 14 AA
FORM 2 — REVISED MAR 1975