




# Garden Insect Spray

Kills many outdoor insects which destroy vegetables, roses, flowers, trees and shrubs.



<b>ACTIVE INGREDIENTS:</b>	
*O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	25.00%
Xylene	65.98%
<b>INERT INGREDIENTS</b>	9.02%
<b>TOTAL</b>	<b>100.00%</b>

\*Diazinon® Trademark of Ciba Geigy

**KEEP OUT OF REACH OF CHILDREN**  
**WARNING**  
**STATEMENT OF PRACTICAL TREATMENT**  
 If swallowed drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.  
 If inhaled remove victim to fresh air. Apply respiration if indicated.  
 If on skin wash contaminated skin with soap and water.  
 If in eyes flush eyes with plenty of water for at least 15 minutes. Get medical attention immediately.  
**SEE RIGHT SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS**

**NET CONTENTS: 32 FL. OZ. (1 QUART)**

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS & DOMESTIC ANIMALS**

**WARNING**  
 May be fatal if swallowed. May be absorbed through skin. Avoid breathing spray mist. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and food stuffs. Do not use on household pets or humans. Do not permit children or pets to go onto sprayed grass until the spray has completely dried. Do not use in the home. Do not contaminate ornamental fish ponds. **NOTE TO PHYSICIAN:** Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine sulfate is antidotal and may be administered in conjunction with 2-PAM.

**ENVIRONMENTAL HAZARDS**  
 This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes or estuaries. Snails and crabs may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop in the treatment area. Cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

**PHYSICAL OR CHEMICAL HAZARDS**  
 Do not use, pour, spill, or store near heat or open flame.

**DIRECTIONS FOR USE**  
 This violation of Federal Law to use this product in a manner inconsistent with its labeling.

How to mix and apply for vegetables, measure the area to be sprayed. For all other applications, apply as a thorough cover spray. Make applications when insects first appear. Repeat applications as noted in the rate chart. Use coarse drenching sprays for soil and surface application. Use medium fine mist sprays for trees and plants. Thorough coverage is important. Flush sprayer with clean water after each use.

See rate chart below for complete insect list-use dosages shown unless otherwise indicated.  
 Do not use on certain ferns, pansies, hibiscus, papaya, olea, gardenia and orchids since injury may occur.

**STORAGE AND DISPOSAL**  
**STORAGE:** Do not store near heat or open flames.  
**DISPOSAL:** Do not reuse empty container. Wrap container and put in trash collection.

Neither Germain's, Inc., nor the Seller MAKES ANY REPRESENTATIONS OR WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, WITH RESPECT TO THE CONTENTS HEREOF, WHETHER AS TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ANY OTHER MATTER, EXCEPT AS INDICATED ON THIS LABEL. Neither Germain's, Inc., nor the Seller shall be liable for any injury, damage or loss resulting from the handling, storage or use of this product not in strict accordance with the directions given herein.

SOLD BY **32 GERMAIN'S, Inc.**  
 Since 1871  
 Los Angeles, Ca. 90068

Product of U.S.A.  
 Made in U.S.A.  
 EPA Reg. #6993-57  
 1979 by Germain's, Inc.  
 EPA Est. 6993 CA 1

PROBLEM AREA	PESTS - RATES AND DIRECTIONS
<b>Vegetables*</b>	For Application Thoroughly spray all plant parts. Generally applications at 7 to 10 day intervals may be required. Do not feed treated vines to livestock. For all of the following vegetable crops use 1 1/2 oz. in 3 gals. of water to cover approximately 1,000 sq. ft. of garden.
<b>Beans - Peas</b>	Aphids, Cucumber Beetles, Grasshoppers, Mexican Bean Beetles, Mites
<b>Cabbage, Cauliflower, Collards, Broccoli, Kale, Radishes, Turnips</b>	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbage Worms, Dolerous Leaf Miners
<b>Carrots, Celery, Endive, Lettuce, Parsnip, Spinach</b>	Aphids, Flea Beetles, Leafhoppers, Dolerous Leaf Miners, Mites
<b>Cantaloupes, Cucumbers, Muskmelon, Squash (Summer and Winter), Watermelon</b>	Aphids, Thrips, Mites
<b>Peppers, Potatoes, Tomatoes</b>	Aphids, Colorado Potato Beetles, Flea Beetles, Armyworms (tomato), Flea Beetles, Leafhoppers, Southern Armyworms (tomato and tomato), Vinegar Flies (Drosophila spp. tomato)
<b>Onions</b>	Onion Thrips
<b>Sweet Corn</b>	Aphids, Flea Beetles, Grasshoppers, Mites
*After spraying, do not pick vegetables within: 1 day - Peas, Tomatoes, Sweet Corn 3 days - Cantaloupes, Muskmelons, Winter Squash, Watermelons 5 days - Broccoli, Cauliflower, Peppers 7 days - Beans, Cabbage, Cucumbers, Summer Squash 10 days - Carrots, Celery, Collards, Endive, Kale, Lettuce, Onions, Parsnips, Radishes, Spinach, Turnips 35 days - Potatoes	
<b>Rose &amp; Flowers</b>	Wet the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage. 1 1/2 oz. in 3 gals. of water. Apply as a thorough foliar spray.
<b>Azaleas, Camellias, Roses, Chrysanthemums, Carnations, Gardenia, Lilac, Rhododendron</b>	Aphids, Eucalyptus Scale Crawlers, Flea Beetles, Dolerous Leaf Miners, Leafhoppers, Mealybugs, Mites (Carnation Bud, Carnation Shoot, Cyclamen European Red, Two-spotted Spider), Trips, Whiteflies
<b>Trees &amp; Shrubs</b>	Wet leaves and branches to the dripping point. Try to penetrate dense foliage. 1 1/2 oz. in 3 gals. of water. Apply as thorough foliar spray.
<b>Azalea, Birch, Boxwood, Camellia, Hawthorn, Holly, Locust, Maple, Pine, Spruce, Willow</b>	Aphids, Bagworms, Holly Bud Moths, Leaf Miners, Leafhoppers, Mealybugs, Mites, Leaf Tiers, Scale Crawlers
<b>Fruits &amp; Nuts</b>	Start spraying after blooming. Applications should be made according to infestation or when aphids, mites, and crawler stages of scale insects are first noted. If infestations warrant, repeat applications at 7 to 10 day intervals. 1 1/2 oz. in 3 gals. of water. Apply as a full cover spray.
<b>Apricots, Peaches, Nectarines</b>	Aphids, Apricot Mealybugs (apricot), Mites (Clove (brown), two spotted Spider), Scale Crawlers (olive, San Jose)
<b>Cherries</b>	Black Cherry Aphids, Cherry Fruit Flies, Cherry Fruitworms (Maggots), Fruit Tree Leaf Rollers, Mites (Cherry Rust), Scale Crawlers (San Jose)
<b>Figs</b>	Dried Fruit Beetles, Vinegar (Drosophila spp.)
<b>Plum, Prunes</b>	Aphids (Leaf Curl, Plum Mealy Plum Thistle), Mites (Clove (brown), European Red)
<b>Strawberries</b>	Aphids, Two spotted Mites, Strawberry Leaf Rollers
<b>Grapes</b>	Aphids, Grape Berry Moths, Grape Leafhoppers, Leafhoppers, Scale Crawlers
<b>Grapefruit, Lemons, Oranges</b>	Citrus Thrips, Scale Crawlers (Citrus Snow, Cottony cushion) 1 1/2 oz. in 3 gals. of water. Apply as a full cover spray after blooming and again in the summer. Thorough spray coverage of the trunk, limbs and buds is essential.
*After spraying do not pick fruit within: 1 day - Strawberries, Figs 10 days - Apricots, Cherries, Grapes, Nectarines, Plums, Prunes 21 day - Grapefruit, Lemons, Oranges, Peaches	