

**GERMAIN'S  
IMPROVED  
BROADLEAF  
WEED KILLER  
KILLS**

BEDSTRAW  
BLACK MEDIC  
BUCKHORN  
BURDOCK  
CHICORY  
CHICKWEED  
CLOVER  
DANDELION  
DOCK  
GROUND IVY  
HEAL-ALL  
HENBIT  
KNOTWEED  
LAMBSQUARTER  
LESPEDESA  
MALLOW  
MORNING GLORY  
OXALIS  
PEPPERGRASS  
PIGWEEED  
PLANTAINS  
POISON IVY  
PURSLANE  
RAGWEED  
SHEEP SOREL  
SPEEDWELL  
SPURGE  
WILD CARROT  
WILD GARLIC  
WILD LETTUCE  
WILD ONION  
YARROW

**GERMAIN'S®**

**IMPROVED**

# Broadleaf Weed Killer

For control of Broadleaf Weeds (Dandelions, Clover, Spurge, Chickweed, Plantain and certain others) in grass lawns.



| ACTIVE INGREDIENTS   | BY WT.         |
|--|----------------|
| ***Dimethylamine salt of 2,4-dichlorophenoxyacetic acid  | 12.78%         |
| ***Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid                             | 5.68%          |
| ***Dimethylamine salt of Dicamba (3,5-dichloro-o-anisic acid)                                    | 1.51%          |
| INERT INGREDIENTS  | 79.83%         |
| <b>TOTAL</b>   | <b>100.00%</b> |
| *2,4-dichlorophenoxyacetic acid (2,4-D) equivalent 0.94 pounds per gallon or 10.52%.             |                |
| **2-(2-methyl-4-chlorophenoxy) propionic acid (MCPP) equivalent 0.42 pounds per gallon or 4.75%. |                |
| ***3,5-dichloro-o-anisic acid (Dicamba) equivalent 0.097 pounds per gallon or 1.10%.             |                |
| U.S. PAT. 3,224,186  |                |

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**  
SEE BACK AND RIGHT PANEL FOR ADDITIONAL CAUTIONS.

**NET CONTENTS 16 FL. OZ. (1 PT.)**

**GERMAIN'S  
IMPROVED  
BROADLEAF  
WEED KILLER**

**CAUTION:** Harmful if swallowed. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist.

Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not contaminate domestic or irrigation water. Do not pour spray solutions near these plants. Do not store in contact with fertilizers, seeds, insecticides, or fungicides. Do not spray on carpetgrass, dichondra nor lawns or turf where desirable Bentgrass or Clovers are present. When container is empty, wash thoroughly and destroy. Never re-use. Use only lawn sprayers for this use. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs. This product may be injurious to St. Augustine grass.

**CONTENTS WILL TREAT 5,400 SQUARE FEET**

**GERMAIN'S IMPROVED BROADLEAF WEED KILLER**

Germain's Improved Broadleaf Weed Killer is a selective herbicide providing a broad spectrum of weed-kill. It is especially effective against weeds such as Dandelion, Chickweed, Spotted Spurge, and Oxalis. In addition to these, it kills virtually all broadleaf weeds, some of which are listed on the right side panel.

Improved Broadleaf Weed Killer can be applied to grass lawns containing Bluegrass, Fescues, Bermuda and Zoysia grasses. Some injury can be expected to turf when used on Centipeda or St. Augustine grass lawns. "Spot Spray" weeds only. Clover present will be damaged if sprayed. Do not use on Dichondra and Carpetgrass lawns. Do not allow spray to get on vegetables or ornamental plants since injury may result.

**DIRECTIONS FOR USE**

For best results, apply Germain's Improved Broadleaf Weed Killer in Spring or early Fall when weeds are actively growing. Avoid spraying during long, excessively dry or hot periods unless adequate irrigation is available. Do not irrigate within 24 hours of application. Do not apply to newly seeded lawns until after the second or third mowing. Grass seed can be safely sown 3 to 4 weeks after application at recommended rate. For best absorption of material by the weeds, avoid cutting or mowing lawn for 2-3 days before and after spraying.

**DOSAGE:** Place 1/2 fluid oz. or 1 tablespoonful of Germain's Improved Broadleaf Weed Killer in Germain's Deluxe Lawn Sprayer (15 gallons) or other similar hose-end applicator. Fill with water to the 1 gallon mark and apply over 170 sq. ft. of lawn area. Below are dosage rates for larger areas.

| AMOUNT OF CHEMICAL TO USE | FILL SPRAYER TO | AREA TO SPRAY |
|---------------------------|-----------------|---------------|
| (TABLESPOONFUL)           | (FL. OZ.)       | (GALLONS)     |
| 5                         | 2 1/2           | 5             |
| 10                        | 5               | 10            |
| 15                        | 7 1/2           | 15            |
| 20                        | 10              | 20            |
| 32 or 1 Pint              | 16              | 32            |
|                           |                 | 850 sq. ft.   |
|                           |                 | 1,700 sq. ft. |
|                           |                 | 2,550 sq. ft. |
|                           |                 | 3,400 sq. ft. |
|                           |                 | 5,400 sq. ft. |

When using a pressure (tank type) sprayer, measure 2 fl. oz. or 4 table-spoonfuls of chemical into tank, add 1 gallon of water. This dosage will treat 670 sq. ft. or can be used as a spot treatment.

**NOTE:** Whether using a hose-end sprayer or pressure sprayer, avoid sprays which are too fine because they can drift and damage ornamental or garden plants.

GERMAIN'S, INC., and Seller make no warranty, express or implied, as to merchantability, fitness for a particular purpose or otherwise, concerning the use of this product other than as indicated on the label. Germain's, Inc., and Seller shall not be liable for any injury, damage or loss resulting from the handling, storing or use of this product not in strict accordance with the directions given herein.

**GERMAIN'S, inc.**  
Since 1871  
Los Angeles, California 90058

Product of U.S.A. • Made in U.S.A. • ©1971 by Germain's, Inc.  
Stock No. 20-901-68 • USDA Reg. No. 6993-41

ACCEPTED  
JUN - 8 - 1971  
UNDER NO 6993-41